

A note on the polynomial approximation of vertex singularities in the boundary element method in three dimensions *

Alexei Bespalov †

Abstract

We study polynomial approximations of vertex singularities of the type $r^\lambda |\log r|^\beta$ on three-dimensional surfaces. The analysis focuses on the case when $\lambda > -\frac{1}{2}$. This assumption is a minimum requirement to guarantee that the above singular function is in the energy space for boundary integral equations with hypersingular operators. Thus, the approximation results for such singularities are needed for the error analysis of boundary element methods on piecewise smooth surfaces. Moreover, to our knowledge, the approximation of strong singularities ($-\frac{1}{2} < \lambda \leq 0$) by high-order polynomials is missing in the existing literature. In this note we prove an estimate for the error of polynomial approximation of the above vertex singularities on quasi-uniform meshes discretising a polyhedral surface. The estimate gives an upper bound for the error in terms of the mesh size h and the polynomial degree p .

Key words: p -approximation, hp -approximation on quasi-uniform meshes, boundary element method, singularities

AMS Subject Classification: 41A10, 65N15, 65N38

1 Introduction

In this note we analyse polynomial approximations of vertex singularities inherent to solutions of boundary integral equations (BIE) on a polyhedral surface Γ . In particular, denoting by r the distance to a vertex of Γ , we study approximations of singularities of the type $r^\lambda |\log r|^\beta$ under a minimum assumption on λ ensuring that this singular function is in the space $H^{1/2}(\Gamma)$ (the energy space for the BIE with hypersingular operator on Γ).

It is well known that solutions to BIE on piecewise smooth surfaces exhibit a singular behaviour in neighbourhoods of edges and vertices of the surface. In [16, 17] explicit formulas are given to specify this behaviour for polyhedral and piecewise plane open surfaces. In particular, it has been shown that solutions of BIE can be decomposed into a number of singular functions and a smooth remainder. Moreover, taking enough singularity terms in this decomposition, one

*Supported by EPSRC under grant no. EP/E058094/1.

†BICOM, Department of Mathematical Sciences, Brunel University, Uxbridge, West London UB8 3PH, UK. email: a1bespalov@yahoo.com. Supported by the INTAS Young Scientist Fellowship grant (project no. 06-1000014-5945) and by the Russian Science Support Foundation.

can obtain the smooth remainder as regular as needed. Let r be the distance to a vertex v of Γ and let ρ be the distance to one of the edges $e \subset \partial\Gamma$ such that $\bar{e} \ni v$. Then typical singularities are:

- (i) vertex singularities of the type $r^\lambda |\log r|^{\beta_1}$;
- (ii) edge singularities of the type $\rho^\gamma |\log \rho|^{\beta_2}$;
- (iii) combined edge-vertex singularities of the type $r^{\lambda-\gamma} \rho^\gamma |\log r|^{\beta_3}$;

here, λ and γ are real parameters to be specified below and β_i ($i = 1, \dots, 3$) are non-negative integers.

The admissible values of λ and γ depend on the problem under consideration. Let us consider the following model problem: *Find $u \in H^{1/2}(\Gamma)$ such that*

$$\langle Wu, v \rangle = \langle f, v \rangle \quad \forall v \in H^{1/2}(\Gamma). \quad (1.1)$$

Here, $f \in H^{-1/2}(\Gamma)$ is a given functional, W is the hypersingular operator

$$Wu(x) := -\frac{1}{4\pi} \frac{\partial}{\partial n_x} \int_{\Gamma} u(y) \frac{\partial}{\partial n_y} \frac{1}{|x-y|} dS_y, \quad W : H^{1/2}(\Gamma) \rightarrow H^{-1/2}(\Gamma);$$

$\langle \cdot, \cdot \rangle = \langle \cdot, \cdot \rangle_{L^2(\Gamma)}$ denotes the extension of the $L^2(\Gamma)$ -inner product by duality, and $H^{-1/2}(\Gamma)$ is the dual space of $H^{1/2}(\Gamma)$. The latter space is defined in §2.

As it follows from [17], for *sufficiently smooth* given f , the singularity exponents λ and γ satisfy

$$\lambda \geq \lambda_1 > 0 \quad \text{and} \quad \gamma \geq \gamma_1 > 1/2. \quad (1.2)$$

We note that in the case of an open surface Γ , the energy space for problem (1.1) is $\tilde{H}^{1/2}(\Gamma)$ and for sufficiently smooth given f there hold

$$\lambda \geq \lambda_1 > 0 \quad \text{and} \quad \gamma \geq \gamma_1 \geq 1/2. \quad (1.3)$$

Thus, conditions (1.2) (or (1.3)) appear, if singularities in the solution to problem (1.1) are caused solely by the geometry of the surface. However, for singular right-hand sides f in (1.1), it may also occur that

$$\lambda \geq \lambda_1 > -1/2 \quad \text{and} \quad \gamma \geq \gamma_1 > 0, \quad (1.4)$$

which are the minimum requirements to ensure $u \in H^{1/2}(\Gamma)$.

In the framework of the p -version of the boundary element method (BEM), approximations of singularities (i)–(iii) were first analysed in [15] under assumptions (1.2) on λ , γ . These assumptions guarantee that all singular functions (i)–(iii) are $H^1(\Gamma)$ -regular. Due to this fact, a rigorous analysis of polynomial approximations of these singularities in $L^2(\Gamma)$ and in $H^1(\Gamma)$ was performed in [15]. Then, using interpolation between these spaces, the paper culminated in the optimal a priori error estimate for the p -version of the BEM with hypersingular operator on a (closed) polyhedral surface Γ (for smooth right-hand side f).

Later, in [5], we extended the results of [15] to the case of open surfaces, where singularity exponents λ, γ satisfy (1.3). Moreover, our analysis in [5] for the p -BEM and then in [6] for the hp -BEM with quasi-uniform meshes, covers the least regular cases (λ, γ satisfying (1.4)), but only for edge and vertex-edge singularities (on both open and closed piecewise plane surfaces). In these cases the corresponding singularities are not in $H^1(\Gamma)$ and one cannot apply the results of [15]. To the author's knowledge, the analysis of the high-order polynomial approximations of the least regular vertex singularities is missing in the existing literature. With this note we aim to fill this gap. As in [5, 6], we perform the error analysis on a scale of fractional order Sobolev spaces. However, in contrast to [5], the analysis of p -approximations in this note relies on explicit definitions of corresponding norms by the K-method of interpolation.

Let us note that efficient approximations of singularities (i)–(iii) under minimum assumptions for λ, γ as in (1.4) can be helpful also for the analysis of high order BEM for time-harmonic Maxwell equations in the exterior and/or interior of polyhedral domains. In fact, it is known (see [11]) that solutions to these problems of electromagnetism are vector fields whose components exhibit singularities analogous to those in (i)–(iii). It has been shown recently that the analysis of polynomial approximations of singular vector fields inherent to the solution of the electric field integral equation on a plane open surface $\tilde{\Gamma}$ (in the energy space $\tilde{\mathbf{H}}^{-1/2}(\text{div}, \tilde{\Gamma})$) can be reduced to the analysis of scalar singularities (i)–(iii) in Sobolev spaces $H^{1/2}(\tilde{\Gamma})$ and $\tilde{H}^{1/2}(\tilde{\Gamma})$ (see [9]). Besides efficient approximation of singularities, the use of high order methods for solving problems of electromagnetism can be advantageous also from the point of view of minimising numerical dispersion errors (see [1, 2, 13]).

We also note that for BIE with weakly singular operators, where the energy space is $H^{-1/2}(\Gamma)$ (or $\tilde{H}^{-1/2}(\Gamma)$, if Γ is an open surface), the minimum assumptions for singularity parameters are

$$\lambda \geq \lambda_1 > -3/2 \quad \text{and} \quad \gamma \geq \gamma_1 > -1.$$

Polynomial approximations of singularities (i)–(iii) under these minimum assumptions were studied in [7, 8] in the context of the p -BEM and the hp -BEM with quasi-uniform meshes.

The rest of the paper is organised as follows. In the next section we introduce a quasi-uniform mesh discretising a Lipschitz polyhedral surface, define corresponding sets of piecewise polynomials, recall definitions of Sobolev spaces and norms, and collect several auxiliary results. Section 3 is focused on p -approximations of vertex singularities on a separate element of the fixed size. Then in §4 we prove the main result (Theorem 4.1), which states an error estimate (in terms of the mesh parameter h and polynomial degree p) for the approximation of vertex singularities by piecewise polynomials on quasi-uniform meshes.

2 Preliminaries

Throughout the paper, Γ denotes a Lipschitz polyhedral surface with plane faces and straight edges. In what follows, $h > 0$ and $p \geq 1$ will always specify the mesh parameter and a polynomial degree, respectively. We will denote by C a generic positive constant which does not depend on h or p .

For any domain $\Omega \subset \mathbf{R}^n$ we will denote $\rho_\Omega = \sup\{\text{diam}(B); B \text{ is a ball in } \Omega\}$. By $A \simeq B$ we mean that A is equivalent to B , i.e., there exists a constant $C > 0$ such that $CB \leq A \leq C^{-1}B$ where B and A may depend on a parameter (usually h or p) but C does not.

Let $\mathcal{M} = \{\Delta_h\}$ be a family of meshes $\Delta_h = \{\Gamma_j; j = 1, \dots, J\}$ on Γ , where Γ_j are open triangles or parallelograms such that $\bar{\Gamma} = \cup_{j=1}^J \bar{\Gamma}_j$. For any $\Gamma_j \in \Delta_h$ we will denote $h_j = \text{diam}(\Gamma_j)$ and $\rho_j = \rho_{\Gamma_j}$. Let $h = \max_j h_j$. In this paper we will consider a family \mathcal{M} of quasi-uniform meshes Δ_h on Γ in the sense that there exist positive constants σ_1, σ_2 independent of h such that for any $\Gamma_j \in \Delta_h$ and arbitrary $\Delta_h \in \mathcal{M}$

$$h \leq \sigma_1 h_j, \quad h_j \leq \sigma_2 \rho_j. \quad (2.1)$$

Let $Q = (0, 1)^2$ and $T = \{(x_1, x_2); 0 < x_1 < 1, 0 < x_2 < x_1\}$ be the reference square and triangle, respectively. Then for any $\Gamma_j \in \Delta_h$ one has $\Gamma_j = M_j(K)$, where M_j is an affine mapping with Jacobian $|J_j| \simeq h_j^2$ and $K = Q$ or T as appropriate.

Further, $\mathcal{P}_p(I)$ denotes the set of polynomials of degree $\leq p$ on an interval $I \subset \mathbf{R}$. Moreover, $\mathcal{P}_p^1(T)$ is the set of polynomials on T of total degree $\leq p$, and $\mathcal{P}_p^2(Q)$ is the set of polynomials on Q of degree $\leq p$ in each variable. Let $K \subset \mathbf{R}^2$ be an arbitrary triangle or parallelogram, and let $K = M(T)$ or $K = M(Q)$ with an invertible affine mapping M . Then by $\mathcal{P}_p(K)$ we will denote the set of polynomials v on K such that $v \circ M \in \mathcal{P}_p^1(T)$ if K is a triangle and $v \circ M \in \mathcal{P}_p^2(Q)$ if K is a parallelogram (in particular, we will use this notation for $K = Q$ and $K = T$). For given p , we then consider the space of continuous, piecewise polynomials on the mesh $\Delta_h \in \mathcal{M}$,

$$S^{hp}(\Gamma) := \{v \in C^0(\Gamma); v|_{\Gamma_j} \in \mathcal{P}_p(\Gamma_j), j = 1, \dots, J\}.$$

Let us recall the Sobolev norms and spaces that will be used, see [14, 12]. For a domain $\Omega \subset \mathbf{R}^n$ and integer s let $H^s(\Omega)$ be the closure of $C^\infty(\Omega)$ with respect to the norm

$$\|u\|_{H^s(\Omega)}^2 = \|u\|_{H^{s-1}(\Omega)}^2 + |u|_{H^s(\Omega)}^2 \quad (s \geq 1),$$

where

$$|u|_{H^s(\Omega)}^2 = \int_{\Omega} |D^s u(x)|^2 dx, \quad \text{and} \quad H^0(\Omega) = L_2(\Omega).$$

Here, $|D^s u(x)|^2 = \sum_{|\alpha|=s} |D^\alpha u(x)|^2$ in the usual notation with multi-index $\alpha = (\alpha_1, \dots, \alpha_n)$ and with respect to Cartesian coordinates $x = (x_1, \dots, x_n)$. For non-integer s , the Sobolev spaces are defined by interpolation. We use the real K-method of interpolation (see [14]), where, for two normed spaces A_0 and A_1 , the interpolation space $(A_0, A_1)_{\theta, 2}$ ($0 < \theta < 1$) is equipped with the norm

$$\|a\|_{(A_0, A_1)_{\theta, 2}} := \left(\int_0^\infty t^{-2\theta} \inf_{a=a_0+a_1} (\|a_0\|_{A_0}^2 + t^2 \|a_1\|_{A_1}^2) \frac{dt}{t} \right)^{1/2}.$$

Using this method we define

$$H^s(\Omega) = \left(L_2(\Omega), H^1(\Omega) \right)_{s, 2} \quad (0 < s < 1).$$

Furthermore, if Ω has Lipschitz boundary, we set

$$\tilde{H}^r(\Omega) = \left(L_2(\Omega), H_0^s(\Omega) \right)_{\frac{r}{s}, 2} \quad (1/2 < s \leq 1, 0 < r < s).$$

Here, $H_0^s(\Omega)$ ($0 < s \leq 1$) is the completion of $C_0^\infty(\Omega)$ in $H^s(\Omega)$ and we identify $H_0^1(\Omega)$ and $\tilde{H}^1(\Omega)$. It is well-known that the norms in $H^s(\Omega)$, $H_0^s(\Omega)$ and $\tilde{H}^s(\Omega)$ are equivalent for $0 < s < 1/2$. For $1/2 < s < 1$, only the norms in $H_0^s(\Omega)$ and $\tilde{H}^s(\Omega)$ are equivalent.

For $s \in [-1, 0)$ the spaces are defined by duality:

$$H^s(\Omega) = (\tilde{H}^{-s}(\Omega))', \quad \tilde{H}^s(\Omega) = (H^{-s}(\Omega))'.$$

Now let us collect several technical lemmas. We will need the following scaling result.

Lemma 2.1 *Let K^h and K be two open subsets of \mathbf{R}^n such that $K^h = M(K)$ under an invertible affine mapping M . Let $\text{diam } K^h \simeq \rho_{K^h} \simeq h$ and $\text{diam } K \simeq \rho_K \simeq 1$. If $u \in H^m(K^h)$ with integer $m \geq 0$, then $\hat{u} = u \circ M \in H^m(K)$ and there exists a positive constant C depending on m but not on h or u such that*

$$|\hat{u}|_{H^m(K)} \leq Ch^{m-\frac{n}{2}} |u|_{H^m(K^h)}. \quad (2.2)$$

Analogously for any $\hat{u} \in H^m(K)$ there holds

$$|u|_{H^m(K^h)} \leq Ch^{\frac{n}{2}-m} |\hat{u}|_{H^m(K)}. \quad (2.3)$$

Moreover, if $\hat{u} \in H^s(K)$ with real $s \in [0, m]$, then

$$C_1 h^{\frac{n}{2}} \|\hat{u}\|_{H^s(K)} \leq \|u\|_{H^s(K^h)} \leq C_2 h^{\frac{n}{2}-s} \|\hat{u}\|_{H^s(K)}. \quad (2.4)$$

For the proof of (2.2), (2.3) see [10, Theorem 3.1.2]. Inequalities (2.4) then follow by interpolation (see [3, Lemma 4.3]).

The next theorem states the result on the hp -approximations of smooth functions (for the proof see [6, Proposition 4.1]).

Theorem 2.1 *Let $m > 1$. Then for $u \in H^m(\Gamma)$ there exists $u_{hp} \in S^{hp}(\Gamma)$ such that for $s \in [0, 1]$*

$$\|u - u_{hp}\|_{H^s(\Gamma)} \leq Ch^{\mu-s} p^{-(m-\tilde{s})} \|u\|_{H^m(\Gamma)},$$

where $\mu = \min \{m, p + 1\}$ and

$$\tilde{s} = \begin{cases} 1/2 & \text{if } s \in [0, 1/2), \\ 1/2 + \varepsilon, \varepsilon > 0 & \text{if } s = 1/2, \\ s & \text{if } s \in (1/2, 1]. \end{cases} \quad (2.5)$$

The following two lemmas have been also proved in [6], cf. Lemma 3.4 and Lemma 3.5 therein.

Lemma 2.2 Let K^h be a triangle (respectively, a parallelogram) satisfying the assumptions of Lemma 2.1, and let l^h be a side of K^h with vertices v_1, v_2 . Let $w_{hp} \in \mathcal{P}_p(l^h)$ be such that $w_{hp}(v_1) = w_{hp}(v_2) = 0$, and $\|w_{hp}\|_{L_2(l^h)} \leq f(h, p)$. Then there exists $u_{hp} \in \mathcal{P}_{2p+1}(K^h)$ (respectively, $u_{hp} \in \mathcal{P}_p(K^h)$) such that $u_{hp} = w_{hp}$ on l^h , $u_{hp} = 0$ on $\partial K^h \setminus l^h$, and for $0 \leq s \leq 1$

$$\|u_{hp}\|_{H^s(K^h)} \leq C h^{1/2-s} p^{-1+2s} f(h, p).$$

Lemma 2.3 Let $\Delta_h = \{\Gamma_j\}$ be a quasi-uniform mesh on Γ . Then for $0 < s < 1$

$$\|u\|_{H^s(\Gamma)}^2 \geq \sum_j \|u\|_{H^s(\Gamma_j)}^2 \quad \forall u \in H^s(\Gamma),$$

and for $1/2 < s < 1$ there holds

$$\|u\|_{H^s(\Gamma)}^2 \leq C \sum_j \left(h_j^{-2s} \|u\|_{L_2(\Gamma_j)}^2 + |u|_{H^s(\Gamma_j)}^2 \right) \quad \forall u \in H^s(\Gamma). \quad (2.6)$$

The positive constant C in (2.6) is independent of u and the mesh Δ .

3 p -approximation on a separate element of the fixed size

We start with a model situation on the reference square $Q = (0, 1)^2$. This will lead us to the p -approximation result on a separate element (either a triangle or a parallelogram) of the fixed size.

For the model situation, let $\kappa > 1$ and denote $S_\kappa = \{x \in Q; \kappa^{-1}x_1 < x_2 < \kappa x_1\}$. We consider the following singular function over the square Q :

$$u(r, \theta) = r^\lambda |\log r|^\beta \chi(r) w(\theta), \quad (3.1)$$

where (r, θ) denote local polar coordinates with origin at $(0, 0)$, $\lambda > -1/2$, $\beta \geq 0$ is an integer, $w(\theta)$ is sufficiently smooth, and χ is a C^∞ cut-off function satisfying

$$\chi(r) = 1 \quad \text{for } 0 \leq r \leq \delta/2 \quad \text{and} \quad \chi(r) = 0 \quad \text{for } r \geq \delta. \quad (3.2)$$

Here, $\delta \in (0, 1)$ is small enough. If $\lambda = 0$, we will assume that β is a positive integer, so that the function u has only a logarithmic singularity in this case. Observing that $u \in H^s(S_{\kappa_0})$ for $\kappa_0 > 1$ and for any $s \in [0, \lambda + 1)$, we study polynomial approximations of u . We emphasize again that for $\lambda \in (-1/2, 0]$ the function u is not $H^1(\Gamma)$ -regular, and one cannot apply the results of [4, 15, 5], were λ was assumed to be positive.

Theorem 3.1 Let u be given by (3.1). Then there exists a sequence $u_p \in \mathcal{P}_{p+2}^2(Q)$, $p = 1, 2, \dots$, such that $u_p = 0$ at the origin $(0, 0)$,

$$\|u - u_p\|_{H^s(S_{\kappa_0})} \leq C p^{-2(\lambda+1-s)} (1 + \log p)^\beta, \quad 0 \leq s < \min\{1, \lambda + 1\}, \quad (3.3)$$

and for any straight line $\ell \ni (0, 0)$ there holds

$$\|u - u_p\|_{L_2(\ell \cap \bar{S}_{\kappa_0})} \leq C p^{-2(\lambda+1/2)} (1 + \log p)^\beta. \quad (3.4)$$

Although the results of [4] and [15] cannot be applied directly, we will use the approach developed in these papers (see, in particular, Theorem 5.1 in [4] and Theorem 8.1 in [15]). First, we extend u smoothly from S_{κ_0} to $S_{\kappa} \supset S_{\kappa_0}$. This can be done by multiplying (3.1) by a C^∞ cut-off function $\tilde{\chi}(\theta)$ such that for $\kappa > \kappa_0$

$$\begin{aligned}\tilde{\chi}(\theta) &= 1 & \text{for } \arctan \kappa_0^{-1} \leq \theta \leq \arctan \kappa_0, \\ \tilde{\chi}(\theta) &= 0 & \text{for } \theta \leq \arctan \kappa^{-1} \text{ and } \theta \geq \arctan \kappa.\end{aligned}$$

We will retain the notation u for the extended function. Let

$$\xi(x_1, x_2) = (x_1 - \kappa x_2)(\kappa x_1 - x_2) = r^2 \Phi_1(\theta)$$

and

$$u_0(x_1, x_2) = \frac{u(x_1, x_2)}{\xi(x_1, x_2)} = r^{\lambda-2} |\log r|^\beta \chi(r) \Phi_2(\theta),$$

where $\Phi_2(\theta)$ is smooth. Introducing a cut-off function ω such that

$$\omega \in C^\infty(\mathbf{R}), \quad \omega(z) = 0 \text{ for } z \leq 1, \quad \omega(z) = 1 \text{ for } z \geq 2, \quad (3.5)$$

we define for a small $\Delta \in (0, 1)$

$$\omega^\Delta(r) = \omega\left(\frac{r}{\Delta}\right), \quad \tilde{\omega}^\Delta(r) = 1 - \omega^\Delta(r), \quad r \geq 0.$$

Then we decompose u_0 as

$$u_0(x) = \frac{u(x)}{\xi(x)} = u_0(x) \omega^\Delta(r) + u_0(x) \tilde{\omega}^\Delta(r) =: v_0(x) + w_0(x). \quad (3.6)$$

The function v_0 in (3.6) is smooth and vanishes for $0 \leq r \leq \Delta$. Moreover, for any non-negative integers k and l there exists a positive constant $C(k+l)$ independent of Δ such that for $(x_1, x_2) \in Q$ and for $i = 1, 2$

$$\left| \frac{\partial^{k+l} v_0}{\partial x_1^k \partial x_2^l} \right| \leq C(k+l) \begin{cases} 0 & \text{for } 0 < r < \Delta, \\ x_i^{\lambda-2-k-l} |\log \Delta|^\beta & \text{otherwise.} \end{cases} \quad (3.7)$$

Polynomial approximations of functions satisfying (3.7) (and not necessarily having the explicit form given above) were investigated in [4] when proving Theorem 5.1 therein, and were also studied in [15, Theorem 8.1]. The estimate for the approximation error in the $H^1(S_\kappa)$ -norm immediately follows from [4], while [15] gives also the estimate in the $L_2(S_\kappa)$ -norm and then, by interpolation, in the norm of $H^s(S_\kappa)$ with $0 \leq s \leq 1$. Moreover, [15, Lemma 8.2] estimates the approximation error in the $L_2(\ell \cap S_\kappa)$ -norm, where ℓ is the line $x_1 = \tilde{\kappa} x_2$ ($\kappa_0^{-1} \leq \tilde{\kappa} \leq \kappa_0$). We summarise the mentioned results in the following lemma.

Lemma 3.1 *Let $\Delta = p^{-2}$. If v_0 satisfies (3.7), then there exists a sequence $v_p \in \mathcal{P}_{p+2}^2(Q)$, $p = 1, 2, \dots$, such that $v_p(0, 0) = 0$ and for any $0 \leq s \leq 1$*

$$\|\xi v_0 - v_p\|_{H^s(S_{\kappa_0})} \leq C p^{-2(\lambda+1-s)} (1 + \log p)^\beta.$$

Moreover,

$$\|\xi v_0 - v_p\|_{L_2(\ell \cap \bar{S}_{\kappa_0})} \leq C p^{-2(\lambda+1/2)} (1 + \log p)^\beta,$$

where ℓ denotes the line $x_1 = \tilde{\kappa} x_2$ ($\kappa_0^{-1} \leq \tilde{\kappa} \leq \kappa_0$).

The function w_0 in (3.6) has a small support, $\text{supp } w_0 \subset \bar{K}_\Delta = \{x \in \bar{S}_\kappa; 0 \leq r \leq 2\Delta\}$. In the next lemma we show that the function ξw_0 being approximated by zero leads to the same estimates as in (3.3), (3.4).

Lemma 3.2 *Let $\Delta = p^{-2}$. Then for $0 \leq s < \min\{1, \lambda + 1\}$*

$$\|\xi w_0\|_{H^s(S_{\kappa_0})} \leq C p^{-2(\lambda+1-s)} (1 + \log p)^\beta. \quad (3.8)$$

$$\|\xi w_0\|_{L_2(\ell \cap \bar{S}_{\kappa_0})} \leq C p^{-2(\lambda+1/2)} (1 + \log p)^\beta, \quad (3.9)$$

where ℓ is the same as in Lemma 3.1.

Proof. First, we prove (3.8) for $s = 0$. For sufficiently small $\Delta > 0$ one has (hereafter, $\theta_1 = \arctan \kappa^{-1}$, $\theta_2 = \arctan \kappa$)

$$\begin{aligned} \|\xi w_0\|_{L_2(S_{\kappa_0})}^2 &\leq \|\xi w_0\|_{L_2(S_\kappa)}^2 = \|\xi w_0\|_{L_2(K_\Delta)}^2 \\ &\leq C \int_0^{2\Delta} \int_{\theta_1}^{\theta_2} r^{2\lambda} |\log r|^{2\beta} r \, d\theta dr \leq C \Delta^{2\lambda+2} |\log \Delta|^{2\beta}, \quad \lambda > -1, \end{aligned} \quad (3.10)$$

where $C > 0$ is independent of Δ . Let $0 < s < \min\{1, \lambda + 1\}$. Then

$$\|\xi w_0\|_{H^s(S_{\kappa_0})}^2 = \int_0^\infty t^{-2s} \inf_{\xi w_0 = w_1 + w_2} \left(\|w_1\|_{L_2(S_{\kappa_0})}^2 + t^2 \|w_2\|_{H^1(S_{\kappa_0})}^2 \right) \frac{dt}{t}. \quad (3.11)$$

For any $t \in (0, \Delta)$ we define

$$\omega_t(r) = \omega\left(\frac{r}{t}\right), \quad \tilde{\omega}_t(r) = 1 - \omega_t(r), \quad r \geq 0,$$

where ω is as in (3.5). Then by (3.11) we have

$$\|\xi w_0\|_{H^s(S_{\kappa_0})}^2 \leq \int_0^\Delta t^{-2s-1} \left(\|\xi w_0 \tilde{\omega}_t\|_{L_2(S_{\kappa_0})}^2 + t^2 \|\xi w_0 \omega_t\|_{H^1(S_{\kappa_0})}^2 \right) dt$$

$$+ \int_{\Delta}^{\infty} t^{-2s-1} \|\xi w_0\|_{L_2(S_{\kappa_0})}^2 dt. \quad (3.12)$$

Now we estimate the norms on the right-hand side of (3.12). Since $\tilde{\omega}_t(r) = 0$ for $r \geq 2t$, we obtain similarly to (3.10)

$$\|\xi w_0 \tilde{\omega}_t\|_{L_2(S_{\kappa_0})}^2 \leq C \int_0^{2t} r^{2\lambda+1} |\log r|^{2\beta} dr \leq C t^{2\lambda+2} |\log t|^{2\beta}. \quad (3.13)$$

To estimate the norm $\|\xi w_0 \omega_t\|_{H^1(S_{\kappa_0})}$ we evaluate the derivatives of $\xi w_0 \omega_t$. We will use the following inequalities

$$\begin{aligned} \left| \frac{\partial r}{\partial x_i} \right| &\leq 1, & \left| \frac{\partial \theta}{\partial x_i} \right| &\leq \frac{1}{x_i} \leq C(\kappa) \frac{1}{r}, \quad x \in S_{\kappa}, \quad i = 1, 2; \\ \left| \frac{d\omega^{\Delta}(r)}{dr} \right| &= \left| \frac{d\tilde{\omega}^{\Delta}(r)}{dr} \right| = \begin{cases} 0 & \text{for } 0 < r < \Delta \text{ or } r > 2\Delta, \\ \left| \omega' \left(\frac{r}{\Delta} \right) \right| \frac{1}{\Delta} & \text{for } \Delta \leq r \leq 2\Delta \end{cases} \\ &\leq C r^{-1} \quad \text{for } r > 0, \end{aligned}$$

and a similar estimate for $\left| \frac{d\omega_t(r)}{dr} \right|$. Hence, for any $x \in S_{\kappa}$ we find by simple calculations

$$\begin{aligned} \left| \frac{\partial}{\partial x_i} (\xi w_0 \omega_t) \right| &= \left| \frac{\partial}{\partial x_i} \left(r^{\lambda} |\log r|^{\beta} \chi(r) \tilde{\chi}(\theta) w(\theta) \tilde{\omega}^{\Delta}(r) \omega_t(r) \right) \right| \\ &\leq C \left[\left| \frac{\partial}{\partial x_i} (r^{\lambda} |\log r|^{\beta}) \right| + r^{\lambda} |\log r|^{\beta} \left(\left| \frac{d\chi}{dr} \right| + \left| \frac{d\tilde{\chi}}{d\theta} \right| \frac{1}{r} + \left| \frac{dw}{d\theta} \right| \frac{1}{r} + \left| \frac{d\tilde{\omega}^{\Delta}}{dr} \right| + \left| \frac{d\omega_t}{dr} \right| \right) \right] \\ &\leq C r^{\lambda-1} |\log r|^{\beta}, \quad i = 1, 2. \end{aligned} \quad (3.14)$$

Since $\xi w_0 \omega_t$ vanishes on ∂S_{κ} and outside the domain $K_{\Delta}^1 = \{x \in S_{\kappa}; t < r < 2\Delta\}$, we deduce from (3.14) that

$$\begin{aligned} \|\xi w_0 \omega_t\|_{H^1(S_{\kappa_0})}^2 &\leq C \|\xi w_0 \omega_t\|_{H^1(K_{\Delta}^1)}^2 \leq C \int_t^{2\Delta} \int_{\theta_1}^{\theta_2} r^{2\lambda-2} |\log r|^{2\beta} r d\theta dr \\ &\leq C \int_t^{2\Delta} r^{2\lambda-1} |\log r|^{2\beta} dr \leq C \begin{cases} t^{2\lambda} |\log t|^{2\beta} & \text{if } \lambda < 0, \\ \Delta^{2\lambda} |\log \Delta|^{2\beta} & \text{if } \lambda > 0. \end{cases} \end{aligned} \quad (3.15)$$

If $\lambda = 0$, we introduce a small $\varepsilon \in (0, 2 - 2s)$ and estimate the norm $\|\xi w_0 \omega_t\|_{H^1(S_{\kappa_0})}^2$ as follows

$$\|\xi w_0 \omega_t\|_{H^1(S_{\kappa_0})}^2 \leq C \int_t^{2\Delta} r^{-1} |\log r|^{2\beta} dr \leq C |\log t|^{2\beta} \int_t^{2\Delta} r^{-1-\varepsilon} r^{\varepsilon} dr \leq C \Delta^{\varepsilon} t^{-\varepsilon} |\log t|^{2\beta}. \quad (3.16)$$

Using estimates (3.10), (3.13), (3.15) for the norms on the right-hand side of (3.12) we obtain for $0 < s < \min\{1, \lambda + 1\}$

$$\begin{aligned} \|\xi w_0\|_{H^s(S_{\kappa_0})}^2 &\leq C \int_0^\Delta t^{2\lambda+1-2s} |\log t|^{2\beta} dt + C \Delta^{2\lambda+2} |\log \Delta|^{2\beta} \int_\Delta^\infty t^{-2s-1} dt \\ &\leq C \Delta^{2(\lambda+1-s)} |\log \Delta|^{2\beta} \quad \text{if } -1 < \lambda < 0 \end{aligned} \quad (3.17)$$

and

$$\begin{aligned} \|\xi w_0\|_{H^s(S_{\kappa_0})}^2 &\leq C \int_0^\Delta t^{2\lambda+1-2s} |\log t|^{2\beta} dt + C \Delta^{2\lambda} |\log \Delta|^{2\beta} \int_0^\Delta t^{-2s+1} dt \\ &\quad + C \Delta^{2\lambda+2} |\log \Delta|^{2\beta} \int_\Delta^\infty t^{-2s-1} dt \\ &\leq C \Delta^{2(\lambda+1-s)} |\log \Delta|^{2\beta} \quad \text{if } \lambda > 0. \end{aligned} \quad (3.18)$$

In the case when $\lambda = 0$ we proceed similarly and use (3.16) instead of (3.15). Then recalling that $0 < \varepsilon < 2 - 2s$ we have for $0 < s < 1$

$$\begin{aligned} \|\xi w_0\|_{H^s(S_{\kappa_0})}^2 &\leq C \int_0^\Delta t^{-2s-1} (t^2 + t^{2-\varepsilon} \Delta^\varepsilon) |\log t|^{2\beta} dt + C \Delta^2 |\log \Delta|^{2\beta} \int_\Delta^\infty t^{-2s-1} dt \\ &\leq C \Delta^{2-2s} |\log \Delta|^{2\beta} + C \Delta^\varepsilon \int_0^\Delta t^{-2s+1-\varepsilon} |\log t|^{2\beta} dt \\ &\leq C \Delta^{2(1-s)} |\log \Delta|^{2\beta} \quad \text{if } \lambda = 0. \end{aligned} \quad (3.19)$$

Taking $\Delta = p^{-2}$ and using estimates (3.10), (3.17)–(3.19) we prove (3.8).

Let ℓ be the line $x_1 = \tilde{\kappa} x_2$, where $\kappa_0^{-1} \leq \tilde{\kappa} \leq \kappa_0$. Then, recalling that $\text{supp } w_0 \subset \bar{K}_\Delta$, we find by simple calculations

$$\|\xi w_0\|_{L_2(\ell \cap \bar{S}_{\kappa_0})}^2 \leq C(\lambda, \beta, \tilde{\kappa}) \int_0^{2\Delta(1+\tilde{\kappa}^2)^{-1/2}} z^{2\lambda} |\log z|^{2\beta} dz \leq C \Delta^{2\lambda+1} |\log \Delta|^{2\beta}.$$

Setting $\Delta = p^{-2}$ we obtain (3.9). □

Proof of Theorem 3.1. The desired statement follows from Lemmas 3.1 and 3.2 making use of decomposition (3.6). □

Now we consider an element (triangle or parallelogram) $K \subset \mathbf{R}^2$ of the fixed size (i.e., we assume that $\text{diam } K \simeq \rho_K \simeq 1$).

Theorem 3.2 *Let $K \subset \mathbf{R}^2$ and suppose that $O = (0, 0)$ is a vertex of K . Let u be given by (3.1) on K . Then there exists a sequence $u_p \in \mathcal{P}_p(K)$, $p = 1, 2, \dots$ such that for $0 \leq s < \min\{1, \lambda + 1\}$*

$$\|u - u_p\|_{H^s(K)} \leq C p^{-2(\lambda+1-s)} (1 + \log p)^\beta.$$

Moreover, $u_p(0, 0) = 0$, $u_p = 0$ on the sides $l_i \subset \partial K$, $\bar{l}_i \not\ni O$, and

$$\|u - u_p\|_{L_2(l_k)} \leq C p^{-2(\lambda+1/2)} (1 + \log p)^\beta \quad \text{for each side } l_k \subset \partial K, O \in \bar{l}_k.$$

The proof is based on Theorem 3.1 and repeats exactly the arguments in [15, Theorem 8.2].

4 hp -approximation on quasi-uniform meshes

In this section we prove the result on the approximation of vertex singularities by piecewise polynomials defined on the quasi-uniform mesh Δ_h discretising polyhedral surface Γ . Let us fix a vertex v of Γ . We will consider the vertex singularity u given by (3.1), where (r, θ) now refers to local polar coordinates (with origin at v) on each face of Γ containing v .

Theorem 4.1 *Let u be given by (3.1) with $\lambda > -\frac{1}{2}$ and an integer $\beta \geq 0$. Then there exists $u_{hp} \in S^{hp}(\Gamma)$ with $p \geq \lambda$ such that for $0 \leq s < \min\{1, \lambda + 1\}$*

$$\|u - u_{hp}\|_{H^s(\Gamma)} \leq C h^{\lambda+1-s} p^{-2(\lambda+1-s)} (1 + \log(p/h))^{\beta+\nu}, \quad (4.1)$$

where $\nu = \frac{1}{2}$ if $p = \lambda$, and $\nu = 0$ otherwise.

If $1 \leq p < \lambda$, then there exists $u_{hp} \in S^{hp}(\Gamma)$ satisfying for $s \in [0, 1]$

$$\|u - u_{hp}\|_{H^s(\Gamma)} \leq C h^{p+1-s}. \quad (4.2)$$

Proof. Note that assumption $1 \leq p < \lambda$ implies $\lambda > 1$. This case was considered in [6, Theorem 6.1], where estimate (4.2) was proved.

To prove (4.1) we decompose u as $u = \varphi_1 + \varphi_2$, where

$$\varphi_1 := u\chi(r/h_0), \quad \varphi_2 := u(1 - \chi(r/h_0)), \quad h_0 = (\sigma_1\sigma_2)^{-1}h, \quad (4.3)$$

χ is the cut-off function in (3.1), and σ_1, σ_2 are the same as in (2.1).

The singular function φ_1 has small support, $\text{supp } \varphi_1 \subset \bar{A}_v := \cup\{\bar{\Gamma}_j; v \in \bar{\Gamma}_j\}$. Let $K^h = \Gamma_j \subset A_v$ and let $K \subset \mathbf{R}^2$ be a triangle or parallelogram such that $K^h = M_h(K)$ under the affine mapping $M_h : x_i = h\hat{x}_i$, $i = 1, 2$, $x \in K^h$, $\hat{x} \in K$. Then $O = (0, 0)$ is a vertex of K and for $h < \frac{1}{2}$ we have

$$\hat{\varphi}_1(\hat{x}) = \varphi_1(h\hat{x}_1, h\hat{x}_2) = h^\lambda \hat{r}^\lambda \sum_{k=0}^{\beta} \binom{\beta}{k} |\log h|^k |\log \hat{r}|^{\beta-k} \chi(\sigma_1\sigma_2\hat{r})w(\hat{\theta}),$$

where $\hat{r} = (\hat{x}_1^2 + \hat{x}_2^2)^{1/2}$, $\hat{\theta} = \arctan(\hat{x}_2/\hat{x}_1)$.

Let $\mathcal{A} = \{l_i\}$ contain those sides $l_i \subset \partial K$ for which $O \in \bar{l}_i$, and let \mathcal{B} be the union of the other sides of K . Then applying Theorem 3.2 to each function $\hat{r}^\lambda |\log \hat{r}|^k \chi(\sigma_1 \sigma_2 \hat{r}) w(\hat{\theta})$, $k = 0, \dots, \beta$, we find a polynomial $\hat{\phi} \in \mathcal{P}_p(K)$ such that $\hat{\phi}(0, 0) = 0$, $\hat{\phi} = 0$ on \mathcal{B} ,

$$\|\hat{\varphi}_1 - \hat{\phi}\|_{H^s(K)} \leq C(\beta) h^\lambda p^{-2(\lambda+1-s)} (1 + \log(p/h))^\beta, \quad 0 \leq s < \min\{1, \lambda + 1\}, \quad (4.4)$$

$$\|\hat{\varphi}_1 - \hat{\phi}\|_{L_2(l)} \leq C(\beta) h^\lambda p^{-2(\lambda+1/2)} (1 + \log(p/h))^\beta \quad \text{for every } l \in \mathcal{A}. \quad (4.5)$$

Let us define $\phi_j := \hat{\phi} \circ M_h^{-1}$. Then $\phi_j \in \mathcal{P}_p(\Gamma_j)$, $\phi_j = 0$ at the vertex v and on the sides $l_i^h \in \mathcal{B}_j = M_h(\mathcal{B})$. Furthermore, making use of Lemma 2.1, we obtain by (4.4), (4.5)

$$\|\varphi_1 - \phi_j\|_{H^s(\Gamma_j)} \leq C h^{\lambda+1-s} p^{-2(\lambda+1-s)} (1 + \log(p/h))^\beta, \quad 0 \leq s < \min\{1, \lambda + 1\}, \quad (4.6)$$

$$\|\varphi_1 - \phi_j\|_{L_2(l^h)} \leq C h^{\lambda+1/2} p^{-2(\lambda+1/2)} (1 + \log(p/h))^\beta \quad \text{for every } l^h \in \mathcal{A}_j = M_h(\mathcal{A}). \quad (4.7)$$

Suppose that $\Gamma_i, \Gamma_j \subset A_v$ are two elements having the common edge $l^h = \bar{\Gamma}_i \cap \bar{\Gamma}_j$ (these elements may lie on different faces of Γ). Let $\phi_i \in \mathcal{P}_p(\Gamma_i)$ and $\phi_j \in \mathcal{P}_p(\Gamma_j)$ be the approximations of φ_1 constructed above and satisfying estimates (4.6), (4.7). Then the jump $g = (\phi_j - \phi_i)|_{l^h}$ vanishes at the end points of l^h and

$$\|g\|_{L_2(l^h)} \leq C h^{\lambda+1/2} p^{-2(\lambda+1/2)} (1 + \log(p/h))^\beta.$$

If Γ_i is a parallelogram, we use Lemma 2.2 to find a polynomial $z \in \mathcal{P}_p(\Gamma_i)$ such that

$$z = g \quad \text{on } l^h, \quad z = 0 \quad \text{on } \partial\Gamma_i \setminus l^h, \quad (4.8)$$

and for $0 \leq s \leq 1$

$$\|z\|_{H^s(\Gamma_i)} \leq C h^{\lambda+1-s} p^{-2(\lambda+1-s)} (1 + \log(p/h))^\beta. \quad (4.9)$$

In the case that Γ_i is a triangle, we note that (4.6) and (4.7) also hold for a polynomial ψ_j of degree $\left[\frac{p-1}{2}\right]$ (with different constants C for the upper bounds in (4.6) and (4.7)). Then Lemma 2.2 yields a polynomial $z \in \mathcal{P}_p(\Gamma_i)$ which satisfies (4.8), (4.9) for Γ_i being a triangle.

Setting $\tilde{\phi} = \phi_i + z$ on Γ_i and $\tilde{\phi} = \phi_j$ on Γ_j we find a continuous piecewise polynomial $\tilde{\phi}$ such that the norms $\|\varphi_1 - \tilde{\phi}\|_{H^s(\Gamma_i)}$ and $\|\varphi_1 - \tilde{\phi}\|_{H^s(\Gamma_j)}$ are bounded as in (4.6) for $0 \leq s < \min\{1, \lambda + 1\}$.

Repeating the above procedure we construct a continuous function $\psi_1 \in C^0(\bar{A}_v)$ such that $\psi_1 = 0$ on ∂A_v , $\psi_1 \in \mathcal{P}_p(\Gamma_j)$ for each $\Gamma_j \subset A_v$, and

$$\|\varphi_1 - \psi_1\|_{H^s(\Gamma_j)} \leq C h^{\lambda+1-s} p^{-2(\lambda+1-s)} (1 + \log(p/h))^\beta, \quad 0 \leq s < \min\{1, \lambda + 1\}. \quad (4.10)$$

Now we extend ψ_1 by zero onto $\Gamma \setminus A_v$ (keeping the notation ψ_1 for the extension). Then $\psi_1 \in S^{hp}(\Gamma)$ and there holds for $0 \leq s < \min\{1, \lambda + 1\}$

$$\|\varphi_1 - \psi_1\|_{H^s(\Gamma)} \leq C h^{\lambda+1-s} p^{-2(\lambda+1-s)} (1 + \log(p/h))^\beta. \quad (4.11)$$

In fact, for $s = 0$ estimate (4.11) on Γ immediately follows from inequalities (4.10) on individual elements. If $1/2 < s < \min\{1, \lambda + 1\}$, then we use Lemma 2.3:

$$\begin{aligned} \|\varphi_1 - \psi_1\|_{H^s(\Gamma)}^2 &\leq C \left(h^{-2s} \|\varphi_1 - \psi_1\|_{L_2(\Gamma)}^2 + \sum_{j:\Gamma_j \subset \Gamma} \|\varphi_1 - \psi_1\|_{H^s(\Gamma_j)}^2 \right) \\ &\leq C \left(h^{-2s} \|\varphi_1 - \psi_1\|_{L_2(A_v)}^2 + \sum_{j:\Gamma_j \subset A_v} \|\varphi_1 - \psi_1\|_{H^s(\Gamma_j)}^2 \right), \end{aligned}$$

and (4.11) follows again from (4.10), because the number ν_v of elements in A_v is independent of h ($\nu_v \leq \frac{\omega_v}{\theta_0}$, where ω_v is the total length of the closed piecewise smooth arc cut out in the unit sphere \mathbf{S}^2 by the edges of Γ having v as an endpoint, θ_0 is the minimal angle of elements in the mesh).

Finally, for $0 < s \leq 1/2$, estimate (4.11) follows via interpolation between $H^0(\Gamma)$ and $H^{s'}(\Gamma)$ for some $s' \in (\frac{1}{2}, \min\{1, \lambda + 1\})$.

For the function φ_2 (see (4.3)) one has

$$\varphi_2 = r^\lambda |\log r|^\beta \chi(r) (1 - \chi(r/h_0)) w(\theta) \in H^m(\Gamma),$$

where m depends on the regularity of $w(\theta)$, m is fixed and as large as required. Furthermore,

$$\text{supp } \varphi_2 \subset \bar{R}^h, \quad \text{where } R^h = \{x \in \Gamma; \frac{\delta}{2}h_0 < r(x) < \delta \text{ on each face at the vertex } v\},$$

where δ is the same as in (3.2).

To bound the norm $\|\varphi_2\|_{H^k(\Gamma)}$ we need the following inequalities:

$$\left| \frac{\partial^{l+m} r}{\partial x_1^l \partial x_2^m} \right| \leq C r^{1-l-m}, \quad \left| \frac{\partial^{l+m} \theta}{\partial x_1^l \partial x_2^m} \right| \leq C r^{-l-m}$$

for any integer $l, m \geq 0$, and

$$\begin{aligned} \left| \frac{\partial^l}{\partial r^l} (1 - \chi(r/h_0)) \right| &= \begin{cases} 0 & \text{for } 0 < r < \frac{\delta}{2}h_0 \text{ and } r > \delta h_0, \\ |\chi^{(l)}| h_0^{-l} & \text{for } \frac{\delta}{2}h_0 \leq r \leq \delta h_0 \end{cases} \\ &\leq C r^{-l} \quad \text{for } r > 0 \end{aligned}$$

with any integer $l \geq 1$.

Hence, we find by simple calculations

$$\|\varphi_2\|_{H^k(\Gamma)}^2 \leq C (\log(1/h))^{2\beta} \int_{\delta h_0/2}^{\delta} r^{2(\lambda-k)} r dr, \quad 0 \leq k \leq m. \quad (4.12)$$

Further, due to Theorem 2.1, there exists $\psi_2 \in S^{hp}(\Gamma)$ such that for $s \in [0, 1]$

$$\|\varphi_2 - \psi_2\|_{H^s(\Gamma)} \leq C h^{\mu-s} p^{-(k-\bar{s})} \|\varphi_2\|_{H^k(\Gamma)}, \quad (4.13)$$

where $k \in (1, m]$ is integer, $\mu = \min \{k, p + 1\}$, and \tilde{s} is defined by (2.5).

If $\lambda + 1 \leq k \leq m$, then (4.12) and (4.13) yield

$$\|\varphi_2 - \psi_2\|_{H^s(\Gamma)} \leq Ch^{\mu-s+\lambda-k+1} p^{-(k-\tilde{s})} \log^{\beta+\bar{\nu}}(1/h), \quad s \in [0, 1], \quad (4.14)$$

where $\bar{\nu} = \frac{1}{2}$ if $k = \lambda + 1$, and $\bar{\nu} = 0$ if $k > \lambda + 1$.

If $p > 2\lambda + \frac{3}{2}$, we select an integer k satisfying

$$2\lambda + \frac{5}{2} < k \leq p + 1.$$

Then $\mu = k > \frac{3}{2}$ and $p^{-(k-\tilde{s})} \leq p^{-2(\lambda+1-s)}$ for any $s \in [0, 1]$.

If $\lambda < p \leq 2\lambda + \frac{3}{2}$ (i.e., p is bounded), we choose an integer k such that

$$\max \{1, \lambda + 1\} < k \leq p + 1,$$

and if $p = \lambda$, then we take $k = \lambda + 1 = p + 1$. In both these cases $\mu = k > 1$ and $p^{-(k-\tilde{s})} \leq C(\lambda) p^{-2(\lambda+1-s)}$ for any $s \in [0, 1]$.

Thus, for any $p \geq \lambda$, selecting k as indicated above we find by (4.14)

$$\|\varphi_2 - \psi_2\|_{H^s(\Gamma)} \leq Ch^{\lambda+1-s} p^{-2(\lambda+1-s)} \log^{\beta+\nu}(1/h), \quad s \in [0, 1], \quad (4.15)$$

where $\nu = \frac{1}{2}$ if $p = \lambda$ and $\nu = 0$ otherwise.

Now combination of (4.11) and (4.15) gives (4.1) with $u_{hp} := \psi_1 + \psi_2 \in S^{hp}(\Gamma)$. \square

Remark 4.1 *If Γ is an open piecewise plane surface and the function u in (3.1) vanishes on $\partial\Gamma$, then $u \in \tilde{H}^s(\Gamma)$ for any $0 \leq s < \min \{1, \lambda + 1\}$. In this case the same arguments as in the proof of Theorem 4.1 lead to even stronger result: if $p \geq \lambda$, then there exists $u_{hp} \in S_0^{hp}(\Gamma) := S^{hp}(\Gamma) \cap H_0^1(\Gamma)$ such that for $0 \leq s < \min \{1, \lambda + 1\}$*

$$\|u - u_{hp}\|_{\tilde{H}^s(\Gamma)} \leq Ch^{\lambda+1-s} p^{-2(\lambda+1-s)} (1 + \log(p/h))^{\beta+\nu},$$

where ν is the same as in (4.1); if $1 \leq p < \lambda$, then there exists $u_{hp} \in S_0^{hp}(\Gamma)$ satisfying for $s \in [0, 1]$

$$\|u - u_{hp}\|_{\tilde{H}^s(\Gamma)} \leq Ch^{p+1-s}.$$

References

- [1] M. AINSWORTH, *Discrete dispersion relation for hp-version finite element approximation at high wave number*, SIAM J. Numer. Anal., 42 (2004), pp. 553–575.
- [2] ———, *Dispersive properties of high-order Nédélec/edge element approximations of time-harmonic Maxwell equations*, Phil. Trans. R. Soc. Lond. A, 362 (2004), pp. 471–491.

- [3] M. AINSWORTH, W. MCLEAN, AND T. TRAN, *The conditioning of boundary element equations on locally refined meshes and preconditioning by diagonal scaling*, SIAM J. Numer. Anal., 36 (1999), pp. 1901–1932.
- [4] I. BABUŠKA AND M. SURI, *The optimal convergence rate of the p -version of the finite element method*, SIAM J. Numer. Anal., 24 (1987), pp. 750–776.
- [5] A. BESPALOV AND N. HEUER, *The p -version of the boundary element method for hypersingular operators on piecewise plane open surfaces*, Numer. Math., 100 (2005), pp. 185–209.
- [6] ———, *The hp -version of the boundary element method with quasi-uniform meshes in three dimensions*, Report 06/1, BICOM, Brunel University, UK, 2006.
- [7] ———, *The p -version of the boundary element method for weakly singular operators on piecewise plane open surfaces*, Numer. Math., 106 (2007), pp. 69–97.
- [8] ———, *The hp -version of the boundary element method with quasi-uniform meshes for weakly singular operators on surfaces*, Report 07/3, BICOM, Brunel University, UK, 2007.
- [9] ———, *On the p -version of the boundary element method for the electric field integral equation on plane screens* (in preparation).
- [10] P. G. CIARLET, *The Finite Element Method for Elliptic Problems*, North-Holland, Amsterdam, 1978.
- [11] M. COSTABEL AND M. DAUGE, *Singularities of electromagnetic fields in polyhedral domains*, Arch. Rational Mech. Anal., 151 (2000), pp. 221–276.
- [12] P. GRISVARD, *Elliptic Problems in Nonsmooth Domains*, Pitman Publishing Inc., Boston, 1985.
- [13] N. HEUER AND C. PURER, *The p -version of the FEM for slowly oscillating singularities in one dimension*, Report 06/5, BICOM, Brunel University, UK, 2006.
- [14] J. L. LIONS AND E. MAGENES, *Non-Homogeneous Boundary Value Problems and Applications I*, Springer-Verlag, New York, 1972.
- [15] C. SCHWAB AND M. SURI, *The optimal p -version approximation of singularities on polyhedra in the boundary element method*, SIAM J. Numer. Anal., 33 (1996), pp. 729–759.
- [16] T. VON PETERSDORFF AND E. P. STEPHAN, *Decompositions in edge and corner singularities for the solution of the Dirichlet problem of the Laplacian in a polyhedron*, Math. Nachr., 149 (1990), pp. 71–104.
- [17] ———, *Regularity of mixed boundary value problems in \mathbf{R}^3 and boundary element methods on graded meshes*, Math. Methods Appl. Sci., 12 (1990), pp. 229–249.