



Letter

Measurements of azimuthal anisotropy of nonprompt D^0 mesons in PbPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV

The CMS Collaboration ^{*}

CERN, Geneva, Switzerland



ARTICLE INFO

Editor: M. Doser

Keywords:

CMS
Heavy ions
QPG
Flow
Heavy flavor

ABSTRACT

Measurements of the elliptic (v_2) and triangular (v_3) azimuthal anisotropy coefficients are presented for D^0 mesons produced in b hadron decays (nonprompt D^0 mesons) in lead-lead collisions at $\sqrt{s_{NN}} = 5.02$ TeV. The results are compared with previously published charm meson anisotropies measured using prompt D^0 mesons. The data were collected with the CMS detector in 2018 with an integrated luminosity of 0.58 nb^{-1} . Azimuthal anisotropy is sensitive to the interactions of quarks with the hot and dense medium created in heavy ion collisions. Comparing results for prompt and nonprompt D^0 mesons can assist in understanding the mass dependence of these interactions. The nonprompt results show lower magnitudes of v_2 and v_3 and weaker dependences on the meson transverse momentum and collision centrality than those found for prompt D^0 mesons. The results are in agreement with theoretical predictions that include a mass dependence in the interactions of quarks with the medium.

1. Introduction

In ultra-relativistic heavy ion collisions, cold nuclear matter transforms into a state of strongly coupled matter where quarks and gluons are deconfined from hadrons, called the quark-gluon plasma (QGP) [1–4]. One of the main features of the QGP is the collective motion of its constituents. A signature of this collective motion is the presence of a momentum anisotropy of the particles that emerge from the collision. In the plane perpendicular to the beam axis (transverse plane), this anisotropy results from the conversion to momentum space of the initial transverse spatial anisotropy of the colliding system, a phenomenon known as collective anisotropic flow. To quantify this effect, azimuthal (Fourier) coefficients of order n , $v_n = \langle \cos[n(\phi - \Psi_n)] \rangle$ can be used, where ϕ is the single-particle azimuthal angle and Ψ_n is the first angle where the n th harmonic component has its maximum outgoing particle density [5,6]. The measured v_n values for low-momentum light hadrons can be described by relativistic hydrodynamics [7–9]. The second Fourier coefficient, v_2 (referred to as elliptic flow), arises from the combined effects of the elongated shape and the event-by-event fluctuations of the overlap region of the colliding nuclei [10]. The third coefficient, v_3 (referred to as triangular flow), is predominantly due to the fluctuations.

Because of their large mass, bottom (b) and charm (c) quarks are produced at the earliest stage of the collision [11], but their azimuthal

distributions can be affected by their interaction with the medium as they travel through the QGP [12]. At low transverse momentum (p_T), it is expected that charm quarks thermalize and, therefore, they should follow the motion of the gluons and light flavor quarks [13]. In the high p_T region, the azimuthal asymmetry could be caused by a path length dependence of the parton energy loss [14,15]. Hence, studying the flow coefficients of heavy quarks in heavy ion collisions can provide important inputs for understanding the properties of the QGP. Significant v_2 and v_3 coefficients of charm hadrons in lead-lead (PbPb) collisions have already been measured at the CERN LHC, proving that charm quarks exhibit collective flow [16–22]. An early measurement by the CMS Collaboration of the elliptic flow of J/ψ from b hadron decays was consistent with zero, albeit with large uncertainties [23]. Recently, the ALICE Collaboration measured the v_2 of electrons from b hadron decays [24], and the ATLAS Collaboration measured both v_2 and v_3 of muons from b hadron decays [19]. These results indicate that b hadron elliptic flow is nonzero, and also show a clear mass ordering, with smaller measured values of Fourier coefficients than those found for particles containing lighter flavor quarks. Compared to these J/ψ and lepton measurements, the study of the $b \rightarrow D^0$ channel is a promising opportunity due to the large branching fraction of this decay, approximately 70% [25]. This enables lower p_T coverage, thereby increasing sensitivity to the low- p_T flow of b hadrons. In addition, be-

^{*} E-mail address: cms-publication-committee-chair@cern.ch.

cause of its large mass, the D^0 momentum is more closely correlated with the b quark momentum than is the case for leptons.

In this Letter, the first v_2 and v_3 measurements of D^0 mesons from b hadron decay (nonprompt D^0 mesons) in large collision systems are reported, using PbPb collisions at a center-of-mass energy per nucleon pair of $\sqrt{s_{NN}} = 5.02$ TeV. Results for D^0 mesons in the rapidity range $|y| < 1$ are shown as a function of their p_T , spanning 1–30 GeV/c, and in three classes of PbPb centrality: 0–10%, 10–30%, and 30–50%. These centrality percentages represent the fraction of the total inelastic hadronic cross section, with 0% corresponding to full overlap of the two colliding nuclei. The broad range in p_T enables a comprehensive study of different flow generation mechanisms. The results are compared with azimuthal anisotropy measurements of D^0 mesons that do not come from b hadron decays (prompt D^0 mesons) [18], as well as with theoretical predictions. Tabulated results are provided in the HEP-Data record for this analysis [26].

2. Experimental setup and data sample

The CMS apparatus [27] is a multipurpose, nearly hermetic detector, designed to trigger on [28,29] and identify electrons, muons, photons, and (charged and neutral) hadrons [30–32]. A global algorithm [33] aims to reconstruct all individual particles in an event, combining information provided by the all-silicon inner tracker and by the crystal electromagnetic and brass-scintillator hadron calorimeters, operating inside a 3.8 T superconducting solenoid, with data from the gas-ionization muon detectors embedded in the flux-return yoke outside the solenoid. The forward hadron (HF) calorimeter uses steel as an absorber and quartz fibers as the sensitive material. The two halves of the HF are located 11.2 m from the interaction region, one on each end, and together they provide coverage in the pseudorapidity range $3.0 < |\eta| < 5.2$. The HF calorimeters are subdivided into “towers” with $\Delta\eta \times \Delta\phi = 0.175 \times 0.175$, and energy deposited in a tower is treated as a detected hadron in this analysis. They serve as luminosity monitors and the total energy deposited in both HF calorimeters is used for centrality determination [34]. Events are filtered using a two-tiered trigger system. The first level, composed of custom hardware processors, uses information from the calorimeters and muon detectors [28]. The second level, known as the high-level trigger [29], consists of a farm of processors running a version of the full event reconstruction software.

The data analyzed in this Letter consist of minimum bias (MB) PbPb events corresponding to an integrated luminosity of 0.58 nb^{-1} [35,36]. The MB selection requires signals above readout thresholds in the range of ~ 6 – 12 GeV on both sides of the HF calorimeters [29]. The events are further filtered to remove background events (beam-gas interactions and nonhadronic collisions) by applying the procedure described in Ref. [37]. The final results include only events with at least one vertex associated with two or more tracks (primary vertex) within 15 cm from the nominal interaction point along the beam direction, and at least two towers with energy larger or equal to 4 GeV in each of the HF detectors. Finally, in order to suppress the contamination from events with multiple ion collisions, an event is rejected if the shapes of the clusters in the pixel part of the inner tracker are not compatible with those expected in single PbPb collisions. The total event selection efficiency is $(97 \pm 2)\%$.

Simulated events from Monte Carlo (MC) generators are used to study the kinematics of D^0 mesons in PbPb collisions. The D^0 mesons are generated with PYTHIA 8.212 [38], tune CP5 [39], while their decays are modeled with EVTGEN 1.3 [40]. Both prompt and nonprompt D^0 samples are produced. The decay products of the D^0 mesons are then embedded into MB events generated using HYDJET 1.9 [41], tuned to reproduce the charged particle multiplicity and spectra. The response of the CMS detector to these combined events is simulated with GEANT4 [42].

3. Analysis procedure

Inclusive (both prompt and nonprompt) D^0 and \bar{D}^0 meson candidates are reconstructed via their decay channels: $D^0 \rightarrow \pi^+ + K^-$ and $\bar{D}^0 \rightarrow \pi^- + K^+$. Both D^0 and \bar{D}^0 are referred to as D^0 in this Letter. All tracks used in the analysis have $p_T > 1$ GeV/c and $|\eta| < 1.2$. To ensure the best track quality, only tracks that satisfy high purity [32] criteria are considered. In addition, the relative uncertainty in the track p_T measurement is required to be less than 10%. The number of hits along the track trajectory in the tracker must be greater or equal to 11. The χ^2 of the track fit divided by the total number of degrees of freedom of the fit, and also divided by the total number of tracker layers having a hit along the track path, must be less than 0.18. Candidates are formed by combining pairs of tracks from oppositely-charged particles and requiring an invariant mass (m_{inv}) within a $\pm 200 \text{ MeV}/c^2$ window of the world-average D^0 meson mass of $1865 \text{ MeV}/c^2$ [25]. Since no particle identification is available, both possible particle assignments (pion and kaon) are considered for each pair of selected candidate decay tracks.

Further selection is done by applying a boosted decision tree (BDT) algorithm implemented using the TMVA package [43]. For the BDT training, the signal candidates are taken from the simulated events in which reconstructed D^0 mesons are required to match the generated nonprompt D^0 particles. The background training sample consists of two components combined together: combinatorial background and prompt D^0 production. The ratio of these two components in the background training sample is the same as found in the data. The combinatorial background sample is composed of D^0 candidates reconstructed from data whose mass is three to six standard deviations away from the nominal D^0 meson mass (~ 1.795 – 1.830 and ~ 1.900 – $1.935 \text{ GeV}/c^2$). The prompt D^0 meson component of the background sample consists of D^0 candidates that correspond to simulated prompt D^0 mesons in the MC events. In this way, the training is optimized to favor the D^0 mesons produced in b hadron decays. Training is performed separately for each p_T range and centrality class. The variables related to D^0 mesons used to discriminate the signal from the background are: χ^2 probability for the D^0 vertex fit, the distance between the D^0 and primary vertices and its significance, as well as the angle between the momentum of the D^0 meson candidate and the line connecting the primary and D^0 vertices (pointing angle). In addition, related to the decay products of the D^0 meson candidate, the variables used are: the significance of the distances of closest approach (DCA) to the primary vertex (both along and perpendicular to the beam direction), and the number of hits assigned to a particular track. These variables are chosen by analyzing their importance in the decision tree and the correlation matrix among all variables. The BDT cut is optimized to provide the maximal nonprompt D^0 meson signal significance for each analysis bin. Each D^0 candidate is weighted by a correction factor, defined as the inverse of the product of the acceptance and selection efficiencies as functions of p_T and centrality. This factor is calculated from the MC sample using the ratio of the number of reconstructed D^0 mesons (that satisfy selection criteria) to the total number of generated nonprompt D^0 mesons in a certain kinematic range. Potential discrepancies between data and simulated p_T spectra could lead to an inaccuracy in the correction factor. Therefore, before performing this correction, the p_T spectrum of D^0 mesons in the simulation is weighted to match that in the data. The weighting factor is calculated in each centrality range from the ratio of the uncorrected p_T distribution of nonprompt D^0 mesons, extracted from the invariant mass and DCA template fits to data, and the model distribution, obtained with the same reconstruction and selection criteria applied to data.

The measurement of inclusive D^0 meson v_n coefficients employs the scalar-product (SP) method [44] used in previous CMS publications for prompt D^0 mesons [17,18].




Fig. 1. An example of the fit to the invariant mass spectrum (left panel) and an example of the template fit of the inclusive D^0 meson yields, extracted as a function of DCA (right panel). The former fit is used for determining the total D^0 yields and the latter for determining the fraction of nonprompt D^0 mesons.

In this method, the v_n coefficients of all D^0 candidates ($v_n^{\text{sig+bkg}}$) are determined using

$$v_n \{ \text{SP} \} \equiv \frac{\langle Q_n^{\text{D}^0} Q_{nA}^* \rangle}{\sqrt{\frac{\langle Q_{nA} Q_{nB}^* \rangle \langle Q_{nA} Q_{nC}^* \rangle}{\langle Q_{nB} Q_{nC}^* \rangle}}}, \quad (1)$$

where the Q -vectors are defined as $Q_n \equiv \sum_{j=1}^n w_j e^{in\phi_j}$, the asterisk symbol (*) means complex conjugation, and the subscripts A, B, and C refer to different subevents. In each event, the sum for Q_{nA} and Q_{nB} is over all hadrons detected in HF which are above a threshold energy, while the sum for Q_{nC} includes all reconstructed tracks above a p_T threshold.

The $Q_n^{\text{D}^0}$ signifies the flow vector of a D^0 meson candidate within a particular kinematic range. The weight w_j is a dimensionless quantity and corresponds to the energy deposited in the HF tower in GeV for Q_{nA} and Q_{nB} , or to the track p_T in GeV/c, at azimuthal angle ϕ_j for Q_{nC} , or $w_j = 1$ in the case of D^0 meson candidates. The subevent A (B) uses the negative (positive) side of the HF when the D^0 meson candidate is at positive pseudorapidity, and vice versa. This choice of subevents avoids autocorrelations and results in an η gap of at least three units between the D^0 meson daughters and particles from the underlying subevent, thereby suppressing short-range correlations. Flattening and recentering procedures are applied to the Q -vectors related to HF and the tracker, for removing detector acceptance effects [45,46]. The averages $\langle Q_{nA} Q_{nB}^* \rangle$, $\langle Q_{nA} Q_{nC}^* \rangle$, and $\langle Q_{nB} Q_{nC}^* \rangle$ are found by considering all selected events, while the average $\langle Q_n^{\text{D}^0} Q_{nA}^* \rangle$ includes all D^0 meson candidates in all selected events.

To extract the inclusive D^0 meson flow harmonics, the individual D^0 candidates are divided into bins of their flow vector SPs (i.e., Eq. (1) but not averaged over all D^0 candidates). A separate mass spectrum is generated in each of these SP bins and an invariant mass fit is performed to get the SP distribution of the D^0 signal. The inclusive D^0 v_n values can be obtained by:

$$v_n^{\text{D}^0} = \frac{\sum v_n^i Y_i}{\sum Y_i}, \quad (2)$$

where v_n^i is the center of the i th SP bin, and the Y_i is the D^0 yield in the same bin.

The m_{inv} distributions are fitted with five components, as shown in Fig. 1: the sum of two Gaussian functions with the same mean but different widths for the D^0 signal, $S(m_{\text{inv}})$; an additional Gaussian function to describe the invariant mass shape of those D^0 candi-

dates that were given an incorrect mass assignment when swapping the pion and kaon designations, $SW(m_{\text{inv}})$; two Crystal Ball functions [47] to describe the processes $D^0 \rightarrow \pi^+\pi^-$ ($S(m_{\pi^+\pi^-})$) and $D^0 \rightarrow K^+K^-$ ($S(m_{K^+K^-})$); and a third-order polynomial to model the combinatorial background, $Bkg(m_{\text{inv}})$. The contributions from the processes $D^0 \rightarrow \pi^+\pi^-$ and $D^0 \rightarrow K^+K^-$ are also resulting from using an incorrect K or π assignment. The ratio between the $S(m_{\text{inv}})$ and $SW(m_{\text{inv}})$ function yields is fixed according to values obtained from simulated events. The widths and means of the $S(m_{\text{inv}})$, $SW(m_{\text{inv}})$, and Crystal Ball functions are initialized using results obtained from simulation studies. To account for potential differences in momentum resolution between the model and the data, the widths are allowed to vary during the data fitting process. Specifically, a single scale factor, common to all three functions, is employed to adjust the component of the widths arising from momentum resolution. In addition, the means are also allowed to be different between the simulation and data.

The fit of the invariant mass spectrum gives the yield for inclusive D^0 mesons. For the extraction of the two individual components (prompt and nonprompt), a similar procedure as in Refs. [48,49] is followed. Distributions of the DCA between the D^0 meson trajectory and the primary vertex are fitted with a linear combination of prompt and nonprompt D^0 DCA templates obtained from MC in each bin of p_T , centrality, and v_n^i . The data distribution was obtained by extracting the D^0 signal from fits to the invariant mass distribution in each DCA interval. In generating the simulated templates, the difference of the DCA values with and without detector effects for each D^0 meson is scaled. The same scale factor is chosen for both prompt and nonprompt D^0 mesons, with its value found by minimizing the χ^2 of the combined template fit.

The left panel of Fig. 1 shows an example of a fit to the mass spectrum for D^0 candidates in the p_T interval 6–8 GeV/c for the centrality class 30–50% and the SP range $0.05 < v_2^i < 0.10$. The right panel shows an example of the template fit of the inclusive D^0 meson yields, extracted as a function of DCA in the same D^0 kinematic region, but averaged over all v_n .

4. Systematic uncertainties

Systematic uncertainties are estimated by varying the conditions and procedures of the v_n measurements. The absolute differences between the nominal and the alternative analyses are considered as one root-mean-square deviations. In the fit to m_{inv} , the functional form of the combinatorial background was varied using a second-order polynomial and an exponential function. In the 0–10% centrality bin, the uncertainty decreases with p_T up to 5 GeV/c, remains constant from




Fig. 2. The elliptic, v_2 (upper panels), and the triangular, v_3 (lower panels), flow coefficients of nonprompt and prompt (from Ref. [18]) D^0 mesons as functions of their p_T and in three bins of centrality. The bars and the boxes represent statistical and systematic uncertainties, respectively.

Table 1

Summary of systematic uncertainties in the absolute differences between the nominal and the alternative analyses for v_2 and v_3 .

Uncertainty source	v_2	v_3
Bkg. mass PDF	0.002–0.012	0.001–0.005
D^0 efficiency correction	0.0001–0.0005	0.0001–0.0002
DCA template fit	0.006–0.013	0.002–0.008
BDT selection	0.001	0.006

5 to 10 GeV/c, and increases above 10 GeV/c. In more peripheral collisions, the uncertainty from this source increases monotonically with p_T over the entire p_T range. The mass shape for the signal region is varied by using a triple-Gaussian function and negligible differences are observed. The effect of the efficiency is studied by changing the efficiency correction values. In this alternative approach, the efficiency is based solely on simulation and does not consider any potential differences between the MC and data spectra. The largest effect is observed at low p_T . The systematic uncertainty from the determination of the nonprompt D^0 meson fraction is evaluated by performing the template fit using an alternative variable, the DCA significance (DCA/(DCA uncertainty)). The differences observed between data and simulation for the DCA measurement and uncertainty are correlated, as they come from the momentum resolution of the same particles. Therefore, this effect is partially canceled in the DCA significance, obtained by dividing the DCA by its uncertainty. However, as DCA significance arises from the D^0 reconstruction, the widths of the distributions cannot be scaled to further match the data, in the same way it was done for DCA. The difference in results between the DCA and DCA significance template fits is quoted as a systematic uncertainty. This variation of the template fit gives the largest contribution to the systematic uncertainty, especially at the highest p_T in central collisions. The effect of the BDT selection is estimated from the MC samples by comparing the results with and without applying it, in bins of v_n , and taking the difference as a systematic uncertainty. No p_T dependence is observed for this source. Table 1 summarizes the estimates of the systematic uncertainties for v_2 and v_3 from each source. The total systematic uncertainty is obtained by adding individual uncertainties in quadrature.

5. Results

The nonprompt D^0 meson flow harmonics are presented in Fig. 2 together with values for prompt D^0 mesons from Ref. [18]. The results, averaged in the range $2 < p_T < 10$ GeV/c, of $v_2 = 0.0289 \pm 0.0069$,

0.0477 ± 0.0054 , and 0.060 ± 0.010 in the centrality intervals 0–10%, 10–30%, and 30–50%, respectively, indicate nonzero values of elliptic flow for B mesons. The v_2 coefficient in the case of b hadron daughters has its maximum value at p_T of ~ 5 GeV/c and is significantly lower than in the prompt D^0 meson case. This difference becomes more pronounced going from central to peripheral collisions. These results for the mass ordering of collective flow extend the previously observed relationship between charm and light flavor hadrons, where the flow of charm hadrons has a lower magnitude than that for light flavor particles. Measurements also suggest an increase of v_2 towards peripheral collisions, as in the case of light hadrons. This observation agrees with the paradigm of flow as a consequence of initial space anisotropy [10]. The elliptic flow of nonprompt D^0 mesons is found to be of the same magnitude as what the ATLAS Collaboration observed for nonprompt muons [19].

The nonprompt v_3 coefficient results have large statistical uncertainties so that neither the p_T nor the centrality dependence can be determined. As was the case for v_2 , the triangular flow signal is weaker for nonprompt than for prompt D^0 mesons, but the difference is not as large as that for the elliptic flow. Measurements, averaged over all three centrality intervals, suggest positive v_3 for nonprompt D^0 mesons in the $4 < p_T < 6$ GeV/c range with a significance of 3.0 standard deviations. The ATLAS Collaboration v_3 results for b hadrons decaying into muons with $p_T > 4$ GeV/c were found to be consistent with zero [19].

Fig. 3 shows the measured nonprompt D^0 meson v_n coefficients compared with theoretical calculations that have different modeling of the b quark flow. The PHSD model is a microscopic off-shell transport model based on a Boltzmann transport equation approach, and includes only collisional energy loss [50]. The TAMU model computes the space-time evolution of the heavy-quark phase space distribution in the QGP using the Fokker–Planck equation, implemented via Langevin dynamics, and also has no radiative energy loss [53]. The LGR model uses the Langevin approach with gluon radiation, emphasizing the in-medium diffusion dynamics. This model aims to describe heavy-quark evolutions from low to intermediate p_T range, $p_T < 15$ GeV/c [54,55]. The LBT model is based on a linearized Boltzmann approach coupled to a hydrodynamic background, and employs both collisional and radiative energy loss [51,52]. The CUJET3.0 is essentially a jet energy loss framework based on a nonperturbative QGP medium [56,57]. The LBT and CUJET3.0 approaches are applicable only for the high- p_T range and do not have predictions for lower momenta. The CUJET3.0 model implements hadronization through a parton fragmentation function, while all other models combine coalescence and fragmentation. All models can qualitatively describe the p_T dependence of the data. The LGR model re-




Fig. 3. The elliptic, v_2 (upper panel), and the triangular, v_3 (lower panel), flow coefficients of nonprompt D^0 mesons as functions of their p_T and in three bins of centrality. The bars and the boxes represent statistical and systematic uncertainties, respectively. The colored bands show theoretical predictions [50–57].

produces the measurements for the centrality 30–50% and in the low- and the intermediate- p_T ranges, where b quarks are expected to follow Brownian motion, i.e., small random momentum fluctuations caused by collisions with thermal particles. For the central events, 0–10%, in the range $4 < p_T < 10$ GeV/c, the results from this model lie systematically below the data. In the same p_T range, the PHSD model provides a good description of the data, for centralities 0–10% and 10–30%, but it cannot describe the data at higher centralities. In contrast, this same PHSD model underpredicts the v_2 Fourier coefficients for D^0 mesons that are not the product of b hadron decays [18]. The TAMU model has predictions in the centrality range 20–40%, therefore only an indirect comparison with data is possible, but the predictions describe well the measurement performed in the centrality range 10–30%. At higher p_T , where the anisotropy is driven by the path-length dependence of parton energy loss, the LBT model gives lower predictions than CUJET3.0, but both models match data within uncertainties. The PHSD, LBT, and LGR models account for event-by-event fluctuations of the initial geometry, but PHSD is the only model that has predictions for v_3 coefficients. While precision is limited for both the model and the data, they both reach similar maximum values in all centralities.

6. Summary

In summary, the elliptic (v_2) and triangular (v_3) flow harmonics of D^0 mesons that originate in b hadron decays (nonprompt D^0 mesons) are measured in lead-lead collisions at $\sqrt{s_{NN}} = 5.02$ TeV. The v_2 results show a weak transverse momentum (p_T) dependence and suggest a slight increase for more peripheral collisions. An indication of a nonzero v_3 coefficient is found for nonprompt D^0 mesons with $4 < p_T < 6$ GeV/c. The magnitudes of the flow coefficients are lower for nonprompt D^0 than for prompt D^0 mesons. This magnitude difference is more pronounced in the case of v_2 . The qualitative agreement between the results and theoretical models, which include a mass hierarchy in quark interactions with the quark-gluon plasma, extends our understanding of heavy quark interactions with the medium.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

Release and preservation of data used by the CMS Collaboration as the basis for publications is guided by the CMS policy as stated in “[CMS data preservation, re-use and open access policy](#)”.

Acknowledgements

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centers and personnel of the Worldwide LHC Computing Grid and other centers for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC, the CMS detector, and the supporting computing infrastructure provided by the following funding agencies: BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil); MES and BNSF (Bulgaria); CERN; CAS, MoST, and NSFC (China); Minciencias (Colombia); MSES and CSF (Croatia); RIF (Cyprus); SENESCYT (Ecuador); MoER, ERC PUT and ERDF (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); BMBF, DFG, and HGF (Germany); GSRI (Greece); NKFIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIP and NRF (Republic of Korea); MES (Latvia); LAS (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MES and NSC (Poland); FCT (Portugal); MESTD (Serbia); MCIN/AEI and PCTI (Spain); MoSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); MHESI and NSTDA (Thailand); TUBITAK and TENMAK (Turkey); NASU (Ukraine); STFC (United Kingdom); DOE and NSF (USA).

Individuals have received support from the Marie-Curie program and the European Research Council and Horizon 2020 Grant, contract Nos. 675440, 724704, 752730, 758316, 765710, 824093, 884104, and COST Action CA16108 (European Union); the Leventis Foundation; the Alfred P. Sloan Foundation; the Alexander von Humboldt Foundation; the Belgian Federal Science Policy Office; the Fonds pour la Formation à la Recherche dans l’Industrie et dans l’Agriculture (FRIA-Belgium); the Agentschap voor Innovatie door Wetenschap en Technologie (IWT-Belgium); the F.R.S.-FNRS and FWO (Belgium) under the “Excellence of Science – EOS” – be.h project n. 30820817; the Beijing Municipal Science & Technology Commission, No. Z191100007219010; The Ministry of Education, Youth and Sports (MEYS) of the Czech Republic;

the Hellenic Foundation for Research and Innovation (HFRI), Project Number 2288 (Greece); the Deutsche Forschungsgemeinschaft (DFG), under Germany's Excellence Strategy – EXC 2121 “Quantum Universe” – 390833306, and under project number 400140256 - GRK2497; the Hungarian Academy of Sciences, the New National Excellence Program - ÚNKP, the NKFIH research grants K 124845, K 124850, K 128713, K 128786, K 129058, K 131991, K 133046, K 138136, K 143460, K 143477, 2020-2.2.1-ED-2021-00181, and TKP2021-NKTA-64 (Hungary); the Council of Science and Industrial Research, India; the Latvian Council of Science; the Ministry of Education and Science, project no. 2022/WK/14, and the National Science Center, contracts Opus 2021/41/B/ST2/01369 and 2021/43/B/ST2/01552 (Poland); the Fundação para a Ciência e a Tecnologia, grant CEECIND/01334/2018 (Portugal); the National Priorities Research Program by Qatar National Research Fund; MCIN/AEI/10.13039/501100011033, ERDF “a way of making Europe”, and the Programa Estatal de Fomento de la Investigación Científica y Técnica de Excelencia María de Maeztu, grant MDM-2017-0765 and Programa Severo Ochoa del Principado de Asturias (Spain); the Chulalongkorn Academic into Its 2nd Century Project Advancement Project, and the National Science, Research and Innovation Fund via the Program Management Unit for Human Resources & Institutional Development, Research and Innovation, grant B05F650021 (Thailand); the Kavli Foundation; the Nvidia Corporation; the Super-Micro Corporation; the Welch Foundation, contract C-1845; and the Weston Havens Foundation (USA).

References






















- [1] I. Arsene, et al., BRAHMS, Quark-gluon plasma and color glass condensate at RHIC? The perspective from the BRAHMS experiment, Nucl. Phys. A 757 (2005) 1, <https://doi.org/10.1016/j.nuclphysa.2005.02.130>, arXiv:nucl-ex/0410020.
- [2] B.B. Back, et al., PHOBOS, The PHOBOS perspective on discoveries at RHIC, Nucl. Phys. A 757 (2005) 28, <https://doi.org/10.1016/j.nuclphysa.2005.03.084>, arXiv:nucl-ex/0410022.
- [3] J. Adams, et al., STAR, Experimental and theoretical challenges in the search for the quark-gluon plasma: the STAR collaboration's critical assessment of the evidence from RHIC collisions, Nucl. Phys. A 757 (2005) 102, <https://doi.org/10.1016/j.nuclphysa.2005.03.085>, arXiv:nucl-ex/0501009.
- [4] K. Adcox, et al., PHENIX, Formation of dense partonic matter in relativistic nucleus-nucleus collisions at RHIC: experimental evaluation by the PHENIX collaboration, Nucl. Phys. A 757 (2005) 184, <https://doi.org/10.1016/j.nuclphysa.2005.03.086>, arXiv:nucl-ex/0410003.
- [5] S.A. Voloshin, Y. Zhang, Flow study in relativistic nuclear collisions by Fourier expansion of azimuthal particle distributions, Z. Phys. C 70 (1996) 665, <https://doi.org/10.1007/s002880050141>, arXiv:hep-ph/9407282.
- [6] W. Busza, K. Rajagopal, W. van der Schee, Heavy ion collisions: the big picture, and the big questions, Annu. Rev. Nucl. Part. Sci. 68 (2018) 339, <https://doi.org/10.1146/annurev-nucl-101917-020852>, arXiv:1802.04801.
- [7] J.-Y. Ollitrault, Anisotropy as a signature of transverse collective flow, Phys. Rev. D 46 (1992) 229, <https://doi.org/10.1103/PhysRevD.46.229>.
- [8] U. Heinz, R. Snellings, Collective flow and viscosity in relativistic heavy-ion collisions, Annu. Rev. Nucl. Part. Sci. 63 (2013) 123, <https://doi.org/10.1146/annurev-nucl-102212-170540>, arXiv:1301.2826.
- [9] C. Gale, S. Jeon, B. Schenke, Hydrodynamic modeling of heavy-ion collisions, Int. J. Mod. Phys. A 28 (2013) 1340011, <https://doi.org/10.1142/S0217751X13400113>, arXiv:1301.5893.
- [10] B. Alver, G. Roland, Collision-geometry fluctuations and triangular flow in heavy-ion collisions, Phys. Rev. C 81 (2010) 054905, <https://doi.org/10.1103/PhysRevC.81.054905>, arXiv:1003.0194.
- [11] F. Prino, R. Rapp, Open heavy flavor in QCD matter and in nuclear collisions, J. Phys. G 43 (2016) 093002, <https://doi.org/10.1088/0954-3899/43/9/093002>, arXiv:1603.00529.
- [12] R. Rapp, et al., Extraction of heavy-flavor transport coefficients in QCD matter, Nucl. Phys. A 979 (2018) 21, <https://doi.org/10.1016/j.nuclphysa.2018.09.002>, arXiv:1803.03824.
- [13] H. van Hees, V. Greco, R. Rapp, Heavy-quark probes of the quark-gluon plasma and interpretation of recent data taken at the BNL relativistic heavy ion collider, Phys. Rev. C 73 (2006) 034913, <https://doi.org/10.1103/PhysRevC.73.034913>, arXiv:nucl-th/0508055.
- [14] M. Gyulassy, I. Vitev, X.-N. Wang, High p_T azimuthal asymmetry in noncentral A + A at RHIC, Phys. Rev. Lett. 86 (2001) 2537, <https://doi.org/10.1103/PhysRevLett.86.2537>, arXiv:nucl-th/0012092.
- [15] E.V. Shuryak, Azimuthal asymmetry at large p_T seem to be too large for a pure “jet quenching”, Phys. Rev. C 66 (2002) 027902, <https://doi.org/10.1103/PhysRevC.66.027902>, arXiv:nucl-th/0112042.
- [16] ALICE Collaboration, D-meson azimuthal anisotropy in midcentral Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Rev. Lett. 120 (2018) 102301, <https://doi.org/10.1103/PhysRevLett.120.102301>, arXiv:1707.01005.
- [17] CMS Collaboration, Measurement of prompt D^0 meson azimuthal anisotropy in Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Rev. Lett. 120 (2018) 202301, <https://doi.org/10.1103/PhysRevLett.120.202301>, arXiv:1708.03497.
- [18] CMS Collaboration, Measurement of prompt D^0 and D^0 meson azimuthal anisotropy and search for strong electric fields in PbPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Lett. B 816 (2021) 136253, <https://doi.org/10.1016/j.physletb.2021.136253>, arXiv:2009.12628.
- [19] ATLAS Collaboration, Measurement of azimuthal anisotropy of muons from charm and bottom hadrons in Pb+Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV with the ATLAS detector, Phys. Lett. B 807 (2020) 135595, <https://doi.org/10.1016/j.physletb.2020.135595>, arXiv:2003.03565.
- [20] ALICE Collaboration, J/ψ elliptic and triangular flow in Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, J. High Energy Phys. 10 (2020) 141, [https://doi.org/10.1007/JHEP10\(2020\)141](https://doi.org/10.1007/JHEP10(2020)141), arXiv:2005.14518.
- [21] ALICE Collaboration, Transverse-momentum and event-shape dependence of D-meson flow harmonics in Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Lett. B 813 (2021) 136054, <https://doi.org/10.1016/j.physletb.2020.136054>, arXiv:2005.11131.
- [22] CMS Collaboration, Probing charm quark dynamics via multiparticle correlations in PbPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Rev. Lett. 129 (2022) 022001, <https://doi.org/10.1103/PhysRevLett.129.022001>, arXiv:2112.12236.
- [23] CMS Collaboration, Suppression and azimuthal anisotropy of prompt and non-prompt J/ψ production in PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Eur. Phys. J. C 77 (2017) 252, <https://doi.org/10.1140/epjc/s10052-017-4781-1>, arXiv:1610.00613.
- [24] ALICE Collaboration, Elliptic flow of electrons from beauty-hadron decays in Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Rev. Lett. 126 (2021) 162001, <https://doi.org/10.1103/PhysRevLett.126.162001>, arXiv:2005.11130.
- [25] R.L. Workman, et al., Particle Data Group, Review of particle physics, Prog. Theor. Exp. Phys. 2022 (2022) 083C01, <https://doi.org/10.1093/ptep/ptac097>.
- [26] HEPData record for this analysis, <https://doi.org/10.17182/hepdata.131598.2022>.
- [27] CMS Collaboration, The CMS experiment at the CERN LHC, J. Instrum. 03 (2008) S08004, <https://doi.org/10.1088/1748-0221/3/08/S08004>.
- [28] CMS Collaboration, Performance of the CMS level-1 trigger in proton-proton collisions at $\sqrt{s} = 13$ TeV, J. Instrum. 15 (2020) P10017, <https://doi.org/10.1088/1748-0221/15/10/P10017>, arXiv:2006.10165.
- [29] CMS Collaboration, The CMS trigger system, J. Instrum. 12 (2017) P01020, <https://doi.org/10.1088/1748-0221/12/01/P01020>, arXiv:1609.02366.
- [30] CMS Collaboration, Electron and photon reconstruction and identification with the CMS experiment at the CERN LHC, J. Instrum. 16 (2021) P05014, <https://doi.org/10.1088/1748-0221/16/05/P05014>, arXiv:2012.06888.
- [31] CMS Collaboration, Performance of the CMS muon detector and muon reconstruction with proton-proton collisions at $\sqrt{s} = 13$ TeV, J. Instrum. 13 (2018) P06015, <https://doi.org/10.1088/1748-0221/13/06/P06015>, arXiv:1804.04528.
- [32] CMS Collaboration, Description and performance of track and primary-vertex reconstruction with the CMS tracker, J. Instrum. 9 (2014) P10009, <https://doi.org/10.1088/1748-0221/9/10/P10009>, arXiv:1405.6569.
- [33] CMS Collaboration, Particle-flow reconstruction and global event description with the CMS detector, J. Instrum. 12 (2017) P10003, <https://doi.org/10.1088/1748-0221/12/10/P10003>, arXiv:1706.04965.
- [34] CMS Collaboration, Observation and studies of jet quenching in PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, Phys. Rev. C 84 (2011) 024906, <https://doi.org/10.1103/PhysRevC.84.024906>, arXiv:1102.1957.
- [35] CMS Collaboration, Precision luminosity measurement in proton-proton collisions at $\sqrt{s} = 13$ TeV in 2015 and 2016 at CMS, Eur. Phys. J. C 81 (2021) 800, <https://doi.org/10.1140/epjc/s10052-021-09538-2>, arXiv:2104.01927.
- [36] CMS Collaboration, CMS luminosity measurement using nucleus-nucleus collisions at $\sqrt{s_{NN}} = 5.02$ TeV in 2018, in: CMS Physics Analysis Summary CMS-PAS-LUM-18-001, 2022, <https://cds.cern.ch/record/2809613>.
- [37] CMS Collaboration, Charged-particle nuclear modification factors in PbPb and pPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, J. High Energy Phys. 04 (2017) 039, [https://doi.org/10.1007/JHEP04\(2017\)039](https://doi.org/10.1007/JHEP04(2017)039), arXiv:1611.01664.
- [38] T. Sjöstrand, S. Ask, J.R. Christiansen, R. Corke, N. Desai, P. Ilten, S. Mrenna, S. Prestel, C.O. Rasmussen, P.Z. Skands, An introduction to PYTHIA 8.2, Comput. Phys. Commun. 191 (2015) 159, <https://doi.org/10.1016/j.cpc.2015.01.024>, arXiv:1410.3012.
- [39] CMS Collaboration, Extraction and validation of a new set of CMS PYTHIA 8 tunes from underlying-event measurements, Eur. Phys. J. C 80 (2020) 4, <https://doi.org/10.1140/epjc/s10052-019-7499-4>, arXiv:1903.12179.
- [40] D.J. Lange, The evtgen particle decay simulation package, Nucl. Instrum. Methods A 462 (2001) 152, [https://doi.org/10.1016/S0168-9002\(01\)00089-4](https://doi.org/10.1016/S0168-9002(01)00089-4).
- [41] I.P. Lokhtin, A.M. Snigirev, A model of jet quenching in ultrarelativistic heavy ion collisions and high- p_T hadron spectra at RHIC, Eur. Phys. J. C 45 (2006) 211, <https://doi.org/10.1140/epjc/s2005-02426-3>, arXiv:hep-ph/0506189.
- [42] S. Agostinelli, et al., GEANT4, GEANT4—a simulation toolkit, Nucl. Instrum. Methods A 506 (2003) 250, [https://doi.org/10.1016/S0168-9002\(03\)01368-8](https://doi.org/10.1016/S0168-9002(03)01368-8).

- [43] H. Voss, A. Höcker, J. Stelzer, F. Tegenfeldt, TMVA, the toolkit for multivariate data analysis with ROOT, in: *XIth International Workshop on Advanced Computing and Analysis Techniques in Physics Research (ACAT)*, 2007, p. 40, arXiv:physics/0703039 [PoS(ACAT)040].
- [44] C. Adler, et al., STAR, Elliptic flow from two- and four-particle correlations in Au + Au collisions at $\sqrt{s_{NN}} = 130$ GeV, *Phys. Rev. C* 66 (2002) 034904, <https://doi.org/10.1103/PhysRevC.66.034904>, arXiv:nucl-ex/0206001.
- [45] A.M. Poskanzer, S.A. Voloshin, Methods for analyzing anisotropic flow in relativistic nuclear collisions, *Phys. Rev. C* 58 (1998) 1671, <https://doi.org/10.1103/PhysRevC.58.1671>, arXiv:nucl-ex/9805001.
- [46] C. Alt, et al., NA49, Directed and elliptic flow of charged pions and protons in Pb+Pb collisions at 40A and 158A GeV, *Phys. Rev. C* 68 (2003) 034903, <https://doi.org/10.1103/PhysRevC.68.034903>, arXiv:nucl-ex/0303001.
- [47] M.J. Oreglia, A study of the reactions $\psi' \rightarrow \gamma\gamma\psi$, Ph.D. thesis, Stanford University, 1980, <http://www.slac.stanford.edu/cgi-wrap/getdoc/slac-r-236.pdf>, SLAC Report SLAC-R-236.
- [48] CMS Collaboration, Studies of charm and beauty hadron long-range correlations in pp and pPb collisions at LHC energies, *Phys. Lett. B* 813 (2021) 136036, <https://doi.org/10.1016/j.physletb.2020.136036>, arXiv:2009.07065.
- [49] CMS Collaboration, Studies of beauty suppression via nonprompt D^0 mesons in Pb-Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV, *Phys. Rev. Lett.* 123 (2019) 022001, <https://doi.org/10.1103/PhysRevLett.123.022001>, arXiv:1810.11102.
- [50] T. Song, H. Berrehrah, D. Cabrera, J.M. Torres-Rincon, L. Tolos, W. Cassing, E. Bratkovskaya, Tomography of the quark-gluon plasma by charm quarks, *Phys. Rev. C* 92 (2015) 014910, <https://doi.org/10.1103/PhysRevC.92.014910>, arXiv:1503.03039.
- [51] S. Cao, T. Luo, G.-Y. Qin, X.-N. Wang, Linearized Boltzmann transport model for jet propagation in the quark-gluon plasma: heavy quark evolution, *Phys. Rev. C* 94 (2016) 014909, <https://doi.org/10.1103/PhysRevC.94.014909>, arXiv:1605.06447.
- [52] W.-J. Xing, S. Cao, G.-Y. Qin, H. Xing, Flavor hierarchy of jet quenching in relativistic heavy-ion collisions, *Phys. Lett. B* 805 (2020) 135424, <https://doi.org/10.1016/j.physletb.2020.135424>, arXiv:1906.00413.
- [53] M. He, R.J. Fries, R. Rapp, Heavy flavor at the large hadron collider in a strong coupling approach, *Phys. Lett. B* 735 (2014) 445, <https://doi.org/10.1016/j.physletb.2014.05.050>, arXiv:1401.3817.
- [54] S. Li, C. Wang, W. Renzhuo, J. Liao, Probing the transport properties of quark-gluon plasma via heavy-flavor Boltzmann and Langevin dynamics, *Phys. Rev. C* 99 (2019) 054909, <https://doi.org/10.1103/PhysRevC.99.054909>, arXiv:1901.04600.
- [55] S. Li, J. Liao, Data-driven extraction of heavy quark diffusion in quark-gluon plasma, *Eur. Phys. J. C* 80 (7) (2020) 671, <https://doi.org/10.1140/epjc/s10052-020-8243-9>, arXiv:1912.08965.
- [56] S. Shi, J. Liao, M. Gyulassy, Probing the color structure of the perfect QCD fluids via soft-hard-event-by-event azimuthal correlations, *Chin. Phys. C* 42 (2018) 104104, <https://doi.org/10.1088/1674-1137/42/10/104104>, arXiv:1804.01915.
- [57] S. Shi, J. Liao, M. Gyulassy, Global constraints from RHIC and LHC on transport properties of QCD fluids in CUJET/CIBJET framework, *Chin. Phys. C* 43 (2019) 044101, <https://doi.org/10.1088/1674-1137/43/4/044101>, arXiv:1808.05461.

The CMS Collaboration

A. Tumasyan ¹,







Yerevan Physics Institute, Yerevan, Armenia

W. Adam , J.W. Andrejkovic , T. Bergauer , S. Chatterjee , K. Damanakis , M. Dragicevic ,
A. Escalante Del Valle , P.S. Hussain , M. Jeitler ², N. Krammer , L. Lechner , D. Liko , I. Mikulec ,
P. Paulitsch , F.M. Pitters ², R. Schöfbeck , D. Schwarz , M. Sonawane , S. Templ ,
W. Waltenberger , C.-E. Wulz ²,

















Institut für Hochenergiephysik, Vienna, Austria

M.R. Darwish ³, T. Janssen , T. Kello ⁴, H. Rejeb Sfar , P. Van Mechelen ,












Universiteit Antwerpen, Antwerpen, Belgium

E.S. Bols , J. D'Hondt , A. De Moor , M. Delcourt , H. El Faham , S. Lowette , S. Moortgat ,
A. Morton , D. Müller , A.R. Sahasransu , S. Tavernier , W. Van Doninck , D. Vannerom ,

















Vrije Universiteit Brussel, Brussel, Belgium

B. Clerbaux , G. De Lentdecker , L. Favart , D. Hohov , J. Jaramillo , K. Lee ,
M. Mahdavihorrani , I. Makarenko , A. Malara , S. Paredes , L. Pétré , N. Postiau , L. Thomas ,
M. Vanden Bemden , C. Vander Velde , P. Vanlaer ,

Université Libre de Bruxelles, Bruxelles, Belgium

D. Dobur , J. Knolle , L. Lambrecht , G. Mestdach , C. Rendón , A. Samalan , K. Skovpen , M. Tytgat ,
N. Van Den Bossche , B. Vermassen , L. Wezenbeek ,

Ghent University, Ghent, Belgium

A. Benecke , G. Bruno , F. Bury , C. Caputo , P. David , C. Delaere , I.S. Donertas ,
A. Giammanco , K. Jaffel , Sa. Jain , V. Lemaitre , K. Mondal , A. Taliencio , T.T. Tran , P. Vischia ,
S. Wertz ,

Université Catholique de Louvain, Louvain-la-Neuve, Belgium

G.A. Alves , E. Coelho , C. Hensel , A. Moraes , P. Rebello Teles ,

Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil

W.L. Aldá Júnior ¹, M. Alves Gallo Pereira ¹, M. Barroso Ferreira Filho ¹, H. Brandao Malbouisson ¹,
W. Carvalho ¹, J. Chinellato ⁵, E.M. Da Costa ¹, G.G. Da Silveira ^{1,6}, D. De Jesus Damiao ¹,
V. Dos Santos Sousa ¹, S. Fonseca De Souza ¹, J. Martins ^{1,7}, C. Mora Herrera ¹, K. Mota Amarilo ¹,
L. Mundim ¹, H. Nogima ¹, A. Santoro ¹, S.M. Silva Do Amaral ¹, A. Sznajder ¹, M. Thiel ¹,
F. Torres Da Silva De Araujo ^{1,8}, A. Vilela Pereira ¹,

Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

C.A. Bernardes ^{1,6}, L. Calligaris ¹, T.R. Fernandez Perez Tomei ¹, E.M. Gregores ¹, P.G. Mercadante ¹,
S.F. Novaes ¹, Sandra S. Padula ¹,

Universidade Estadual Paulista, Universidade Federal do ABC, São Paulo, Brazil

A. Aleksandrov ¹, G. Antchev ¹, R. Hadjiiska ¹, P. Iaydjiev ¹, M. Misheva ¹, M. Rodozov, M. Shopova ¹,
G. Sultanov ¹,

Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria

A. Dimitrov ¹, T. Ivanov ¹, L. Litov ¹, B. Pavlov ¹, P. Petkov ¹, A. Petrov ¹, E. Shumka ¹,

University of Sofia, Sofia, Bulgaria

S. Thakur ¹,

Instituto De Alta Investigación, Universidad de Tarapacá, Casilla 7 D, Arica, Chile

T. Cheng ¹, T. Javaid ^{1,9}, M. Mittal ¹, L. Yuan ¹,

Beihang University, Beijing, China

M. Ahmad ¹, G. Bauer ¹⁰, Z. Hu ¹, S. Lezki ¹, K. Yi ^{1,10,11},

Department of Physics, Tsinghua University, Beijing, China

G.M. Chen ^{1,9}, H.S. Chen ^{1,9}, M. Chen ^{1,9}, F. Iemmi ¹, C.H. Jiang, A. Kapoor ¹, H. Liao ¹, Z.-A. Liu ^{1,12},
V. Milosevic ¹, F. Monti ¹, R. Sharma ¹, J. Tao ¹, J. Thomas-Wilsker ¹, J. Wang ¹, H. Zhang ¹, J. Zhao ¹,

Institute of High Energy Physics, Beijing, China

A. Agapitos ¹, Y. An ¹, Y. Ban ¹, A. Levin ¹, C. Li ¹, Q. Li ¹, X. Lyu, Y. Mao, S.J. Qian ¹, X. Sun ¹,
D. Wang ¹, J. Xiao ¹, H. Yang,

State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

M. Lu ¹, Z. You ¹,

Sun Yat-Sen University, Guangzhou, China

N. Lu ¹,

University of Science and Technology of China, Hefei, China

X. Gao ^{1,4}, D. Leggat, H. Okawa ¹, Y. Zhang ¹,

Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) - Fudan University, Shanghai, China

Z. Lin ¹, C. Lu ¹, M. Xiao ¹,

Zhejiang University, Hangzhou, Zhejiang, China

C. Avila ¹, D.A. Barbosa Trujillo, A. Cabrera ¹, C. Florez ¹, J. Fraga ¹,

Universidad de Los Andes, Bogota, Colombia

J. Mejia Guisao ^{ID}, F. Ramirez ^{ID}, M. Rodriguez ^{ID}, J.D. Ruiz Alvarez ^{ID},

Universidad de Antioquia, Medellin, Colombia

D. Giljanovic ^{ID}, N. Godinovic ^{ID}, D. Lelas ^{ID}, I. Puljak ^{ID},

University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia

Z. Antunovic, M. Kovac ^{ID}, T. Sculac ^{ID},

University of Split, Faculty of Science, Split, Croatia

V. Brigljevic ^{ID}, B.K. Chitroda ^{ID}, D. Ferencek ^{ID}, S. Mishra ^{ID}, M. Roguljic ^{ID}, A. Starodumov ^{ID,13}, T. Susa ^{ID},

Institute Rudjer Boskovic, Zagreb, Croatia

A. Attikis ^{ID}, K. Christoforou ^{ID}, M. Kolosova ^{ID}, S. Konstantinou ^{ID}, J. Mousa ^{ID}, C. Nicolaou, F. Ptochos ^{ID},
P.A. Razis ^{ID}, H. Rykaczewski, H. Saka ^{ID}, A. Stepennov ^{ID},

University of Cyprus, Nicosia, Cyprus

M. Finger ^{ID}, M. Finger Jr. ^{ID}, A. Kveton ^{ID},

Charles University, Prague, Czech Republic

E. Ayala ^{ID},

Escuela Politecnica Nacional, Quito, Ecuador

E. Carrera Jarrin ^{ID},

Universidad San Francisco de Quito, Quito, Ecuador

A.A. Abdelalim ^{ID,14,15}, E. Salama ^{ID,16,17},

Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt

M. Abdullah Al-Mashad ^{ID}, M.A. Mahmoud ^{ID},

Center for High Energy Physics (CHEP-FU), Fayoum University, El-Fayoum, Egypt

S. Bhowmik ^{ID}, R.K. Dewanjee ^{ID}, K. Ehataht ^{ID}, M. Kadastik, T. Lange ^{ID}, S. Nandan ^{ID}, C. Nielsen ^{ID}, J. Pata ^{ID},
M. Raidal ^{ID}, L. Tani ^{ID}, C. Veelken ^{ID},

National Institute of Chemical Physics and Biophysics, Tallinn, Estonia

P. Eerola ^{ID}, H. Kirschenmann ^{ID}, K. Osterberg ^{ID}, M. Voutilainen ^{ID},

Department of Physics, University of Helsinki, Helsinki, Finland

S. Bharthuar ^{ID}, E. Brücken ^{ID}, F. Garcia ^{ID}, J. Havukainen ^{ID}, M.S. Kim ^{ID}, R. Kinnunen, T. Lampén ^{ID},
K. Lassila-Perini ^{ID}, S. Lehti ^{ID}, T. Lindén ^{ID}, M. Lotti, L. Martikainen ^{ID}, M. Myllymäki ^{ID}, J. Ott ^{ID},
M.m. Rantanen ^{ID}, H. Siikonen ^{ID}, E. Tuominen ^{ID}, J. Tuominiemi ^{ID},

Helsinki Institute of Physics, Helsinki, Finland

P. Luukka ^{ID}, H. Petrow ^{ID}, T. Tuuva,

Lappeenranta-Lahti University of Technology, Lappeenranta, Finland

C. Amendola ^{ID}, M. Besancon ^{ID}, F. Couderc ^{ID}, M. Dejardin ^{ID}, D. Denegri, J.L. Faure, F. Ferri ^{ID}, S. Ganjour ^{ID},
P. Gras ^{ID}, G. Hamel de Monchenault ^{ID}, V. Lohezic ^{ID}, J. Malcles ^{ID}, J. Rander, A. Rosowsky ^{ID}, M.Ö. Sahin ^{ID},
A. Savoy-Navarro ^{ID,18}, P. Simkina ^{ID}, M. Titov ^{ID},

IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France

C. Baldenegro Barrera^{ID}, F. Beaudette^{ID}, A. Buchot Perraguin^{ID}, P. Busson^{ID}, A. Cappati^{ID}, C. Charlot^{ID},
 F. Damas^{ID}, O. Davignon^{ID}, B. Diab^{ID}, G. Falmagne^{ID}, B.A. Fontana Santos Alves^{ID}, S. Ghosh^{ID},
 R. Granier de Cassagnac^{ID}, A. Hakimi^{ID}, B. Harikrishnan^{ID}, G. Liu^{ID}, J. Motta^{ID}, M. Nguyen^{ID},
 C. Ochando^{ID}, L. Portales^{ID}, R. Salerno^{ID}, U. Sarkar^{ID}, J.B. Sauvan^{ID}, Y. Sirois^{ID}, A. Tarabini^{ID},
 E. Vernazza^{ID}, A. Zabi^{ID}, A. Zghiche^{ID},

Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France

J.-L. Agram^{ID,19}, J. Andrea^{ID}, D. Apparú^{ID}, D. Bloch^{ID}, G. Bourgatte^{ID}, J.-M. Brom^{ID}, E.C. Chabert^{ID},
 C. Collard^{ID}, D. Darej, U. Goerlach^{ID}, C. Grimault, A.-C. Le Bihan^{ID}, P. Van Hove^{ID},

Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France

S. Beauceron^{ID}, B. Blancon^{ID}, G. Boudoul^{ID}, A. Carle, N. Chanon^{ID}, J. Choi^{ID}, D. Contardo^{ID}, P. Depasse^{ID},
 C. Dozen^{ID,20}, H. El Mamouni, J. Fay^{ID}, S. Gascon^{ID}, M. Gouzevitch^{ID}, G. Grenier^{ID}, B. Ille^{ID}, I.B. Laktineh,
 M. Lethuillier^{ID}, L. Mirabito, S. Perries, L. Torterotot^{ID}, M. Vander Donckt^{ID}, P. Verdier^{ID}, S. Viret,

Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France

I. Lomidze^{ID}, T. Toriashvili^{ID,21}, Z. Tsamalaidze^{ID,13},

Georgian Technical University, Tbilisi, Georgia

V. Botta^{ID}, L. Feld^{ID}, K. Klein^{ID}, M. Lipinski^{ID}, D. Meuser^{ID}, A. Pauls^{ID}, N. Röwert^{ID}, M. Teroerde^{ID},

RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany

S. Diekmann^{ID}, A. Dodonova^{ID}, N. Eich^{ID}, D. Eliseev^{ID}, M. Erdmann^{ID}, P. Fackeldey^{ID}, D. Fasanella^{ID},
 B. Fischer^{ID}, T. Hebbeker^{ID}, K. Hoepfner^{ID}, F. Ivone^{ID}, M.y. Lee^{ID}, L. Mastrolorenzo, M. Merschmeyer^{ID},
 A. Meyer^{ID}, S. Mondal^{ID}, S. Mukherjee^{ID}, D. Noll^{ID}, A. Novak^{ID}, F. Nowotny, A. Pozdnyakov^{ID}, Y. Rath,
 W. Redjeb^{ID}, H. Reithler^{ID}, A. Schmidt^{ID}, S.C. Schuler, A. Sharma^{ID}, A. Stein^{ID}, L. Vigilante,
 S. Wiedenbeck^{ID}, S. Zaleski,

RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

C. Dziwok^{ID}, G. Flügge^{ID}, W. Haj Ahmad^{ID,22}, O. Hlushchenko, T. Kress^{ID}, A. Nowack^{ID}, O. Pooth^{ID},
 A. Stahl^{ID}, T. Ziemons^{ID}, A. Zotz^{ID},

RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany

H. Aarup Petersen^{ID}, M. Aldaya Martin^{ID}, P. Asmuss, S. Baxter^{ID}, M. Bayatmakou^{ID}, O. Behnke^{ID},
 A. Bermúdez Martínez^{ID}, S. Bhattacharya^{ID}, A.A. Bin Anuar^{ID}, F. Blekman^{ID,23}, K. Borrás^{ID,24}, D. Brunner^{ID},
 A. Campbell^{ID}, A. Cardini^{ID}, C. Cheng, F. Colombina^{ID}, S. Consuegra Rodríguez^{ID}, G. Correia Silva^{ID},
 M. De Silva^{ID}, L. Didukh^{ID}, G. Eckerlin, D. Eckstein^{ID}, L.I. Estevez Banos^{ID}, O. Filatov^{ID}, E. Gallo^{ID,23},
 A. Geiser^{ID}, A. Giraldi^{ID}, G. Greau, A. Grohsjean^{ID}, V. Guglielmi^{ID}, M. Guthoff^{ID}, A. Jafari^{ID,25},
 N.Z. Jomhari^{ID}, B. Kaeck^{ID}, M. Kasemann^{ID}, H. Kaveh^{ID}, C. Kleinwort^{ID}, R. Kogler^{ID}, M. Komm^{ID},
 D. Krücker^{ID}, W. Lange, D. Leyva Pernia^{ID}, K. Lipka^{ID,26}, W. Lohmann^{ID,27}, R. Mankel^{ID},
 I.-A. Melzer-Pellmann^{ID}, M. Mendizabal Morentin^{ID}, J. Metwally, A.B. Meyer^{ID}, G. Milella^{ID}, M. Mormile^{ID},
 A. Mussgiller^{ID}, A. Nürnberg^{ID}, Y. Otariid, D. Pérez Adán^{ID}, A. Raspereza^{ID}, B. Ribeiro Lopes^{ID}, J. Rübenach,
 A. Saggio^{ID}, A. Saibel^{ID}, M. Savitskyi^{ID}, M. Scham^{ID,28,24}, V. Scheurer, S. Schnake^{ID,24}, P. Schütze^{ID},
 C. Schwanenberger^{ID,23}, M. Shchedrolosiev^{ID}, R.E. Sosa Ricardo^{ID}, D. Stafford, N. Tonon^{ID,†},

M. Van De Klundert^{1b}, F. Vazzoler^{1b}, A. Ventura Barroso^{1b}, R. Walsh^{1b}, D. Walter^{1b}, Q. Wang^{1b}, Y. Wen^{1b},
K. Wichmann, L. Wiens^{1b,24}, C. Wissing^{1b}, S. Wuchterl^{1b}, Y. Yang^{1b}, A. Zimmermann Castro Santos^{1b},

Deutsches Elektronen-Synchrotron, Hamburg, Germany

A. Albrecht^{1b}, S. Albrecht^{1b}, M. Antonello^{1b}, S. Bein^{1b}, L. Benato^{1b}, M. Bonanomi^{1b}, P. Connor^{1b},
K. De Leo^{1b}, M. Eich, K. El Morabit^{1b}, F. Feindt, A. Fröhlich, C. Garbers^{1b}, E. Garutti^{1b}, M. Hajheidari,
J. Haller^{1b}, A. Hinzmann^{1b}, H.R. Jabusch^{1b}, G. Kasieczka^{1b}, P. Keicher, R. Klanner^{1b}, W. Korcari^{1b},
T. Kramer^{1b}, V. Kutzner^{1b}, F. Labe^{1b}, J. Lange^{1b}, A. Lobanov^{1b}, C. Matthies^{1b}, A. Mehta^{1b}, L. Moureaux^{1b},
M. Mrowietz, A. Nigamova^{1b}, Y. Nissan, A. Paasch^{1b}, K.J. Pena Rodriguez^{1b}, T. Quadfasel^{1b}, M. Rieger^{1b},
O. Rieger, D. Savoie^{1b}, J. Schindler^{1b}, P. Schleper^{1b}, M. Schröder^{1b}, J. Schwandt^{1b}, M. Sommerhalder^{1b},
H. Stadie^{1b}, G. Steinbrück^{1b}, A. Tews, M. Wolf^{1b},

University of Hamburg, Hamburg, Germany

S. Brommer^{1b}, M. Burkart, E. Butz^{1b}, R. Caspart^{1b}, T. Chwalek^{1b}, A. Dierlamm^{1b}, A. Droll, N. Faltermann^{1b},
M. Giffels^{1b}, J.O. Gosewisch, A. Gottmann^{1b}, F. Hartmann^{1b,29}, M. Horzela^{1b}, U. Husemann^{1b}, M. Klute^{1b},
R. Koppenhöfer^{1b}, M. Link, A. Lintuluoto^{1b}, S. Maier^{1b}, S. Mitra^{1b}, Th. Müller^{1b}, M. Neukum, M. Oh^{1b},
G. Quast^{1b}, K. Rabbertz^{1b}, J. Rauser, M. Schnepf, I. Shvetsov^{1b}, H.J. Simonis^{1b}, N. Trevisani^{1b}, R. Ulrich^{1b},
J. van der Linden^{1b}, R.F. Von Cube^{1b}, M. Wassmer^{1b}, S. Wieland^{1b}, R. Wolf^{1b}, S. Wozniowski^{1b}, S. Wunsch,
X. Zuo^{1b},

Karlsruher Institut fuer Technologie, Karlsruhe, Germany

G. Anagnostou, P. Assiouras^{1b}, G. Daskalakis^{1b}, A. Kyriakis, A. Stakia^{1b},

Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece

M. Diamantopoulou, D. Karasavvas, P. Kontaxakis^{1b}, A. Manousakis-Katsikakis^{1b}, A. Panagiotou,
I. Papavergou^{1b}, N. Saoulidou^{1b}, K. Theofilatos^{1b}, E. Tziaferi^{1b}, K. Vellidis^{1b}, I. Zisopoulos^{1b},

National and Kapodistrian University of Athens, Athens, Greece

G. Bakas^{1b}, T. Chatzistavrou, K. Kousouris^{1b}, I. Papakrivopoulos^{1b}, G. Tsiolitis, A. Zacharopoulou,

National Technical University of Athens, Athens, Greece

K. Adamidis, I. Bestintzanos, I. Evangelou^{1b}, C. Foudas, P. Gianneios^{1b}, C. Kamtsikis, P. Katsoulis,
P. Kokkas^{1b}, P.G. Kosmoglou Kioseoglou^{1b}, N. Manthos^{1b}, I. Papadopoulos^{1b}, J. Strogas^{1b},

University of Ioánnina, Ioánnina, Greece

M. Csanád^{1b}, K. Farkas^{1b}, M.M.A. Gadallah^{1b,30}, S. Lökös^{1b,31}, P. Major^{1b}, K. Mandal^{1b}, G. Pásztor^{1b},
A.J. Rádl^{1b,32}, O. Surányi^{1b}, G.I. Veres^{1b},

MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

M. Bartók^{1b,33}, G. Bencze, C. Hajdu^{1b}, D. Horvath^{1b,34,35}, F. Sikler^{1b}, V. Veszpremi^{1b},

Wigner Research Centre for Physics, Budapest, Hungary

N. Beni^{1b}, S. Czellar, J. Karancsi^{1b,33}, J. Molnar, Z. Szillasi, D. Teyssier^{1b},

Institute of Nuclear Research ATOMKI, Debrecen, Hungary

P. Raics, B. Ujvari^{1b,36},

Institute of Physics, University of Debrecen, Debrecen, Hungary

T. Csorgo ^{ID,32}, F. Nemes ^{ID,32}, T. Novak ^{ID},

Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary

J. Babbar ^{ID}, S. Bansal ^{ID}, S.B. Beri, V. Bhatnagar ^{ID}, G. Chaudhary ^{ID}, S. Chauhan ^{ID}, N. Dhingra ^{ID,37}, R. Gupta, A. Kaur ^{ID}, A. Kaur ^{ID}, H. Kaur ^{ID}, M. Kaur ^{ID}, S. Kumar ^{ID}, P. Kumari ^{ID}, M. Meena ^{ID}, K. Sandeep ^{ID}, T. Sheokand, J.B. Singh ^{ID,38}, A. Singla ^{ID}, A.K. Viridi ^{ID},

Panjab University, Chandigarh, India

A. Ahmed ^{ID}, A. Bhardwaj ^{ID}, A. Chhetri ^{ID}, B.C. Choudhary ^{ID}, A. Kumar ^{ID}, M. Naimuddin ^{ID}, K. Ranjan ^{ID}, S. Saumya ^{ID},

University of Delhi, Delhi, India

S. Baradia ^{ID}, S. Barman ^{ID,39}, S. Bhattacharya ^{ID}, D. Bhowmik, S. Dutta ^{ID}, S. Dutta, B. Gomber ^{ID,40}, M. Maity ³⁹, P. Palit ^{ID}, G. Saha ^{ID}, B. Sahu ^{ID}, S. Sarkar,

Saha Institute of Nuclear Physics, HBNI, Kolkata, India

P.K. Behera ^{ID}, S.C. Behera ^{ID}, P. Kalbhor ^{ID}, J.R. Komaragiri ^{ID,41}, D. Kumar ^{ID,41}, A. Muhammad ^{ID}, L. Panwar ^{ID,41}, R. Pradhan ^{ID}, P.R. Pujahari ^{ID}, A. Sharma ^{ID}, A.K. Sikdar ^{ID}, P.C. Tiwari ^{ID,41}, S. Verma ^{ID},

Indian Institute of Technology Madras, Madras, India

K. Naskar ^{ID,42},

Bhabha Atomic Research Centre, Mumbai, India

T. Aziz, I. Das ^{ID}, S. Dugad, M. Kumar ^{ID}, G.B. Mohanty ^{ID}, P. Suryadevara,

Tata Institute of Fundamental Research-A, Mumbai, India

S. Banerjee ^{ID}, R. Chudasama ^{ID}, M. Guchait ^{ID}, S. Karmakar ^{ID}, S. Kumar ^{ID}, G. Majumder ^{ID}, K. Mazumdar ^{ID}, S. Mukherjee ^{ID}, A. Thachayath ^{ID},

Tata Institute of Fundamental Research-B, Mumbai, India

S. Bahinipati ^{ID,43}, A.K. Das, C. Kar ^{ID}, P. Mal ^{ID}, T. Mishra ^{ID}, V.K. Muraleedharan Nair Bindhu ^{ID,44}, A. Nayak ^{ID,44}, P. Saha ^{ID}, S.K. Swain ^{ID}, D. Vats ^{ID,44},

National Institute of Science Education and Research, An OCC of Homi Bhabha National Institute, Bhubaneswar, Odisha, India

A. Alpana ^{ID}, S. Dube ^{ID}, B. Kansal ^{ID}, A. Laha ^{ID}, S. Pandey ^{ID}, A. Rastogi ^{ID}, S. Sharma ^{ID},

Indian Institute of Science Education and Research (IISER), Pune, India

H. Bakhshiansohi ^{ID,45,46}, E. Khazaie ^{ID,46}, M. Zeinali ^{ID,47},

Isfahan University of Technology, Isfahan, Iran

S. Chenarani ^{ID,48}, S.M. Etesami ^{ID}, M. Khakzad ^{ID}, M. Mohammadi Najafabadi ^{ID},

Institute for Research in Fundamental Sciences (IPM), Tehran, Iran

M. Grunewald ^{ID},

University College Dublin, Dublin, Ireland

M. Abbrescia ^{a,b,ID}, R. Aly ^{a,c,ID,14}, C. Aruta ^{a,b,ID}, A. Colaleo ^{a,ID}, D. Creanza ^{a,c,ID}, N. De Filippis ^{a,c,ID}, M. De Palma ^{a,b,ID}, A. Di Florio ^{a,b,ID}, W. Elmetenawee ^{a,b,ID}, F. Errico ^{a,b,ID}, L. Fiore ^{a,ID}, G. Iaselli ^{a,c,ID}, G. Maggi ^{a,c,ID}, M. Maggi ^{a,ID}, I. Margjeka ^{a,b,ID}, V. Mastrapasqua ^{a,b,ID}, S. My ^{a,b,ID}, S. Nuzzo ^{a,b,ID}, A. Pellecchia ^{a,b,ID}, A. Pompili ^{a,b,ID}, G. Pugliese ^{a,c,ID}, R. Radogna ^{a,ID}, D. Ramos ^{a,ID}, A. Ranieri ^{a,ID},

G. Selvaggi^{a,b, ID}, L. Silvestris^{a, ID}, F.M. Simone^{a,b, ID}, Ü. Sözbilir^{a, ID}, A. Stamerra^{a, ID}, R. Venditti^{a, ID},
P. Verwilligen^{a, ID},

^a INFN Sezione di Bari, Bari, Italy

^b Università di Bari, Bari, Italy

^c Politecnico di Bari, Bari, Italy

G. Abbiendi^{a, ID}, C. Battilana^{a,b, ID}, D. Bonacorsi^{a,b, ID}, L. Borgonovi^{a, ID}, R. Campanini^{a,b, ID}, P. Capiluppi^{a,b, ID},
A. Castro^{a,b, ID}, F.R. Cavallo^{a, ID}, C. Ciocca^{a, ID}, M. Cuffiani^{a,b, ID}, G.M. Dallavalle^{a, ID}, T. Diotallevi^{a,b, ID},
F. Fabbri^{a, ID}, A. Fanfani^{a,b, ID}, P. Giacomelli^{a, ID}, L. Giommi^{a,b, ID}, C. Grandi^{a, ID}, L. Guiducci^{a,b, ID},
S. Lo Meo^{a, ID, 49}, L. Lunerti^{a,b, ID}, S. Marcellini^{a, ID}, G. Masetti^{a, ID}, F.L. Navarra^{a,b, ID}, A. Perrotta^{a, ID},
F. Primavera^{a,b, ID}, A.M. Rossi^{a,b, ID}, T. Rovelli^{a,b, ID}, G.P. Siroli^{a,b, ID},

^a INFN Sezione di Bologna, Bologna, Italy

^b Università di Bologna, Bologna, Italy

S. Costa^{a,b, ID, 50}, A. Di Mattia^{a, ID}, R. Potenza^{a,b}, A. Tricomi^{a,b, ID, 50}, C. Tuve^{a,b, ID},

^a INFN Sezione di Catania, Catania, Italy

^b Università di Catania, Catania, Italy

G. Barbagli^{a, ID}, G. Bardelli^{a,b, ID}, B. Camaiani^{a,b, ID}, A. Cassese^{a, ID}, R. Ceccarelli^{a,b, ID}, V. Ciulli^{a,b, ID},
C. Civinini^{a, ID}, R. D'Alessandro^{a,b, ID}, E. Focardi^{a,b, ID}, G. Latino^{a,b, ID}, P. Lenzi^{a,b, ID}, M. Lizzo^{a,b, ID},
M. Meschini^{a, ID}, S. Paoletti^{a, ID}, R. Seidita^{a,b, ID}, G. Sguazzoni^{a, ID}, L. Viliani^{a, ID},

^a INFN Sezione di Firenze, Firenze, Italy

^b Università di Firenze, Firenze, Italy

L. Benussi^{ID}, S. Bianco^{ID}, S. Meola^{ID, 51}, D. Piccolo^{ID},

INFN Laboratori Nazionali di Frascati, Frascati, Italy

M. Bozzo^{a,b, ID}, P. Chatagnon^{a, ID}, F. Ferro^{a, ID}, E. Robutti^{a, ID}, S. Tosi^{a,b, ID},

^a INFN Sezione di Genova, Genova, Italy

^b Università di Genova, Genova, Italy

A. Benaglia^{a, ID}, G. Boldrini^{a, ID}, F. Brivio^{a,b, ID}, F. Cetorelli^{a,b, ID}, F. De Guio^{a,b, ID}, M.E. Dinardo^{a,b, ID}, P. Dini^{a, ID},
S. Gennai^{a, ID}, A. Ghezzi^{a,b, ID}, P. Govoni^{a,b, ID}, L. Guzzi^{a,b, ID}, M.T. Lucchini^{a,b, ID}, M. Malberti^{a, ID},
S. Malvezzi^{a, ID}, A. Massironi^{a, ID}, D. Menasce^{a, ID}, L. Moroni^{a, ID}, M. Paganoni^{a,b, ID}, D. Pedrini^{a, ID},
B.S. Pinolini^a, S. Ragazzi^{a,b, ID}, N. Redaelli^{a, ID}, T. Tabarelli de Fatis^{a,b, ID}, D. Zuolo^{a,b, ID},

^a INFN Sezione di Milano-Bicocca, Milano, Italy

^b Università di Milano-Bicocca, Milano, Italy

S. Buontempo^{a, ID}, F. Carnevali^{a,b}, N. Cavallo^{a,c, ID}, A. De Iorio^{a,b, ID}, F. Fabozzi^{a,c, ID}, A.O.M. Iorio^{a,b, ID},
L. Lista^{a,b, ID, 52}, P. Paolucci^{a, ID, 29}, B. Rossi^{a, ID}, C. Sciacca^{a,b, ID},

^a INFN Sezione di Napoli, Napoli, Italy

^b Università di Napoli 'Federico II', Napoli, Italy

^c Università della Basilicata, Potenza, Italy

^d Università G. Marconi, Roma, Italy

P. Azzi^{a, ID}, N. Bacchetta^{a, ID, 53}, M. Benettoni^{a, ID}, D. Bisello^{a,b, ID}, P. Bortignon^{a, ID}, A. Bragagnolo^{a,b, ID},
R. Carlin^{a,b, ID}, P. Checchia^{a, ID}, T. Dorigo^{a, ID}, F. Gasparini^{a,b, ID}, U. Gasparini^{a,b, ID}, G. Grosso^a, L. Layer^{a, 54},
E. Lusiani^{a, ID}, M. Margoni^{a,b, ID}, A.T. Meneguzzo^{a,b, ID}, J. Pazzini^{a,b, ID}, P. Ronchese^{a,b, ID}, F. Simonetto^{a,b, ID},
G. Strong^{a, ID}, M. Tosi^{a,b, ID}, H. Yarar^{a,b}, M. Zanetti^{a,b, ID}, P. Zotto^{a,b, ID}, A. Zucchetta^{a,b, ID}, G. Zumerle^{a,b, ID},

^a INFN Sezione di Padova, Padova, Italy

^b Università di Padova, Padova, Italy

^c Università di Trento, Trento, Italy

S. Abu Zeid^{a, ID, 17}, C. Aimè^{a, b, ID}, A. Braghieri^{a, ID}, S. Calzaferri^{a, b, ID}, D. Fiorina^{a, b, ID}, P. Montagna^{a, b, ID},
V. Re^{a, ID}, C. Riccardi^{a, b, ID}, P. Salvini^{a, ID}, I. Vai^{a, ID}, P. Vitulo^{a, b, ID},

^a INFN Sezione di Pavia, Pavia, Italy

^b Università di Pavia, Pavia, Italy

P. Asenov^{a, ID, 55}, G.M. Bilei^{a, ID}, D. Ciangottini^{a, b, ID}, L. Fanò^{a, b, ID}, M. Magherini^{a, b, ID}, G. Mantovani^{a, b},
V. Mariani^{a, b, ID}, M. Menichelli^{a, ID}, F. Moscatelli^{a, ID, 55}, A. Piccinelli^{a, b, ID}, M. Presilla^{a, b, ID}, A. Rossi^{a, b, ID},
A. Santocchia^{a, b, ID}, D. Spiga^{a, ID}, T. Tedeschi^{a, b, ID},

^a INFN Sezione di Perugia, Perugia, Italy

^b Università di Perugia, Perugia, Italy

P. Azzurri^{a, ID}, G. Bagliesi^{a, ID}, V. Bertacchi^{a, c, ID}, R. Bhattacharya^{a, ID}, L. Bianchini^{a, b, ID}, T. Boccali^{a, ID},
E. Bossini^{a, b, ID}, D. Bruschini^{a, c, ID}, R. Castaldi^{a, ID}, M.A. Ciocci^{a, b, ID}, V. D'Amante^{a, d, ID}, R. Dell'Orso^{a, ID},
M.R. Di Domenico^{a, d, ID}, S. Donato^{a, ID}, A. Giassi^{a, ID}, F. Ligabue^{a, c, ID}, G. Mandorli^{a, c, ID}, D. Matos Figueiredo^{a, ID},
A. Messineo^{a, b, ID}, M. Musich^{a, b, ID}, F. Palla^{a, ID}, S. Parolia^{a, ID}, G. Ramirez-Sanchez^{a, c, ID}, A. Rizzi^{a, b, ID},
G. Rolandi^{a, c, ID}, S. Roy Chowdhury^{a, ID}, T. Sarkar^{a, ID}, A. Scribano^{a, ID}, N. Shafiei^{a, b, ID}, P. Spagnolo^{a, ID},
R. Tenchini^{a, ID}, G. Tonelli^{a, b, ID}, N. Turini^{a, d, ID}, A. Venturi^{a, ID}, P.G. Verdini^{a, ID},

^a INFN Sezione di Pisa, Pisa, Italy

^b Università di Pisa, Pisa, Italy

^c Scuola Normale Superiore di Pisa, Pisa, Italy

^d Università di Siena, Siena, Italy

P. Barria^{a, ID}, M. Campana^{a, b, ID}, F. Cavallari^{a, ID}, D. Del Re^{a, b, ID}, E. Di Marco^{a, ID}, M. Diemoz^{a, ID}, E. Longo^{a, b, ID},
P. Meridiani^{a, ID}, G. Organtini^{a, b, ID}, F. Pandolfi^{a, ID}, R. Paramatti^{a, b, ID}, C. Quaranta^{a, b, ID}, S. Rahatlou^{a, b, ID},
C. Rovelli^{a, ID}, F. Santanastasio^{a, b, ID}, L. Soffi^{a, ID}, R. Tramontano^{a, b, ID},

^a INFN Sezione di Roma, Roma, Italy

^b Sapienza Università di Roma, Roma, Italy

N. Amapane^{a, b, ID}, R. Arcidiacono^{a, c, ID}, S. Argiro^{a, b, ID}, M. Arneodo^{a, c, ID}, N. Bartosik^{a, ID}, R. Bellan^{a, b, ID},
A. Bellora^{a, b, ID}, C. Biino^{a, ID}, N. Cartiglia^{a, ID}, M. Costa^{a, b, ID}, R. Covarelli^{a, b, ID}, N. Demaria^{a, ID}, M. Grippo^{a, b, ID},
B. Kiani^{a, b, ID}, F. Legger^{a, ID}, C. Mariotti^{a, ID}, S. Maselli^{a, ID}, A. Mecca^{a, b, ID}, E. Migliore^{a, b, ID}, E. Monteil^{a, b, ID},
M. Monteno^{a, ID}, R. Mulargia^{a, ID}, M.M. Obertino^{a, b, ID}, G. Ortona^{a, ID}, L. Pacher^{a, b, ID}, N. Pastrone^{a, ID},
M. Pelliccioni^{a, ID}, M. Ruspa^{a, c, ID}, K. Shchelina^{a, ID}, F. Siviero^{a, b, ID}, V. Sola^{a, ID}, A. Solano^{a, b, ID}, D. Soldi^{a, b, ID},
A. Staiano^{a, ID}, M. Tornago^{a, b, ID}, D. Trocino^{a, ID}, G. Umoret^{a, b, ID}, A. Vagnerini^{a, b, ID},

^a INFN Sezione di Torino, Torino, Italy

^b Università di Torino, Torino, Italy

^c Università del Piemonte Orientale, Novara, Italy

S. Belforte^{a, ID}, V. Candelise^{a, b, ID}, M. Casarsa^{a, ID}, F. Cossutti^{a, ID}, A. Da Rold^{a, b, ID}, G. Della Ricca^{a, b, ID},
G. Sorrentino^{a, b, ID},

^a INFN Sezione di Trieste, Trieste, Italy

^b Università di Trieste, Trieste, Italy

S. Dogra^{ID}, C. Huh^{ID}, B. Kim^{ID}, D.H. Kim^{ID}, G.N. Kim^{ID}, J. Kim, J. Lee^{ID}, S.W. Lee^{ID}, C.S. Moon^{ID},
Y.D. Oh^{ID}, S.I. Pak^{ID}, M.S. Ryu^{ID}, S. Sekmen^{ID}, Y.C. Yang^{ID},

Kyungpook National University, Daegu, Korea

H. Kim^{ID}, D.H. Moon^{ID},

Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea

E. Asilar ^{ID}, T.J. Kim ^{ID}, J. Park ^{ID},

Hanyang University, Seoul, Korea

S. Choi ^{ID}, S. Han, B. Hong ^{ID}, K. Lee, K.S. Lee ^{ID}, J. Lim, J. Park, S.K. Park, J. Yoo ^{ID},

Korea University, Seoul, Korea

J. Goh ^{ID},

Kyung Hee University, Department of Physics, Seoul, Korea

H.S. Kim ^{ID}, Y. Kim, S. Lee,

Sejong University, Seoul, Korea

J. Almond, J.H. Bhyun, J. Choi ^{ID}, S. Jeon ^{ID}, J. Kim ^{ID}, J.S. Kim, S. Ko ^{ID}, H. Kwon ^{ID}, H. Lee ^{ID}, S. Lee,
B.H. Oh ^{ID}, S.B. Oh ^{ID}, H. Seo ^{ID}, U.K. Yang, I. Yoon ^{ID},

Seoul National University, Seoul, Korea

W. Jang ^{ID}, D.Y. Kang, Y. Kang ^{ID}, D. Kim ^{ID}, S. Kim ^{ID}, B. Ko, J.S.H. Lee ^{ID}, Y. Lee ^{ID}, J.A. Merlin, I.C. Park ^{ID},
Y. Roh, D. Song, I.J. Watson ^{ID}, S. Yang ^{ID},

University of Seoul, Seoul, Korea

S. Ha ^{ID}, H.D. Yoo ^{ID},

Yonsei University, Department of Physics, Seoul, Korea

M. Choi ^{ID}, M.R. Kim ^{ID}, H. Lee, Y. Lee ^{ID}, Y. Lee ^{ID}, I. Yu ^{ID},

Sungkyunkwan University, Suwon, Korea

T. Beyrouthy, Y. Maghrbi ^{ID},

College of Engineering and Technology, American University of the Middle East (AUM), Dasman, Kuwait

K. Dreimanis ^{ID}, G. Pikurs, A. Potrebko ^{ID}, M. Seidel ^{ID}, V. Veckalns ^{ID},

Riga Technical University, Riga, Latvia

M. Ambrozas ^{ID}, A. Carvalho Antunes De Oliveira ^{ID}, A. Juodagalvis ^{ID}, A. Rinkevicius ^{ID}, G. Tamulaitis ^{ID},

Vilnius University, Vilnius, Lithuania

N. Bin Norjoharuddeen ^{ID}, S.Y. Hoh ^{ID,56}, I. Yusuff ^{ID,56}, Z. Zolkapli,

National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia

J.F. Benitez ^{ID}, A. Castaneda Hernandez ^{ID}, H.A. Encinas Acosta, L.G. Gallegos Maríñez, M. León Coello ^{ID},
J.A. Murillo Quijada ^{ID}, A. Sehwat ^{ID}, L. Valencia Palomo ^{ID},

Universidad de Sonora (UNISON), Hermosillo, Mexico

G. Ayala ^{ID}, H. Castilla-Valdez ^{ID}, I. Heredia-De La Cruz ^{ID,57}, R. Lopez-Fernandez ^{ID}, C.A. Mondragon Herrera,
D.A. Perez Navarro ^{ID}, A. Sánchez Hernández ^{ID},

Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico

C. Oropeza Barrera ^{ID}, F. Vazquez Valencia ^{ID},

Universidad Iberoamericana, Mexico City, Mexico

I. Pedraza ^{ID}, H.A. Salazar Ibarguen ^{ID}, C. Uribe Estrada ^{ID},

Benemerita Universidad Autonoma de Puebla, Puebla, Mexico

I. Bubanja, J. Mijuskovic^{[ID](#)},⁵⁸, N. Raicevic^{[ID](#)},

University of Montenegro, Podgorica, Montenegro

A. Ahmad^{[ID](#)}, M.I. Asghar, A. Awais^{[ID](#)}, M.I.M. Awan, M. Gul^{[ID](#)}, H.R. Hoorani^{[ID](#)}, W.A. Khan^{[ID](#)}, M. Shoaib^{[ID](#)},
M. Waqas^{[ID](#)},

National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan

V. Avati, L. Grzanka^{[ID](#)}, M. Malawski^{[ID](#)},

AGH University of Krakow, Faculty of Computer Science, Electronics and Telecommunications, Krakow, Poland

H. Bialkowska^{[ID](#)}, M. Bluj^{[ID](#)}, B. Boimska^{[ID](#)}, M. Górski^{[ID](#)}, M. Kazana^{[ID](#)}, M. Szleper^{[ID](#)}, P. Zalewski^{[ID](#)},

National Centre for Nuclear Research, Swierk, Poland

K. Bunkowski^{[ID](#)}, K. Doroba^{[ID](#)}, A. Kalinowski^{[ID](#)}, M. Konecki^{[ID](#)}, J. Krolikowski^{[ID](#)},

Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland

M. Araujo^{[ID](#)}, P. Bargassa^{[ID](#)}, D. Bastos^{[ID](#)}, A. Boletti^{[ID](#)}, P. Faccioli^{[ID](#)}, M. Gallinaro^{[ID](#)}, J. Hollar^{[ID](#)},
N. Leonardo^{[ID](#)}, T. Niknejad^{[ID](#)}, M. Pisano^{[ID](#)}, J. Seixas^{[ID](#)}, J. Varela^{[ID](#)},

Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal

P. Adzic^{[ID](#)},⁵⁹, M. Dordevic^{[ID](#)}, P. Milenovic^{[ID](#)}, J. Milosevic^{[ID](#)},

VINCA Institute of Nuclear Sciences, University of Belgrade, Belgrade, Serbia

M. Aguilar-Benitez, J. Alcaraz Maestre^{[ID](#)}, A. Álvarez Fernández^{[ID](#)}, M. Barrio Luna, Cristina F. Bedoya^{[ID](#)},
C.A. Carrillo Montoya^{[ID](#)}, M. Cepeda^{[ID](#)}, M. Cerrada^{[ID](#)}, N. Colino^{[ID](#)}, B. De La Cruz^{[ID](#)}, A. Delgado Peris^{[ID](#)},
D. Fernández Del Val^{[ID](#)}, J.P. Fernández Ramos^{[ID](#)}, J. Flix^{[ID](#)}, M.C. Fouz^{[ID](#)}, O. Gonzalez Lopez^{[ID](#)},
S. Goy Lopez^{[ID](#)}, J.M. Hernandez^{[ID](#)}, M.I. Josa^{[ID](#)}, J. León Holgado^{[ID](#)}, D. Moran^{[ID](#)}, C. Perez Dengra^{[ID](#)},
A. Pérez-Calero Yzquierdo^{[ID](#)}, J. Puerta Pelayo^{[ID](#)}, I. Redondo^{[ID](#)}, D.D. Redondo Ferrero^{[ID](#)}, L. Romero,
S. Sánchez Navas^{[ID](#)}, J. Sastre^{[ID](#)}, L. Urda Gómez^{[ID](#)}, J. Vazquez Escobar^{[ID](#)}, C. Willmott,

Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain

J.F. de Trocóniz^{[ID](#)},

Universidad Autónoma de Madrid, Madrid, Spain

B. Alvarez Gonzalez^{[ID](#)}, J. Cuevas^{[ID](#)}, J. Fernandez Menendez^{[ID](#)}, S. Folgueras^{[ID](#)}, I. Gonzalez Caballero^{[ID](#)},
J.R. González Fernández^{[ID](#)}, E. Palencia Cortezon^{[ID](#)}, C. Ramón Álvarez^{[ID](#)}, V. Rodríguez Bouza^{[ID](#)},
A. Soto Rodríguez^{[ID](#)}, A. Trapote^{[ID](#)}, C. Vico Villalba^{[ID](#)},

Universidad de Oviedo, Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA), Oviedo, Spain

J.A. Brochero Cifuentes^{[ID](#)}, I.J. Cabrillo^{[ID](#)}, A. Calderon^{[ID](#)}, J. Duarte Campderros^{[ID](#)}, M. Fernandez^{[ID](#)},
C. Fernandez Madrazo^{[ID](#)}, A. García Alonso, G. Gomez^{[ID](#)}, C. Lasiosa García^{[ID](#)}, C. Martinez Rivero^{[ID](#)},
P. Martinez Ruiz del Arbol^{[ID](#)}, F. Matorras^{[ID](#)}, P. Matorras Cuevas^{[ID](#)}, J. Piedra Gomez^{[ID](#)}, C. Prieels,
A. Ruiz-Jimeno^{[ID](#)}, L. Scodellaro^{[ID](#)}, I. Vila^{[ID](#)}, J.M. Vizan Garcia^{[ID](#)},

Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain

M.K. Jayananda^{[ID](#)}, B. Kailasapathy^{[ID](#)},⁶⁰, D.U.J. Sonnadara^{[ID](#)}, D.D.C. Wickramaratna^{[ID](#)},

University of Colombo, Colombo, Sri Lanka

W.G.D. Dharmaratna^{[ID](#)},⁶¹, K. Liyanage^{[ID](#)}, N. Perera^{[ID](#)}, N. Wickramage^{[ID](#)},

University of Ruhuna, Department of Physics, Matara, Sri Lanka

D. Abbaneo¹, J. Alimena¹, E. Auffray¹, G. Auzinger¹, J. Baechler, P. Baillon[†], D. Barney¹,
 J. Bendavid¹, M. Bianco¹, B. Bilin¹, A. Bocchi¹, E. Brondolin¹, C. Caillol¹, T. Camporesi¹,
 G. Cerminara¹, N. Chernyavskaya¹, S.S. Chhibra¹, S. Choudhury, M. Cipriani¹, L. Cristella¹,
 D. d'Enterria¹, A. Dabrowski¹, A. David¹, A. De Roeck¹, M.M. Defranchis¹, M. Deile¹, M. Dobson¹,
 M. Dünser¹, N. Dupont, F. Fallavollita⁶², A. Florent¹, L. Forthomme¹, G. Franzoni¹, W. Funk¹,
 S. Ghosh¹, S. Giani, D. Gigi, K. Gill¹, F. Glege¹, L. Gouskos¹, E. Govorkova¹, M. Haranko¹,
 J. Hegeman¹, V. Innocente¹, T. James¹, P. Janot¹, J. Kaspar¹, J. Kieseler¹, N. Kratochwil¹,
 S. Laurila¹, P. Lecoq¹, E. Leutgeb¹, C. Lourenço¹, B. Maier¹, L. Malgeri¹, M. Mannelli¹,
 A.C. Marini¹, F. Meijers¹, S. Mersi¹, E. Meschi¹, F. Moortgat¹, M. Mulders¹, S. Orfanelli, L. Orsini,
 F. Pantaleo¹, E. Perez, M. Peruzzi¹, A. Petrilli¹, G. Petrucciani¹, A. Pfeiffer¹, M. Pierini¹, D. Piparo¹,
 M. Pitt¹, H. Qu¹, T. Quast, D. Rabadý¹, A. Racz, G. Reales Gutiérrez, M. Rovere¹, H. Sakulin¹,
 J. Salfeld-Nebgen¹, S. Scarfi¹, M. Selvaggi¹, A. Sharma¹, P. Silva¹, P. Sphicas^{1,63}, A.G. Stahl Leitner¹,
 S. Summers¹, K. Tatar¹, V.R. Tavolaro¹, D. Treille¹, P. Tropea¹, A. Tsirou, J. Wanczyk^{1,64},
 K.A. Wozniak¹, W.D. Zeuner,

CERN, European Organization for Nuclear Research, Geneva, Switzerland

L. Caminada^{1,65}, A. Ebrahimi¹, W. Erdmann¹, R. Horisberger¹, Q. Ingram¹, H.C. Kaestli¹,
 D. Kotlinski¹, C. Lange¹, M. Missiroli^{1,65}, L. Noehte^{1,65}, T. Rohe¹,

Paul Scherrer Institut, Villigen, Switzerland

T.K. Aarrestad¹, K. Androsov^{1,64}, M. Backhaus¹, P. Berger, A. Calandri¹, K. Datta¹, A. De Cosa¹,
 G. Dissertori¹, M. Dittmar, M. Donegà¹, F. Eble¹, M. Galli¹, K. Gedia¹, F. Glessgen¹,
 T.A. Gómez Espinosa¹, C. Grab¹, D. Hits¹, W. Lustermann¹, A.-M. Lyon¹, R.A. Manzoni¹,
 L. Marchese¹, C. Martin Perez¹, A. Mascellani^{1,64}, F. Nessi-Tedaldi¹, J. Niedziela¹, F. Pauss¹,
 V. Perovic¹, S. Pigazzini¹, M.G. Ratti¹, M. Reichmann¹, C. Reissel¹, T. Reitenspiess¹, B. Ristic¹,
 F. Riti¹, D. Ruini, D.A. Sanz Becerra¹, J. Steggemann^{1,64}, D. Valsecchi^{1,29}, R. Wallny¹,

ETH Zurich - Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland

C. Amsler^{1,66}, P. Bärtzchi¹, C. Botta¹, D. Brzdechko, M.F. Canelli¹, K. Cormier¹, A. De Wit¹,
 R. Del Burgo, J.K. Heikkilä¹, M. Huwiler¹, W. Jin¹, A. Jofrehei¹, B. Kilminster¹, S. Leontsinis¹,
 S.P. Liechti¹, A. Macchiolo¹, P. Meiring¹, V.M. Mikuni¹, U. Molinatti¹, I. Neutelings¹, A. Reimers¹,
 P. Robmann, S. Sanchez Cruz¹, K. Schweiger¹, M. Senger¹, Y. Takahashi¹,

Universität Zürich, Zurich, Switzerland

C. Adloff⁶⁷, C.M. Kuo, W. Lin, P.K. Rout¹, S.S. Yu¹,

National Central University, Chung-Li, Taiwan

L. Ceard, Y. Chao¹, K.F. Chen¹, P.s. Chen, H. Cheng¹, W.-S. Hou¹, R. Khurana, G. Kole¹, Y.y. Li¹,
 R.-S. Lu¹, E. Paganis¹, A. Psallidas, A. Steen¹, H.y. Wu, E. Yazgan¹, P.r. Yu,

National Taiwan University (NTU), Taipei, Taiwan

C. Asawatangtrakuldee¹, N. Srimanobhas¹,

Chulalongkorn University, Faculty of Science, Department of Physics, Bangkok, Thailand

D. Agyel^{ID}, F. Boran^{ID}, Z.S. Demiroglu^{ID}, F. Dolek^{ID}, I. Dumanoglu^{ID,68}, E. Eskut^{ID}, Y. Guler^{ID,69},
 E. Gurpinar Guler^{ID,69}, C. Isik^{ID}, O. Kara, A. Kayis Topaksu^{ID}, U. Kiminsu^{ID}, G. Onengut^{ID}, K. Ozdemir^{ID,70},
 A. Polatoz^{ID}, A.E. Simsek^{ID}, B. Tali^{ID,71}, U.G. Tok^{ID}, S. Turkcapar^{ID}, E. Uslan^{ID}, I.S. Zorbakir^{ID},

Çukurova University, Physics Department, Science and Art Faculty, Adana, Turkey

G. Karapinar⁷², K. Ocalan^{ID,73}, M. Yalvac^{ID,74},

Middle East Technical University, Physics Department, Ankara, Turkey

B. Akgun^{ID}, I.O. Atakisi^{ID}, E. Gülmez^{ID}, M. Kaya^{ID,75}, O. Kaya^{ID,76}, S. Tekten^{ID,77},

Bogazici University, Istanbul, Turkey

A. Cakir^{ID}, K. Cankocak^{ID,68}, Y. Komurcu^{ID}, S. Sen^{ID,68},

Istanbul Technical University, Istanbul, Turkey

O. Aydilek^{ID}, S. Cerci^{ID,71}, B. Hacisahinoglu^{ID}, I. Hos^{ID,78}, B. Isildak^{ID,79}, B. Kaynak^{ID}, S. Ozkorucuklu^{ID},
 C. Simsek^{ID}, D. Sunar Cerci^{ID,71},

Istanbul University, Istanbul, Turkey

B. Grynyov^{ID},

Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkiv, Ukraine

L. Levchuk^{ID},

National Science Centre, Kharkiv Institute of Physics and Technology, Kharkiv, Ukraine

D. Anthony^{ID}, E. Bhal^{ID}, J.J. Brooke^{ID}, A. Bundock^{ID}, E. Clement^{ID}, D. Cussans^{ID}, H. Flacher^{ID},
 M. Glowacki, J. Goldstein^{ID}, H.F. Heath^{ID}, L. Kreczko^{ID}, B. Krikler^{ID}, S. Paramesvaran^{ID},
 S. Seif El Nasr-Storey, V.J. Smith^{ID}, N. Stylianou^{ID,80}, K. Walkingshaw Pass, R. White^{ID},

University of Bristol, Bristol, United Kingdom

A.H. Ball, K.W. Bell^{ID}, A. Belyaev^{ID,81}, C. Brew^{ID}, R.M. Brown^{ID}, D.J.A. Cockerill^{ID}, C. Cooke^{ID}, K.V. Ellis,
 K. Harder^{ID}, S. Harper^{ID}, M.-L. Holmberg^{ID,82}, Sh. Jain^{ID}, J. Linacre^{ID}, K. Manolopoulos, D.M. Newbold^{ID},
 E. Olaiya, D. Petyt^{ID}, T. Reis^{ID}, G. Salvi^{ID}, T. Schuh, C.H. Shepherd-Themistocleous^{ID}, I.R. Tomalin^{ID},
 T. Williams^{ID},

Rutherford Appleton Laboratory, Didcot, United Kingdom

R. Bainbridge^{ID}, P. Bloch^{ID}, S. Bonomally, J. Borg^{ID}, C.E. Brown^{ID}, O. Buchmuller, V. Cacchio, V. Cepaitis^{ID},
 G.S. Chahal^{ID,83}, D. Colling^{ID}, J.S. Dancu, P. Dauncey^{ID}, G. Davies^{ID}, J. Davies, M. Della Negra^{ID}, S. Fayer,
 G. Fedi^{ID}, G. Hall^{ID}, M.H. Hassanshahi^{ID}, A. Howard, G. Iles^{ID}, J. Langford^{ID}, L. Lyons^{ID}, A.-M. Magnan^{ID},
 S. Malik, A. Martelli^{ID}, M. Mieskolainen^{ID}, D.G. Monk^{ID}, J. Nash^{ID,84}, M. Pesaresi, B.C. Radburn-Smith^{ID},
 D.M. Raymond, A. Richards, A. Rose^{ID}, E. Scott^{ID}, C. Seez^{ID}, R. Shukla^{ID}, A. Tapper^{ID}, K. Uchida^{ID},
 G.P. Uttley^{ID}, L.H. Vage, T. Virdee^{ID,29}, M. Vojinovic^{ID}, N. Wardle^{ID}, S.N. Webb^{ID}, D. Winterbottom^{ID},

Imperial College, London, United Kingdom

K. Coldham, J.E. Cole^{ID}, A. Khan, P. Kyberd^{ID}, I.D. Reid^{ID},

Brunel University, Uxbridge, United Kingdom

S. Abdullin^{ID}, A. Brinkerhoff^{ID}, B. Caraway^{ID}, J. Dittmann^{ID}, K. Hatakeyama^{ID}, A.R. Kanuganti^{ID},
 B. McMaster^{ID}, M. Saunders^{ID}, S. Sawant^{ID}, C. Sutantawibul^{ID}, J. Wilson^{ID},

Baylor University, Waco, TX, USA

R. Bartek^{ID}, A. Dominguez^{ID}, R. Uniyal^{ID}, A.M. Vargas Hernandez^{ID},

Catholic University of America, Washington, DC, USA

S.I. Cooper^{ID}, D. Di Croce^{ID}, S.V. Gleyzer^{ID}, C. Henderson^{ID}, C.U. Perez^{ID}, P. Rumerio^{ID,85}, C. West^{ID},

The University of Alabama, Tuscaloosa, AL, USA

A. Akpinar^{ID}, A. Albert^{ID}, D. Arcaro^{ID}, C. Cosby^{ID}, Z. Demiragli^{ID}, C. Erice^{ID}, E. Fontanesi^{ID}, D. Gastler^{ID},
S. May^{ID}, J. Rohlf^{ID}, K. Salyer^{ID}, D. Sperka^{ID}, D. Spitzbart^{ID}, I. Suarez^{ID}, A. Tsatsos^{ID}, S. Yuan^{ID},

Boston University, Boston, MA, USA

G. Benelli^{ID}, B. Burkley^{ID}, X. Coubez²⁴, D. Cutts^{ID}, M. Hadley^{ID}, U. Heintz^{ID}, J.M. Hogan^{ID,86}, T. Kwon^{ID},
G. Landsberg^{ID}, K.T. Lau^{ID}, D. Li^{ID}, J. Luo^{ID}, M. Narain^{ID}, N. Pervan^{ID}, S. Sagir^{ID,87}, F. Simpson^{ID},
E. Usai^{ID}, W.Y. Wong, X. Yan^{ID}, D. Yu^{ID}, W. Zhang,

Brown University, Providence, RI, USA

J. Bonilla^{ID}, C. Brainerd^{ID}, R. Breedon^{ID}, M. Calderon De La Barca Sanchez^{ID}, M. Chertok^{ID}, J. Conway^{ID},
P.T. Cox^{ID}, R. Erbacher^{ID}, G. Haza^{ID}, F. Jensen^{ID}, O. Kukral^{ID}, G. Mocellin^{ID}, M. Mulhearn^{ID}, D. Pellett^{ID},
B. Regnery^{ID}, Y. Yao^{ID}, F. Zhang^{ID},

University of California, Davis, Davis, CA, USA

M. Bachtis^{ID}, R. Cousins^{ID}, A. Datta^{ID}, D. Hamilton^{ID}, J. Hauser^{ID}, M. Ignatenko^{ID}, M.A. Iqbal^{ID}, T. Lam^{ID},
E. Manca^{ID}, W.A. Nash^{ID}, S. Regnard^{ID}, D. Saltzberg^{ID}, B. Stone^{ID}, V. Valuev^{ID},

University of California, Los Angeles, CA, USA

R. Clare^{ID}, J.W. Gary^{ID}, M. Gordon, G. Hanson^{ID}, G. Karapostoli^{ID}, O.R. Long^{ID}, N. Manganelli^{ID}, W. Si^{ID},
S. Wimpenny^{ID},

University of California, Riverside, Riverside, CA, USA

J.G. Branson^{ID}, P. Chang^{ID}, S. Cittolin^{ID}, S. Cooperstein^{ID}, D. Diaz^{ID}, J. Duarte^{ID}, R. Gerosa^{ID},
L. Giannini^{ID}, J. Guiang^{ID}, R. Kansal^{ID}, V. Krutelyov^{ID}, R. Lee^{ID}, J. Letts^{ID}, M. Masciovecchio^{ID},
F. Mokhtar^{ID}, M. Pieri^{ID}, B.V. Sathia Narayanan^{ID}, V. Sharma^{ID}, M. Tadel^{ID}, E. Vourliotis^{ID},
F. Würthwein^{ID}, Y. Xiang^{ID}, A. Yagil^{ID},

University of California, San Diego, La Jolla, CA, USA

N. Amin, C. Campagnari^{ID}, M. Citron^{ID}, G. Collura^{ID}, A. Dorsett^{ID}, V. Dutta^{ID}, J. Incandela^{ID},
M. Kilpatrick^{ID}, J. Kim^{ID}, A.J. Li^{ID}, P. Masterson^{ID}, H. Mei^{ID}, M. Oshiro^{ID}, M. Quinnan^{ID}, J. Richman^{ID},
U. Sarica^{ID}, R. Schmitz^{ID}, F. Setti^{ID}, J. Shephlock^{ID}, P. Siddireddy, D. Stuart^{ID}, S. Wang^{ID},

University of California, Santa Barbara - Department of Physics, Santa Barbara, CA, USA

A. Bornheim^{ID}, O. Cerri, I. Dutta^{ID}, A. Latorre, J.M. Lawhorn^{ID}, J. Mao^{ID}, H.B. Newman^{ID}, T.Q. Nguyen^{ID},
M. Spiropulu^{ID}, J.R. Vlimant^{ID}, C. Wang^{ID}, S. Xie^{ID}, R.Y. Zhu^{ID},

California Institute of Technology, Pasadena, CA, USA

J. Alison^{ID}, S. An^{ID}, M.B. Andrews^{ID}, P. Bryant^{ID}, T. Ferguson^{ID}, A. Harilal^{ID}, C. Liu^{ID}, T. Mudholkar^{ID},
S. Murthy^{ID}, M. Paulini^{ID}, A. Roberts^{ID}, A. Sanchez^{ID}, W. Terrill^{ID},

Carnegie Mellon University, Pittsburgh, PA, USA

J.P. Cumalat^{1b}, W.T. Ford^{1b}, A. Hassani^{1b}, G. Karathanasis^{1b}, E. MacDonald, F. Marini^{1b}, A. Perloff^{1b},
C. Savard^{1b}, N. Schonbeck^{1b}, K. Stenson^{1b}, K.A. Ulmer^{1b}, S.R. Wagner^{1b}, N. Zipper^{1b},

University of Colorado Boulder, Boulder, CO, USA

J. Alexander^{1b}, S. Bright-Thonney^{1b}, X. Chen^{1b}, D.J. Cranshaw^{1b}, J. Fan^{1b}, X. Fan^{1b}, D. Gadkari^{1b},
S. Hogan^{1b}, J. Monroy^{1b}, J.R. Patterson^{1b}, D. Quach^{1b}, J. Reichert^{1b}, M. Reid^{1b}, A. Ryd^{1b}, J. Thom^{1b},
P. Wittich^{1b}, R. Zou^{1b},

Cornell University, Ithaca, NY, USA

M. Albrow^{1b}, M. Alyari^{1b}, G. Apollinari^{1b}, A. Apresyan^{1b}, L.A.T. Bauerdick^{1b}, D. Berry^{1b}, J. Berryhill^{1b},
P.C. Bhat^{1b}, K. Burkett^{1b}, J.N. Butler^{1b}, A. Canepa^{1b}, G.B. Cerati^{1b}, H.W.K. Cheung^{1b}, F. Chlebana^{1b},
K.F. Di Petrillo^{1b}, J. Dickinson^{1b}, V.D. Elvira^{1b}, Y. Feng^{1b}, J. Freeman^{1b}, A. Gandrakota^{1b}, Z. Gece^{1b},
L. Gray^{1b}, D. Green, S. Grünendahl^{1b}, D. Guerrero^{1b}, O. Gutsche^{1b}, R.M. Harris^{1b}, R. Heller^{1b},
T.C. Herwig^{1b}, J. Hirschauer^{1b}, L. Horyn^{1b}, B. Jayatilaka^{1b}, S. Jindariani^{1b}, M. Johnson^{1b}, U. Joshi^{1b},
T. Klijnsma^{1b}, B. Klima^{1b}, K.H.M. Kwok^{1b}, S. Lammel^{1b}, D. Lincoln^{1b}, R. Lipton^{1b}, T. Liu^{1b}, C. Madrid^{1b},
K. Maeshima^{1b}, C. Mantilla^{1b}, D. Mason^{1b}, P. McBride^{1b}, P. Merkel^{1b}, S. Mrenna^{1b}, S. Nahn^{1b},
J. Ngadiuba^{1b}, D. Noonan^{1b}, V. Papadimitriou^{1b}, N. Pastika^{1b}, K. Pedro^{1b}, C. Pena^{1b,88}, F. Ravera^{1b},
A. Reinsvold Hall^{1b,89}, L. Ristori^{1b}, E. Sexton-Kennedy^{1b}, N. Smith^{1b}, A. Soha^{1b}, L. Spiegel^{1b}, J. Strait^{1b},
L. Taylor^{1b}, S. Tkaczyk^{1b}, N.V. Tran^{1b}, L. Uplegger^{1b}, E.W. Vaandering^{1b}, I. Zoi^{1b},

Fermi National Accelerator Laboratory, Batavia, IL, USA

P. Avery^{1b}, D. Bourilkov^{1b}, L. Cadamuro^{1b}, V. Cherepanov^{1b}, R.D. Field, M. Kim, E. Koenig^{1b},
J. Konigsberg^{1b}, A. Korytov^{1b}, E. Kuznetsova^{1b}, K.H. Lo, K. Matchev^{1b}, N. Menendez^{1b}, G. Mitselmakher^{1b},
A. Muthirakalayil Madhu^{1b}, N. Rawal^{1b}, D. Rosenzweig^{1b}, S. Rosenzweig^{1b}, K. Shi^{1b}, J. Wang^{1b}, Z. Wu^{1b},

University of Florida, Gainesville, FL, USA

T. Adams^{1b}, A. Askew^{1b}, N. Bower^{1b}, R. Habibullah^{1b}, V. Hagopian^{1b}, T. Kolberg^{1b}, G. Martinez,
H. Prosper^{1b}, O. Viazlo^{1b}, M. Wulansatiti^{1b}, R. Yohay^{1b}, J. Zhang,

Florida State University, Tallahassee, FL, USA

M.M. Baarmand^{1b}, S. Butalla^{1b}, T. Elkafrawy^{1b,17}, M. Hohlmann^{1b}, R. Kumar Verma^{1b}, M. Rahmani,
F. Yumiceva^{1b},

Florida Institute of Technology, Melbourne, FL, USA

M.R. Adams^{1b}, H. Becerril Gonzalez^{1b}, R. Cavanaugh^{1b}, S. Dittmer^{1b}, O. Evdokimov^{1b}, C.E. Gerber^{1b},
D.J. Hofman^{1b}, D.S. Lemos^{1b}, A.H. Merrit^{1b}, C. Mills^{1b}, G. Oh^{1b}, T. Roy^{1b}, S. Rudrabhatla^{1b}, M.B. Tonjes^{1b},
N. Varelas^{1b}, X. Wang^{1b}, Z. Ye^{1b}, J. Yoo^{1b},

University of Illinois Chicago, Chicago, USA, Chicago, USA

M. Alhusseini^{1b}, K. Dilsiz^{1b,90}, L. Emediato^{1b}, G. Karaman^{1b}, O.K. Köseyan^{1b}, J.-P. Merlo,
A. Mestvirishvili^{1b,91}, J. Nachtman^{1b}, O. Neogi, H. Ogul^{1b,92}, Y. Onel^{1b}, A. Penzo^{1b}, C. Snyder, E. Tiras^{1b,93},

The University of Iowa, Iowa City, IA, USA

O. Amram^{1b}, B. Blumenfeld^{1b}, L. Corcodilos^{1b}, J. Davis^{1b}, A.V. Gritsan^{1b}, S. Kyriacou^{1b}, P. Maksimovic^{1b},
J. Roskes^{1b}, S. Sekhar^{1b}, M. Swartz^{1b}, T.Á. Vámi^{1b},

Johns Hopkins University, Baltimore, MD, USA

A. Abreu^{ID}, L.F. Alcerro Alcerro^{ID}, J. Anguiano^{ID}, P. Baringer^{ID}, A. Bean^{ID}, Z. Flowers^{ID}, T. Isidori^{ID}, J. King^{ID}, G. Krintiras^{ID}, M. Lazarovits^{ID}, C. Le Mahieu^{ID}, C. Lindsey, J. Marquez^{ID}, N. Minafra^{ID}, M. Murray^{ID}, M. Nickel^{ID}, C. Rogan^{ID}, C. Royon^{ID}, R. Salvatico^{ID}, S. Sanders^{ID}, C. Smith^{ID}, Q. Wang^{ID}, G. Wilson^{ID},

The University of Kansas, Lawrence, KS, USA

B. Allmond^{ID}, S. Duric, A. Ivanov^{ID}, K. Kaadze^{ID}, A. Kalogeropoulos^{ID}, D. Kim, Y. Maravin^{ID}, T. Mitchell, A. Modak, K. Nam, D. Roy^{ID},

Kansas State University, Manhattan, KS, USA

F. Rebassoo^{ID}, D. Wright^{ID},

Lawrence Livermore National Laboratory, Livermore, CA, USA

E. Adams^{ID}, A. Baden^{ID}, O. Baron, A. Belloni^{ID}, A. Bethani^{ID}, S.C. Eno^{ID}, N.J. Hadley^{ID}, S. Jabeen^{ID}, R.G. Kellogg^{ID}, T. Koeth^{ID}, Y. Lai^{ID}, S. Lascio^{ID}, A.C. Mignerey^{ID}, S. Nabili^{ID}, C. Palmer^{ID}, C. Papageorgakis^{ID}, L. Wang^{ID}, K. Wong^{ID},

University of Maryland, College Park, MD, USA

D. Abercrombie, W. Busza^{ID}, I.A. Cali^{ID}, Y. Chen^{ID}, M. D'Alfonso^{ID}, J. Eysermans^{ID}, C. Freer^{ID}, G. Gomez-Ceballos^{ID}, M. Goncharov, P. Harris, M. Hu^{ID}, D. Kovalskyi^{ID}, J. Krupa^{ID}, Y.-J. Lee^{ID}, K. Long^{ID}, C. Mironov^{ID}, C. Paus^{ID}, D. Rankin^{ID}, C. Roland^{ID}, G. Roland^{ID}, Z. Shi^{ID}, G.S.F. Stephans^{ID}, J. Wang, Z. Wang^{ID}, B. Wyslouch^{ID}, T.J. Yang^{ID},

Massachusetts Institute of Technology, Cambridge, MA, USA

R.M. Chatterjee, B. Crossman^{ID}, A. Evans^{ID}, J. Hiltbrand^{ID}, B.M. Joshi^{ID}, C. Kapsiak^{ID}, M. Krohn^{ID}, Y. Kubota^{ID}, J. Mans^{ID}, M. Revering^{ID}, R. Rusack^{ID}, R. Saradhy^{ID}, N. Schroeder^{ID}, N. Strobbe^{ID}, M.A. Wadud^{ID},

University of Minnesota, Minneapolis, MN, USA

L.M. Cremaldi^{ID},

University of Mississippi, Oxford, MS, USA

K. Bloom^{ID}, M. Bryson, D.R. Claes^{ID}, C. Fangmeier^{ID}, L. Finco^{ID}, F. Golf^{ID}, C. Joo^{ID}, R. Kamalieddin, I. Kravchenko^{ID}, I. Reed^{ID}, J.E. Siado^{ID}, G.R. Snow[†], W. Tabb^{ID}, A. Wightman^{ID}, F. Yan^{ID}, A.G. Zecchinelli^{ID},

University of Nebraska-Lincoln, Lincoln, NE, USA

G. Agarwal^{ID}, H. Bandyopadhyay^{ID}, L. Hay^{ID}, I. Iashvili^{ID}, A. Kharchilava^{ID}, C. McLean^{ID}, M. Morris^{ID}, D. Nguyen^{ID}, J. Pekkanen^{ID}, S. Rappoccio^{ID}, A. Williams^{ID},

State University of New York at Buffalo, Buffalo, NY, USA

G. Alverson^{ID}, E. Barberis^{ID}, Y. Haddad^{ID}, Y. Han^{ID}, A. Krishna^{ID}, J. Li^{ID}, J. Lidrych^{ID}, G. Madigan^{ID}, B. Marzocchi^{ID}, D.M. Morse^{ID}, V. Nguyen^{ID}, T. Orimoto^{ID}, A. Parker^{ID}, L. Skinnari^{ID}, A. Tishelman-Charny^{ID}, T. Wamorkar^{ID}, B. Wang^{ID}, A. Wisecarver^{ID}, D. Wood^{ID},

Northeastern University, Boston, MA, USA

S. Bhattacharya^{ID}, J. Bueghly, Z. Chen^{ID}, A. Gilbert^{ID}, K.A. Hahn^{ID}, Y. Liu^{ID}, N. Odell^{ID}, M.H. Schmitt^{ID}, M. Velasco,

Northwestern University, Evanston, IL, USA

R. Band ^{id}, R. Bucci, M. Cremonesi, A. Das ^{id}, R. Goldouzian ^{id}, M. Hildreth ^{id}, K. Hurtado Anampa ^{id},
C. Jessop ^{id}, K. Lannon ^{id}, J. Lawrence ^{id}, N. Loukas ^{id}, L. Lutton ^{id}, J. Mariano, N. Marinelli, I. Mcalister,
T. McCauley ^{id}, C. Mcgrady ^{id}, K. Mohrman ^{id}, C. Moore ^{id}, Y. Musienko ^{id,13}, R. Ruchti ^{id}, A. Townsend ^{id},
M. Wayne ^{id}, H. Yockey, M. Zarucki ^{id}, L. Zygala ^{id},

University of Notre Dame, Notre Dame, IN, USA

B. Bylsma, M. Carrigan ^{id}, L.S. Durkin ^{id}, B. Francis ^{id}, C. Hill ^{id}, M. Joyce ^{id}, A. Lesauvage ^{id},
M. Nunez Ornelas ^{id}, K. Wei, B.L. Winer ^{id}, B.R. Yates ^{id},

The Ohio State University, Columbus, OH, USA

F.M. Addesa ^{id}, P. Das ^{id}, G. Dezoort ^{id}, P. Elmer ^{id}, A. Frankenthal ^{id}, B. Greenberg ^{id}, N. Haubrich ^{id},
S. Higginbotham ^{id}, G. Kopp ^{id}, S. Kwan ^{id}, D. Lange ^{id}, D. Marlow ^{id}, I. Ojalvo ^{id}, J. Olsen ^{id}, D. Stickland ^{id},
C. Tully ^{id},

Princeton University, Princeton, NJ, USA

S. Malik ^{id}, S. Norberg,

University of Puerto Rico, Mayaguez, PR, USA

A.S. Bakshi ^{id}, V.E. Barnes ^{id}, R. Chawla ^{id}, S. Das ^{id}, L. Gutay, M. Jones ^{id}, A.W. Jung ^{id}, D. Kondratyev ^{id},
A.M. Koshy, M. Liu ^{id}, G. Negro ^{id}, N. Neumeister ^{id}, G. Paspalaki ^{id}, S. Piperov ^{id}, A. Purohit ^{id}, H. Qiu,
J.F. Schulte ^{id}, M. Stojanovic ^{id}, J. Thieman ^{id}, F. Wang ^{id}, R. Xiao ^{id}, W. Xie ^{id},

Purdue University, West Lafayette, IN, USA

J. Dolen ^{id}, N. Parashar ^{id},

Purdue University Northwest, Hammond, IN, USA

D. Acosta ^{id}, A. Baty ^{id}, T. Carnahan ^{id}, S. Dildick ^{id}, K.M. Ecklund ^{id}, P.J. Fernández Manteca ^{id}, S. Freed,
P. Gardner, F.J.M. Geurts ^{id}, A. Kumar ^{id}, W. Li ^{id}, B.P. Padley ^{id}, R. Redjimi, J. Rotter ^{id}, S. Yang ^{id},
E. Yigitbasi ^{id}, L. Zhang ⁹⁴, Y. Zhang ^{id},

Rice University, Houston, TX, USA

A. Bodek ^{id}, P. de Barbaro ^{id}, R. Demina ^{id}, J.L. Dulemba ^{id}, C. Fallon, T. Ferbel ^{id}, M. Galanti,
A. Garcia-Bellido ^{id}, O. Hindrichs ^{id}, A. Khukhunaishvili ^{id}, P. Parygin ^{id}, E. Popova ^{id}, E. Ranken ^{id},
R. Taus ^{id}, G.P. Van Onsem ^{id},

University of Rochester, Rochester, NY, USA

K. Goulios ^{id},

The Rockefeller University, New York, NY, USA

B. Chiarito, J.P. Chou ^{id}, Y. Gershtein ^{id}, E. Halkiadakis ^{id}, A. Hart ^{id}, M. Heindl ^{id}, D. Jaroslowski ^{id},
O. Karacheban ^{id,27}, I. Laflotte ^{id}, A. Lath ^{id}, R. Montalvo, K. Nash, M. Osherson ^{id}, H. Routray ^{id}, S. Salur ^{id},
S. Schnetzer, S. Somalwar ^{id}, R. Stone ^{id}, S.A. Thayil ^{id}, S. Thomas, H. Wang ^{id},

Rutgers, The State University of New Jersey, Piscataway, NJ, USA

H. Acharya, A.G. Delannoy ^{id}, S. Fiorendi ^{id}, T. Holmes ^{id}, E. Nibigira ^{id}, S. Spanier ^{id},

University of Tennessee, Knoxville, TN, USA

O. Bouhali^{ID,95}, M. Dalchenko^{ID}, A. Delgado^{ID}, R. Eusebi^{ID}, J. Gilmore^{ID}, T. Huang^{ID}, T. Kamon^{ID,96},
H. Kim^{ID}, S. Luo^{ID}, S. Malhotra, R. Mueller^{ID}, D. Overton^{ID}, D. Rathjens^{ID}, A. Safonov^{ID},

Texas A&M University, College Station, TX, USA

N. Akchurin^{ID}, J. Damgov^{ID}, V. Hegde^{ID}, K. Lamichhane^{ID}, S.W. Lee^{ID}, T. Mengke, S. Muthumuni^{ID},
T. Peltola^{ID}, I. Volobouev^{ID}, A. Whitbeck^{ID},

Texas Tech University, Lubbock, TX, USA

E. Appelt^{ID}, S. Greene, A. Gurrola^{ID}, W. Johns^{ID}, A. Melo^{ID}, F. Romeo^{ID}, P. Sheldon^{ID}, S. Tuo^{ID},
J. Velkovska^{ID}, J. Viinikainen^{ID},

Vanderbilt University, Nashville, TN, USA

B. Cardwell^{ID}, B. Cox^{ID}, G. Cummings^{ID}, J. Hakala^{ID}, R. Hirosky^{ID}, A. Ledovsky^{ID}, A. Li^{ID}, C. Neu^{ID},
C.E. Perez Lara^{ID}, B. Tannenwald^{ID},

University of Virginia, Charlottesville, VA, USA

P.E. Karchin^{ID}, N. Poudyal^{ID},

Wayne State University, Detroit, MI, USA

S. Banerjee^{ID}, K. Black^{ID}, T. Bose^{ID}, S. Dasu^{ID}, I. De Bruyn^{ID}, P. Everaerts^{ID}, C. Galloni, H. He^{ID},
M. Herndon^{ID}, A. Herve^{ID}, C.K. Koraka^{ID}, A. Lanaro, A. Loeliger^{ID}, R. Loveless^{ID},
J. Madhusudanan Sreekala^{ID}, A. Mallampalli^{ID}, A. Mohammadi^{ID}, S. Mondal, G. Parida^{ID}, D. Pinna, A. Savin,
V. Shang^{ID}, V. Sharma^{ID}, W.H. Smith^{ID}, D. Teague, H.F. Tsoi^{ID}, W. Vetens^{ID},

University of Wisconsin - Madison, Madison, WI, USA

S. Afanasiev^{ID}, V. Andreev^{ID}, Yu. Andreev^{ID}, T. Aushev^{ID}, M. Azarkin^{ID}, A. Babaev^{ID}, A. Belyaev^{ID},
V. Blinov⁹⁷, E. Boos^{ID}, V. Borshch^{ID}, D. Budkouski^{ID}, V. Chekhovsky, R. Chistov^{ID,97}, M. Danilov^{ID,97},
A. Demiyanov^{ID}, A. Dermenev^{ID}, T. Dimova^{ID,97}, I. Dremin^{ID}, V. Epshteyn^{ID}, A. Ershov^{ID}, G. Gavrillov^{ID},
V. Gavrillov^{ID}, S. Gninenko^{ID}, V. Golovtsov^{ID}, N. Golubev^{ID}, I. Golutvin^{ID}, I. Gorbunov^{ID}, A. Gribushin^{ID},
Y. Ivanov^{ID}, V. Kachanov^{ID}, L. Kardapoltsev^{ID,97}, V. Karjavine^{ID}, A. Karneyeu^{ID}, L. Khein, V. Kim^{ID,97},
M. Kirakosyan, D. Kirpichnikov^{ID}, M. Kirsanov^{ID}, O. Kodolova^{ID,98}, D. Konstantinov^{ID}, V. Korenkov^{ID},
V. Korotkikh, A. Kozyrev^{ID,97}, N. Krasnikov^{ID}, A. Lanev^{ID}, P. Levchenko^{ID}, A. Litomin, N. Lychkovskaya^{ID},
V. Makarenko^{ID}, A. Malakhov^{ID}, V. Matveev^{ID,97}, V. Murzin^{ID}, A. Nikitenko^{ID,99}, S. Obraztsov^{ID},
I. Ovtin^{ID,97}, V. Palichik^{ID}, V. Perelygin^{ID}, S. Petrushanko^{ID}, S. Polikarpov^{ID,97}, V. Popov, O. Radchenko^{ID,97},
M. Savina^{ID}, V. Savrin^{ID}, D. Selivanova^{ID}, V. Shalaev^{ID}, S. Shmatov^{ID}, S. Shulha^{ID}, Y. Skovpen^{ID,97},
S. Slabospitskii^{ID}, V. Smirnov^{ID}, A. Snigirev^{ID}, D. Sosnov^{ID}, V. Sulimov^{ID}, E. Tcherniaev^{ID}, A. Terkulov^{ID},
O. Teryaev^{ID}, I. Tlisova^{ID}, M. Toms^{ID}, A. Toropin^{ID}, L. Uvarov^{ID}, A. Uzunian^{ID}, I. Vardanyan^{ID},
E. Vlasov^{ID}, A. Vorobyev, N. Voytishin^{ID}, B.S. Yuldashev¹⁰⁰, A. Zarubin^{ID}, I. Zhizhin^{ID}, A. Zhokin^{ID},

Authors affiliated with an institute or an international laboratory covered by a cooperation agreement with CERN

[†] Deceased.

¹ Also at Yerevan State University, Yerevan, Armenia.

² Also at TU Wien, Vienna, Austria.

³ Also at Institute of Basic and Applied Sciences, Faculty of Engineering, Arab Academy for Science, Technology and Maritime Transport, Alexandria, Egypt.

⁴ Also at Université Libre de Bruxelles, Bruxelles, Belgium.

⁵ Also at Universidade Estadual de Campinas, Campinas, Brazil.

⁶ Also at Federal University of Rio Grande do Sul, Porto Alegre, Brazil.

⁷ Also at UFMS, Nova Andradina, Brazil.

- ⁸ Also at The University of the State of Amazonas, Manaus, Brazil.
- ⁹ Also at University of Chinese Academy of Sciences, Beijing, China.
- ¹⁰ Also at Nanjing Normal University, Nanjing, China.
- ¹¹ Now at The University of Iowa, Iowa City, Iowa, USA.
- ¹² Also at University of Chinese Academy of Sciences, Beijing, China.
- ¹³ Also at an institute or an international laboratory covered by a cooperation agreement with CERN.
- ¹⁴ Also at Helwan University, Cairo, Egypt.
- ¹⁵ Now at Zewail City of Science and Technology, Zewail, Egypt.
- ¹⁶ Also at British University in Egypt, Cairo, Egypt.
- ¹⁷ Now at Ain Shams University, Cairo, Egypt.
- ¹⁸ Also at Purdue University, West Lafayette, Indiana, USA.
- ¹⁹ Also at Université de Haute Alsace, Mulhouse, France.
- ²⁰ Also at Department of Physics, Tsinghua University, Beijing, China.
- ²¹ Also at Tbilisi State University, Tbilisi, Georgia.
- ²² Also at Erzincan Binali Yildirim University, Erzincan, Turkey.
- ²³ Also at University of Hamburg, Hamburg, Germany.
- ²⁴ Also at RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany.
- ²⁵ Also at Isfahan University of Technology, Isfahan, Iran.
- ²⁶ Also at Bergische University Wuppertal (BUW), Wuppertal, Germany.
- ²⁷ Also at Brandenburg University of Technology, Cottbus, Germany.
- ²⁸ Also at Forschungszentrum Jülich, Juelich, Germany.
- ²⁹ Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland.
- ³⁰ Also at Physics Department, Faculty of Science, Assiut University, Assiut, Egypt.
- ³¹ Also at Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary.
- ³² Also at Wigner Research Centre for Physics, Budapest, Hungary.
- ³³ Also at Institute of Physics, University of Debrecen, Debrecen, Hungary.
- ³⁴ Also at Institute of Nuclear Research ATOMKI, Debrecen, Hungary.
- ³⁵ Now at Universitatea Babeş-Bolyai - Facultatea de Fizica, Cluj-Napoca, Romania.
- ³⁶ Also at Faculty of Informatics, University of Debrecen, Debrecen, Hungary.
- ³⁷ Also at Punjab Agricultural University, Ludhiana, India.
- ³⁸ Also at UPES - University of Petroleum and Energy Studies, Dehradun, India.
- ³⁹ Also at University of Visva-Bharati, Santiniketan, India.
- ⁴⁰ Also at University of Hyderabad, Hyderabad, India.
- ⁴¹ Also at Indian Institute of Science (IISc), Bangalore, India.
- ⁴² Also at Indian Institute of Technology (IIT), Mumbai, India.
- ⁴³ Also at IIT Bhubaneswar, Bhubaneswar, India.
- ⁴⁴ Also at Institute of Physics, Bhubaneswar, India.
- ⁴⁵ Also at Deutsches Elektronen-Synchrotron, Hamburg, Germany.
- ⁴⁶ Now at Department of Physics, Isfahan University of Technology, Isfahan, Iran.
- ⁴⁷ Also at Sharif University of Technology, Tehran, Iran.
- ⁴⁸ Also at Department of Physics, University of Science and Technology of Mazandaran, Behshahr, Iran.
- ⁴⁹ Also at Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Bologna, Italy.
- ⁵⁰ Also at Centro Siciliano di Fisica Nucleare e di Struttura Della Materia, Catania, Italy.
- ⁵¹ Also at Università degli Studi Guglielmo Marconi, Roma, Italy.
- ⁵² Also at Scuola Superiore Meridionale, Università di Napoli 'Federico II', Napoli, Italy.
- ⁵³ Also at Fermi National Accelerator Laboratory, Batavia, Illinois, USA.
- ⁵⁴ Also at Università di Napoli 'Federico II', Napoli, Italy.
- ⁵⁵ Also at Consiglio Nazionale delle Ricerche - Istituto Officina dei Materiali, Perugia, Italy.
- ⁵⁶ Also at Department of Applied Physics, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Malaysia.
- ⁵⁷ Also at Consejo Nacional de Ciencia y Tecnología, Mexico City, Mexico.
- ⁵⁸ Also at IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France.
- ⁵⁹ Also at Faculty of Physics, University of Belgrade, Belgrade, Serbia.
- ⁶⁰ Also at Trincomalee Campus, Eastern University, Sri Lanka, Nilaveli, Sri Lanka.
- ⁶¹ Also at Saegis Campus, Nugegoda, Sri Lanka.
- ⁶² Also at INFN Sezione di Pavia, Università di Pavia, Pavia, Italy.
- ⁶³ Also at National and Kapodistrian University of Athens, Athens, Greece.
- ⁶⁴ Also at Ecole Polytechnique Fédérale Lausanne, Lausanne, Switzerland.
- ⁶⁵ Also at Universität Zürich, Zurich, Switzerland.
- ⁶⁶ Also at Stefan Meyer Institute for Subatomic Physics, Vienna, Austria.
- ⁶⁷ Also at Laboratoire d'Annecy-le-Vieux de Physique des Particules, IN2P3-CNRS, Annecy-le-Vieux, France.
- ⁶⁸ Also at Near East University, Research Center of Experimental Health Science, Mersin, Turkey.
- ⁶⁹ Also at Konya Technical University, Konya, Turkey.
- ⁷⁰ Also at Izmir Bakircay University, Izmir, Turkey.
- ⁷¹ Also at Adiyaman University, Adiyaman, Turkey.
- ⁷² Also at Istanbul Gedik University, Istanbul, Turkey.
- ⁷³ Also at Necmettin Erbakan University, Konya, Turkey.
- ⁷⁴ Also at Bozok Universitetesi Rektörlüğü, Yozgat, Turkey.
- ⁷⁵ Also at Marmara University, Istanbul, Turkey.
- ⁷⁶ Also at Milli Savunma University, Istanbul, Turkey.
- ⁷⁷ Also at Kafkas University, Kars, Turkey.
- ⁷⁸ Also at Istanbul University - Cerrahpasa, Faculty of Engineering, Istanbul, Turkey.

- ⁷⁹ Also at Ozyegin University, Istanbul, Turkey.
- ⁸⁰ Also at Vrije Universiteit Brussel, Brussel, Belgium.
- ⁸¹ Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom.
- ⁸² Also at University of Bristol, Bristol, United Kingdom.
- ⁸³ Also at IPPP Durham University, Durham, United Kingdom.
- ⁸⁴ Also at Monash University, Faculty of Science, Clayton, Australia.
- ⁸⁵ Also at Università di Torino, Torino, Italy.
- ⁸⁶ Also at Bethel University, St. Paul, Minnesota, USA.
- ⁸⁷ Also at Karamanoğlu Mehmetbey University, Karaman, Turkey.
- ⁸⁸ Also at California Institute of Technology, Pasadena, California, USA.
- ⁸⁹ Also at United States Naval Academy, Annapolis, Maryland, USA.
- ⁹⁰ Also at Bingol University, Bingol, Turkey.
- ⁹¹ Also at Georgian Technical University, Tbilisi, Georgia.
- ⁹² Also at Sinop University, Sinop, Turkey.
- ⁹³ Also at Erciyes University, Kayseri, Turkey.
- ⁹⁴ Also at Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) - Fudan University, Shanghai, China.
- ⁹⁵ Also at Texas A&M University at Qatar, Doha, Qatar.
- ⁹⁶ Also at Kyungpook National University, Daegu, Korea.
- ⁹⁷ Also at another institute or international laboratory covered by a cooperation agreement with CERN.
- ⁹⁸ Also at Yerevan Physics Institute, Yerevan, Armenia.
- ⁹⁹ Also at Imperial College, London, United Kingdom.
- ¹⁰⁰ Also at Institute of Nuclear Physics of the Uzbekistan Academy of Sciences, Tashkent, Uzbekistan.