

Exclusive Branching-Fraction Measurements of Semileptonic τ Decays into Three Charged Hadrons, into $\varphi\pi^-\nu_\tau$, and into $\varphi K^-\nu_\tau$

- B. Aubert,¹ M. Bona,¹ D. Boutigny,¹ F. Couderc,¹ Y. Karyotakis,¹ J. P. Lees,¹ V. Poireau,¹ V. Tisserand,¹ A. Zghiche,¹ E. Grauges,² A. Palano,³ J. C. Chen,⁴ N. D. Qi,⁴ G. Rong,⁴ P. Wang,⁴ Y. S. Zhu,⁴ G. Eigen,⁵ I. Ofte,⁵ B. Stugu,⁵ G. S. Abrams,⁶ M. Battaglia,⁶ D. N. Brown,⁶ J. Button-Shafer,⁶ R. N. Cahn,⁶ E. Charles,⁶ M. S. Gill,⁶ Y. Groysman,⁶ R. G. Jacobsen,⁶ J. A. Kadyk,⁶ L. T. Kerth,⁶ Yu. G. Kolomensky,⁶ G. Kukartsev,⁶ D. Lopes Pegna,⁶ G. Lynch,⁶ L. M. Mir,⁶ T. J. Orimoto,⁶ M. Pripstein,⁶ N. A. Roe,⁶ M. T. Ronan,⁶ W. A. Wenzel,⁶ P. del Amo Sanchez,⁷ M. Barrett,⁷ K. E. Ford,⁷ T. J. Harrison,⁷ A. J. Hart,⁷ C. M. Hawkes,⁷ A. T. Watson,⁷ T. Held,⁸ H. Koch,⁸ B. Lewandowski,⁸ M. Pelizaeus,⁸ K. Peters,⁸ T. Schroeder,⁸ M. Steinke,⁸ J. T. Boyd,⁹ J. P. Burke,⁹ W. N. Cottingham,⁹ D. Walker,⁹ D. J. Asgeirsson,¹⁰ T. Cuhadar-Donszelmann,¹⁰ B. G. Fulson,¹⁰ C. Hearty,¹⁰ N. S. Knecht,¹⁰ T. S. Mattison,¹⁰ J. A. McKenna,¹⁰ A. Khan,¹¹ P. Kyberd,¹¹ M. Saleem,¹¹ D. J. Sherwood,¹¹ L. Teodorescu,¹¹ V. E. Blinov,¹² A. D. Bukan,¹² V. P. Druzhinin,¹² V. B. Golubev,¹² A. P. Onuchin,¹² S. I. Serednyakov,¹² Yu. I. Skovpen,¹² E. P. Solodov,¹² K. Yu Todyshev,¹² D. S. Best,¹³ M. Bondioli,¹³ M. Bruinsma,¹³ M. Chao,¹³ S. Curry,¹³ I. Eschrich,¹³ D. Kirkby,¹³ A. J. Lankford,¹³ P. Lund,¹³ M. Mandelkern,¹³ W. Roethel,¹³ D. P. Stoker,¹³ S. Abachi,¹⁴ C. Buchanan,¹⁴ S. D. Foulkes,¹⁵ J. W. Gary,¹⁵ O. Long,¹⁵ B. C. Shen,¹⁵ K. Wang,¹⁵ L. Zhang,¹⁵ H. K. Hadavand,¹⁶ E. J. Hill,¹⁶ H. P. Paar,¹⁶ S. Rahatlou,¹⁶ V. Sharma,¹⁶ J. W. Berryhill,¹⁷ C. Campagnari,¹⁷ A. Cunha,¹⁷ B. Dahmes,¹⁷ T. M. Hong,¹⁷ D. Kovalskyi,¹⁷ J. D. Richman,¹⁷ T. W. Beck,¹⁸ A. M. Eisner,¹⁸ C. J. Flacco,¹⁸ C. A. Heusch,¹⁸ J. Kroeseberg,¹⁸ W. S. Lockman,¹⁸ G. Nesom,¹⁸ T. Schalk,¹⁸ B. A. Schumm,¹⁸ A. Seiden,¹⁸ P. Spradlin,¹⁸ D. C. Williams,¹⁸ M. G. Wilson,¹⁸ J. Albert,¹⁹ E. Chen,¹⁹ C. H. Cheng,¹⁹ A. Dvoretskii,¹⁹ F. Fang,¹⁹ D. G. Hitlin,¹⁹ I. Narsky,¹⁹ T. Piatenko,¹⁹ F. C. Porter,¹⁹ G. Mancinelli,²⁰ B. T. Meadows,²⁰ K. Mishra,²⁰ M. D. Sokoloff,²⁰ F. Blanc,²¹ P. C. Bloom,²¹ S. Chen,²¹ W. T. Ford,²¹ J. F. Hirschauer,²¹ A. Kreisel,²¹ M. Nagel,²¹ U. Nauenberg,²¹ A. Olivas,²¹ W. O. Ruddick,²¹ J. G. Smith,²¹ K. A. Ulmer,²¹ S. R. Wagner,²¹ J. Zhang,²¹ A. Chen,²² E. A. Eckhart,²² A. Soffer,²² W. H. Toki,²² R. J. Wilson,²² F. Winkelmeier,²² Q. Zeng,²² D. D. Altenburg,²³ E. Feltresi,²³ A. Hauke,²³ H. Jasper,²³ J. Merkel,²³ A. Petzold,²³ B. Spaan,²³ T. Brandt,²⁴ V. Klose,²⁴ H. M. Lacker,²⁴ W. F. Mader,²⁴ R. Nogowski,²⁴ J. Schubert,²⁴ K. R. Schubert,²⁴ R. Schwierz,²⁴ J. E. Sundermann,²⁴ A. Volk,²⁴ D. Bernard,²⁵ G. R. Bonneau,²⁵ E. Latour,²⁵ Ch. Thiebaux,²⁵ M. Verderi,²⁵ P. J. Clark,²⁶ W. Grisl,²⁶ F. Muheim,²⁶ S. Playfer,²⁶ A. I. Robertson,²⁶ Y. Xie,²⁶ M. Andreotti,²⁷ D. Bettoni,²⁷ C. Bozzi,²⁷ R. Calabrese,²⁷ G. Cibinetto,²⁷ E. Luppi,²⁷ M. Negrini,²⁷ A. Petrella,²⁷ L. Piemontese,²⁷ E. Prencipe,²⁷ F. Anulli,²⁸ R. Baldini-Ferroli,²⁸ A. Calcaterra,²⁸ R. de Sangro,²⁸ G. Finocchiaro,²⁸ S. Pacetti,²⁸ P. Patteri,²⁸ I. M. Peruzzi,^{28,*} M. Piccolo,²⁸ M. Rama,²⁸ A. Zallo,²⁸ A. Buzzo,²⁹ R. Contri,²⁹ M. Lo Vetere,²⁹ M. M. Macri,²⁹ M. R. Monge,²⁹ S. Passaggio,²⁹ C. Patrignani,²⁹ E. Robutti,²⁹ A. Santroni,²⁹ S. Tosi,²⁹ G. Brandenburg,³⁰ K. S. Chaisanguanthum,³⁰ C. L. Lee,³⁰ M. Morii,³⁰ J. Wu,³⁰ R. S. Dubitzky,³¹ J. Marks,³¹ S. Schenk,³¹ U. Uwer,³¹ D. J. Bard,³² W. Bhimji,³² D. A. Bowerman,³² P. D. Dauncey,³² U. Egede,³² R. L. Flack,³² J. A. Nash,³² M. B. Nikolich,³² W. Panduro Vazquez,³² P. K. Behera,³³ X. Chai,³³ M. J. Charles,³³ U. Mallik,³³ N. T. Meyer,³³ V. Ziegler,³³ J. Cochran,³⁴ H. B. Crawley,³⁴ L. Dong,³⁴ V. Eyges,³⁴ W. T. Meyer,³⁴ S. Prell,³⁴ E. I. Rosenberg,³⁴ A. E. Rubin,³⁴ A. V. Gritsan,³⁵ A. G. Denig,³⁶ M. Fritsch,³⁶ G. Schott,³⁶ N. Arnaud,³⁷ M. Davier,³⁷ G. Grosdidier,³⁷ A. Höcker,³⁷ V. Lepeltier,³⁷ F. Le Diberder,³⁷ A. M. Lutz,³⁷ A. Oyanguren,³⁷ S. Pruvot,³⁷ S. Rodier,³⁷ P. Roudeau,³⁷ M. H. Schune,³⁷ J. Serrano,³⁷ A. Stocchi,³⁷ W. F. Wang,³⁷ G. Wormser,³⁷ D. J. Lange,³⁸ D. M. Wright,³⁸ C. A. Chavez,³⁹ I. J. Forster,³⁹ J. R. Fry,³⁹ E. Gabathuler,³⁹ R. Gamet,³⁹ K. A. George,³⁹ D. E. Hutchcroft,³⁹ D. J. Payne,³⁹ K. C. Schofield,³⁹ C. Touramanis,³⁹ A. J. Bevan,⁴⁰ C. K. Clarke,⁴⁰ F. Di Lodovico,⁴⁰ W. Menges,⁴⁰ R. Sacco,⁴⁰ G. Cowan,⁴¹ H. U. Flaecher,⁴¹ D. A. Hopkins,⁴¹ P. S. Jackson,⁴¹ T. R. McMahon,⁴¹ F. Salvatore,⁴¹ A. C. Wren,⁴¹ D. N. Brown,⁴² C. L. Davis,⁴² J. Allison,⁴³ N. R. Barlow,⁴³ R. J. Barlow,⁴³ Y. M. Chia,⁴³ C. L. Edgar,⁴³ G. D. Lafferty,⁴³ M. T. Naisbit,⁴³ J. C. Williams,⁴³ J. I. Yi,⁴³ C. Chen,⁴⁴ W. D. Hulsbergen,⁴⁴ A. Jawahery,⁴⁴ C. K. Lae,⁴⁴ D. A. Roberts,⁴⁴ G. Simi,⁴⁴ G. Blaylock,⁴⁵ C. Dallapiccola,⁴⁵ S. S. Hertzbach,⁴⁵ X. Li,⁴⁵ T. B. Moore,⁴⁵ S. Saremi,⁴⁵ H. Staengle,⁴⁵ R. Cowan,⁴⁶ G. Sciolla,⁴⁶ S. J. Sekula,⁴⁶ M. Spitznagel,⁴⁶ F. Taylor,⁴⁶ R. K. Yamamoto,⁴⁶ H. Kim,⁴⁷ S. E. McLachlin,⁴⁷ P. M. Patel,⁴⁷ S. H. Robertson,⁴⁷ A. Lazzaro,⁴⁸ V. Lombardo,⁴⁸ F. Palombo,⁴⁸ J. M. Bauer,⁴⁹ L. Cremaldi,⁴⁹ V. Eschenburg,⁴⁹ R. Godang,⁴⁹ R. Kroeger,⁴⁹ D. A. Sanders,⁴⁹ D. J. Summers,⁴⁹ H. W. Zhao,⁴⁹ S. Brunet,⁵⁰ D. Côté,⁵⁰ M. Simard,⁵⁰ P. Taras,⁵⁰ F. B. Viaud,⁵⁰ H. Nicholson,⁵¹ N. Cavallo,^{52,†} G. De Nardo,⁵² F. Fabozzi,^{52,†} C. Gatto,⁵² L. Lista,⁵² D. Monorchio,⁵² P. Paolucci,⁵² D. Piccolo,⁵² C. Sciacca,⁵² M. A. Baak,⁵³ G. Raven,⁵³ H. L. Snoek,⁵³ C. P. Jessop,⁵⁴ J. M. LoSecco,⁵⁴ G. Benelli,⁵⁵ L. A. Corwin,⁵⁵ K. K. Gan,⁵⁵ K. Honscheid,⁵⁵ D. Hufnagel,⁵⁵ P. D. Jackson,⁵⁵ H. Kagan,⁵⁵ R. Kass,⁵⁵ A. M. Rahimi,⁵⁵ J. J. Regensburger,⁵⁵ R. Ter-Antonyan,⁵⁵ Q. K. Wong,⁵⁵ N. L. Blount,⁵⁶ J. Brau,⁵⁶ R. Frey,⁵⁶

- O. Igonkina,⁵⁶ J. A. Kolb,⁵⁶ M. Lu,⁵⁶ C. T. Potter,⁵⁶ R. Rahmat,⁵⁶ N. B. Sinev,⁵⁶ D. Strom,⁵⁶ J. Strube,⁵⁶ E. Torrence,⁵⁶ A. Gaz,⁵⁷ M. Margoni,⁵⁷ M. Morandin,⁵⁷ A. Pompili,⁵⁷ M. Posocco,⁵⁷ M. Rotondo,⁵⁷ F. Simonetto,⁵⁷ R. Stroili,⁵⁷ C. Voci,⁵⁷ M. Benayoun,⁵⁸ H. Briand,⁵⁸ J. Chauveau,⁵⁸ P. David,⁵⁸ L. Del Buono,⁵⁸ Ch. de la Vaissière,⁵⁸ O. Hamon,⁵⁸ B. L. Hartfiel,⁵⁸ Ph. Leruste,⁵⁸ J. Malclès,⁵⁸ J. Ocariz,⁵⁸ L. Roos,⁵⁸ G. Therin,⁵⁸ L. Gladney,⁵⁹ M. Biasini,⁶⁰ R. Covarelli,⁶⁰ C. Angelini,⁶¹ G. Batignani,⁶¹ S. Bettarini,⁶¹ F. Bucci,⁶¹ G. Calderini,⁶¹ M. Carpinelli,⁶¹ R. Cenci,⁶¹ F. Forti,⁶¹ M. A. Giorgi,⁶¹ A. Lusiani,⁶¹ G. Marchiori,⁶¹ M. A. Mazur,⁶¹ M. Morganti,⁶¹ N. Neri,⁶¹ E. Paoloni,⁶¹ G. Rizzo,⁶¹ J. J. Walsh,⁶¹ M. Haire,⁶² D. Judd,⁶² D. E. Wagoner,⁶² J. Biesiada,⁶³ N. Danielson,⁶³ P. Elmer,⁶³ Y. P. Lau,⁶³ C. Lu,⁶³ J. Olsen,⁶³ A. J. S. Smith,⁶³ A. V. Telnov,⁶³ F. Bellini,⁶⁴ G. Cavoto,⁶⁴ A. D'Orazio,⁶⁴ D. del Re,⁶⁴ E. Di Marco,⁶⁴ R. Faccini,⁶⁴ F. Ferrarotto,⁶⁴ F. Ferroni,⁶⁴ M. Gaspero,⁶⁴ L. Li Gioi,⁶⁴ M. A. Mazzoni,⁶⁴ S. Morganti,⁶⁴ G. Piredda,⁶⁴ F. Polci,⁶⁴ F. Safai Tehrani,⁶⁴ C. Voena,⁶⁴ M. Ebert,⁶⁵ H. Schröder,⁶⁵ R. Waldi,⁶⁵ T. Adye,⁶⁶ B. Franek,⁶⁶ E. O. Olaiya,⁶⁶ S. Ricciardi,⁶⁶ F. F. Wilson,⁶⁶ R. Aleksan,⁶⁷ S. Emery,⁶⁷ A. Gaidot,⁶⁷ S. F. Ganzhur,⁶⁷ G. Hamel de Monchenault,⁶⁷ W. Kozanecki,⁶⁷ M. Legendre,⁶⁷ G. Vasseur,⁶⁷ Ch. Yèche,⁶⁷ M. Zito,⁶⁷ X. R. Chen,⁶⁸ H. Liu,⁶⁸ W. Park,⁶⁸ M. V. Purohit,⁶⁸ J. R. Wilson,⁶⁸ M. T. Allen,⁶⁹ D. Aston,⁶⁹ R. Bartoldus,⁶⁹ P. Bechtle,⁶⁹ N. Berger,⁶⁹ R. Claus,⁶⁹ J. P. Coleman,⁶⁹ M. R. Convery,⁶⁹ J. C. Dingfelder,⁶⁹ J. Dorfan,⁶⁹ G. P. Dubois-Felsmann,⁶⁹ D. Dujmic,⁶⁹ W. Dunwoodie,⁶⁹ R. C. Field,⁶⁹ T. Glanzman,⁶⁹ S. J. Gowdy,⁶⁹ M. T. Graham,⁶⁹ P. Grenier,⁶⁹ V. Halyo,⁶⁹ C. Hast,⁶⁹ T. Hry'ova,⁶⁹ W. R. Innes,⁶⁹ M. H. Kelsey,⁶⁹ P. Kim,⁶⁹ D. W. G. S. Leith,⁶⁹ S. Li,⁶⁹ S. Luitz,⁶⁹ V. Luth,⁶⁹ H. L. Lynch,⁶⁹ D. B. MacFarlane,⁶⁹ H. Marsiske,⁶⁹ R. Messner,⁶⁹ D. R. Muller,⁶⁹ C. P. O'Grady,⁶⁹ V. E. Ozcan,⁶⁹ A. Perazzo,⁶⁹ M. Perl,⁶⁹ T. Pulliam,⁶⁹ B. N. Ratcliff,⁶⁹ A. Roodman,⁶⁹ A. A. Salnikov,⁶⁹ R. H. Schindler,⁶⁹ J. Schwiening,⁶⁹ A. Snyder,⁶⁹ J. Stelzer,⁶⁹ D. Su,⁶⁹ M. K. Sullivan,⁶⁹ K. Suzuki,⁶⁹ S. K. Swain,⁶⁹ J. M. Thompson,⁶⁹ J. Va'vra,⁶⁹ N. van Bakel,⁶⁹ A. P. Wagner,⁶⁹ M. Weaver,⁶⁹ A. J. R. Weinstein,⁶⁹ W. J. Wisniewski,⁶⁹ M. Wittgen,⁶⁹ D. H. Wright,⁶⁹ H. W. Wulsin,⁶⁹ A. K. Yarritu,⁶⁹ K. Yi,⁶⁹ C. C. Young,⁶⁹ P. R. Burchat,⁷⁰ A. J. Edwards,⁷⁰ S. A. Majewski,⁷⁰ B. A. Petersen,⁷⁰ L. Wilden,⁷⁰ S. Ahmed,⁷¹ M. S. Alam,⁷¹ R. Bula,⁷¹ J. A. Ernst,⁷¹ V. Jain,⁷¹ B. Pan,⁷¹ M. A. Saeed,⁷¹ F. R. Wappler,⁷¹ S. B. Zain,⁷¹ W. Bugg,⁷² M. Krishnamurthy,⁷² S. M. Spanier,⁷² R. Eckmann,⁷³ J. L. Ritchie,⁷³ A. Satpathy,⁷³ C. J. Schilling,⁷³ R. F. Schwitters,⁷³ J. M. Izen,⁷⁴ X. C. Lou,⁷⁴ S. Ye,⁷⁴ F. Bianchi,⁷⁵ F. Gallo,⁷⁵ D. Gamba,⁷⁵ M. Bomben,⁷⁶ L. Bosisio,⁷⁶ C. Cartaro,⁷⁶ F. Cossutti,⁷⁶ G. Della Ricca,⁷⁶ S. Dittongo,⁷⁶ L. Lanceri,⁷⁶ L. Vitale,⁷⁶ V. Azzolini,⁷⁷ N. Lopez-March,⁷⁷ F. Martinez-Vidal,⁷⁷ Sw. Banerjee,⁷⁸ B. Bhuyan,⁷⁸ C. M. Brown,⁷⁸ D. Fortin,⁷⁸ K. Hamano,⁷⁸ R. Kowalewski,⁷⁸ I. M. Nugent,⁷⁸ J. M. Roney,⁷⁸ R. J. Sobie,⁷⁸ J. J. Back,⁷⁹ P. F. Harrison,⁷⁹ T. E. Latham,⁷⁹ G. B. Mohanty,⁷⁹ M. Pappagallo,^{79,‡} H. R. Band,⁸⁰ X. Chen,⁸⁰ B. Cheng,⁸⁰ S. Dasu,⁸⁰ M. Datta,⁸⁰ K. T. Flood,⁸⁰ J. J. Hollar,⁸⁰ P. E. Kutter,⁸⁰ B. Mellado,⁸⁰ A. Mihalyi,⁸⁰ Y. Pan,⁸⁰ M. Pierini,⁸⁰ R. Prepost,⁸⁰ S. L. Wu,⁸⁰ Z. Yu,⁸⁰ and H. Neal⁸¹

(The BABAR Collaboration)

¹Laboratoire de Physique des Particules, IN2P3/CNRS et Université de Savoie, F-74941 Annecy-Le-Vieux, France²Universitat de Barcelona, Facultat de Fisica, Departament ECM, E-08028 Barcelona, Spain³Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy⁴Institute of High Energy Physics, Beijing 100039, China⁵University of Bergen, Institute of Physics, N-5007 Bergen, Norway⁶Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA⁷University of Birmingham, Birmingham, B15 2TT, United Kingdom⁸Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany⁹University of Bristol, Bristol BS8 1TL, United Kingdom¹⁰University of British Columbia, Vancouver, British Columbia V6T 1Z1, Canada¹¹Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom¹²Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia¹³University of California at Irvine, Irvine, California 92697, USA¹⁴University of California at Los Angeles, Los Angeles, California 90024, USA¹⁵University of California at Riverside, Riverside, California 92521, USA¹⁶University of California at San Diego, La Jolla, California 92093, USA¹⁷University of California at Santa Barbara, Santa Barbara, California 93106, USA¹⁸University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA¹⁹California Institute of Technology, Pasadena, California 91125, USA²⁰University of Cincinnati, Cincinnati, Ohio 45221, USA²¹University of Colorado, Boulder, Colorado 80309, USA²²Colorado State University, Fort Collins, Colorado 80523, USA²³Universität Dortmund, Institut für Physik, D-44221 Dortmund, Germany

²⁴*Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany*²⁵*Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, F-91128 Palaiseau, France*²⁶*University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom*²⁷*Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy*²⁸*Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy*²⁹*Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy*³⁰*Harvard University, Cambridge, Massachusetts 02138, USA*³¹*Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany*³²*Imperial College London, London, SW7 2AZ, United Kingdom*³³*University of Iowa, Iowa City, Iowa 52242, USA*³⁴*Iowa State University, Ames, Iowa 50011-3160, USA*³⁵*Johns Hopkins University, Baltimore, Maryland 21218, USA*³⁶*Universität Karlsruhe, Institut für Experimentelle Kernphysik, D-76021 Karlsruhe, Germany*³⁷*Laboratoire de l'Accélérateur Linéaire, IN2P3/CNRS et Université Paris-Sud 11, Centre Scientifique d'Orsay, B. P. 34, F-91898 ORSAY Cedex, France*³⁸*Lawrence Livermore National Laboratory, Livermore, California 94550, USA*³⁹*University of Liverpool, Liverpool L69 7ZE, United Kingdom*⁴⁰*Queen Mary, University of London, E1 4NS, United Kingdom*⁴¹*University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom*⁴²*University of Louisville, Louisville, Kentucky 40292, USA*⁴³*University of Manchester, Manchester M13 9PL, United Kingdom*⁴⁴*University of Maryland, College Park, Maryland 20742, USA*⁴⁵*University of Massachusetts, Amherst, Massachusetts 01003, USA*⁴⁶*Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA*⁴⁷*McGill University, Montréal, Québec H3A 2T8, Canada*⁴⁸*Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy*⁴⁹*University of Mississippi, University, Mississippi 38677, USA*⁵⁰*Université de Montréal, Physique des Particules, Montréal, Québec H3C 3J7, Canada*⁵¹*Mount Holyoke College, South Hadley, Massachusetts 01075, USA*⁵²*Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy*⁵³*NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands*⁵⁴*University of Notre Dame, Notre Dame, Indiana 46556, USA*⁵⁵*Ohio State University, Columbus, Ohio 43210, USA*⁵⁶*University of Oregon, Eugene, Oregon 97403, USA*⁵⁷*Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy*⁵⁸*Laboratoire de Physique Nucléaire et de Hautes Energies, IN2P3/CNRS, Université Pierre et Marie Curie-Paris6, Université Denis Diderot-Paris7, F-75252 Paris, France*⁵⁹*University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA*⁶⁰*Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy*⁶¹*Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy*⁶²*Prairie View A&M University, Prairie View, Texas 77446, USA*⁶³*Princeton University, Princeton, New Jersey 08544, USA*⁶⁴*Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy*⁶⁵*Universität Rostock, D-18051 Rostock, Germany*⁶⁶*Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom*⁶⁷*DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France*⁶⁸*University of South Carolina, Columbia, South Carolina 29208, USA*⁶⁹*Stanford Linear Accelerator Center, Stanford, California 94309, USA*⁷⁰*Stanford University, Stanford, California 94305-4060, USA*⁷¹*State University of New York, Albany, New York 12222, USA*⁷²*University of Tennessee, Knoxville, Tennessee 37996, USA*⁷³*University of Texas at Austin, Austin, Texas 78712, USA*⁷⁴*University of Texas at Dallas, Richardson, Texas 75083, USA*⁷⁵*Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy*⁷⁶*Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy*⁷⁷*IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain*⁷⁸*University of Victoria, Victoria, British Columbia V8W 3P6, Canada*⁷⁹*Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom*⁸⁰*University of Wisconsin, Madison, Wisconsin 53706, USA*⁸¹*Yale University, New Haven, Connecticut 06511, USA*

(Received 19 July 2007; published 8 January 2008)

Using a data sample corresponding to an integrated luminosity of 342 fb^{-1} collected with the *BABAR* detector at the SLAC PEP-II electron-positron storage ring operating at a center-of-mass energy near 10.58 GeV, we measure $\mathcal{B}(\tau^- \rightarrow \pi^- \pi^- \pi^+ \nu_\tau (\text{ex. } K_S^0)) = (8.83 \pm 0.01 \pm 0.13)\%$, $\mathcal{B}(\tau^- \rightarrow K^- \pi^- \pi^+ \nu_\tau (\text{ex. } K_S^0)) = (0.273 \pm 0.002 \pm 0.009)\%$, $\mathcal{B}(\tau^- \rightarrow K^- \pi^- K^+ \nu_\tau) = (0.1346 \pm 0.0010 \pm 0.0036)\%$, and $\mathcal{B}(\tau^- \rightarrow K^- K^- K^+ \nu_\tau) = (1.58 \pm 0.13 \pm 0.12) \times 10^{-5}$, where the uncertainties are statistical and systematic, respectively. These include significant improvements over previous measurements and a first measurement of $\mathcal{B}(\tau^- \rightarrow K^- K^- K^+ \nu_\tau)$ in which no resonance structure is assumed. We also report a first measurement of $\mathcal{B}(\tau^- \rightarrow \phi \pi^- \nu_\tau) = (3.42 \pm 0.55 \pm 0.25) \times 10^{-5}$, a new measurement of $\mathcal{B}(\tau^- \rightarrow \phi K^- \nu_\tau) = (3.39 \pm 0.20 \pm 0.28) \times 10^{-5}$ and a first upper limit on $\mathcal{B}(\tau^- \rightarrow K^- K^- K^+ \nu_\tau (\text{ex. } \phi))$.

DOI: 10.1103/PhysRevLett.100.011801

PACS numbers: 13.35.Dx, 12.15.Hh, 14.40.Cs, 14.60.Fg

The weak interaction coupling strength between the first and second quark generations [1] can be probed with unprecedented precision in hadronic τ decays having net strangeness of unity in the final state using $e^+ e^- \rightarrow \tau^+ \tau^-$ data collected at the $e^+ e^-$ *B* factories. Inclusive measurements of the strange spectral function, obtained from τ lepton decays to final states containing kaons, provide a direct determination of the strange quark mass and Cabibbo-Kobayashi-Maskawa (CKM) matrix mixing element $|V_{us}|$ [2]. A significant improvement on the precision of $\mathcal{B}(\tau^- \rightarrow K^- \pi^- \pi^+ \nu_\tau)$ in particular, where one measurement [3] disagrees by more than 2 standard deviations from the others [4,5], will have the most immediate impact on the determination of these two fundamental Standard Model (SM) parameters using τ decays [6]. Measurements of $\tau^- \rightarrow \phi \pi^- \nu_\tau$ and $\tau^- \rightarrow \phi K^- \nu_\tau$ provide an interesting laboratory for studying Okubo-Zweig-Iizuka (OZI) suppression [7].

Significant improvements on measurements of $\mathcal{B}(\tau^- \rightarrow \pi^- \pi^- \pi^+ \nu_\tau)$, $\mathcal{B}(\tau^- \rightarrow K^- \pi^- \pi^+ \nu_\tau)$, and $\mathcal{B}(\tau^- \rightarrow K^- \pi^- K^+ \nu_\tau)$ are reported together with a first measurement of $\mathcal{B}(\tau^- \rightarrow K^- K^- K^+ \nu_\tau)$ without resonance assumptions (charge conjugate decays are implied). The data sample corresponds to an integrated luminosity of $\mathcal{L} = 342 \text{ fb}^{-1}$ recorded at a center-of-mass (c.m.) energy (\sqrt{s}) near 10.58 GeV using the *BABAR* detector at the SLAC PEP-II asymmetric-energy $e^+ e^-$ storage ring. With a luminosity-weighted average cross-section of $\sigma_{e^+ e^- \rightarrow \tau^+ \tau^-} = (0.919 \pm 0.003) \text{ nb}$ [8,9], this corresponds to the production of 3.16×10^8 τ -pair events.

The *BABAR* detector [10] has a silicon vertex tracker (SVT), drift chamber (DCH), ring-imaging Cherenkov detector (DIRC), and electromagnetic calorimeter (EMC) all contained in a 1.5-T solenoid. The iron flux return of the solenoid is instrumented to identify muons.

τ -pair events are simulated with higher-order radiative corrections using the KK2F Monte Carlo (MC) generator [8] with τ decays simulated with TAUOLA [11,12] using measured rates [13]. The detector response is simulated with GEANT4 [14]. Simulated events for signal as well as SM background processes [8,11,12,15,16] are reconstructed in the same manner as data. The MC samples are used for selection optimization and systematic error studies. The number of simulated nonsignal events is comparable to the

number expected in the data, with the exception of Bhabha and two-photon events, which are not simulated. Data studies demonstrate that these backgrounds are negligible.

The basic analysis strategy is to select a pure sample of $\tau^- \rightarrow h^- h^- h^+ \nu_\tau$ decays from $e^+ e^- \rightarrow \tau^+ \tau^-$ events by requiring the partner τ^+ to decay leptonically. Within this sample, each of the h^\pm mesons is uniquely identified as a charged pion or kaon and the decay categorized as $\tau^- \rightarrow \pi^- \pi^- \pi^+ \nu_\tau$, $\tau^- \rightarrow K^- \pi^- \pi^+ \nu_\tau$, $\tau^- \rightarrow K^- \pi^- K^+ \nu_\tau$, or $\tau^- \rightarrow K^- K^- K^+ \nu_\tau$. An efficiency migration matrix, \mathcal{E}_{ij} , initially obtained from MC simulations, is used to correct for efficiency losses from all stages of event selection and includes cross feed between the four signal channels where i (j) is the selected (true) decay mode index. The \mathcal{E}_{ij} matrix is modified using data control samples of kaons and pions from $D^{*+} \rightarrow \pi^+ D^0$, $D^0 \rightarrow \pi^+ K^-$ decays to account for small differences between MC calculations and data. The number of decay mode j signal events measured in the sample, $\mathbf{N}_j^{\text{Sig}}$, is then

$$\mathbf{N}_j^{\text{Sig}} = \sum_i (\mathcal{E}^{-1})_{ji} (\mathbf{N}_i^{\text{Data}} - \mathbf{N}_i^{\text{Bkg}}) \quad (1)$$

where $\mathbf{N}_i^{\text{Data}}$ is the number of data events selected in decay channel i and $\mathbf{N}_i^{\text{Bkg}}$ is the estimated number of background events in decay channel i arising from sources other than $\tau^- \rightarrow h^- h^- h^+ \nu_\tau$. The branching fraction for decay mode j is then $\mathcal{B}_j = \frac{\mathbf{N}_j^{\text{Sig}}}{2\mathcal{L}\sigma_{e^+ e^- \rightarrow \tau^+ \tau^-}}$.

The $\tau^- \rightarrow h^- h^- h^+ \nu_\tau$ sample is obtained by selecting events with four well reconstructed tracks having zero total charge, where none of the tracks originate from the conversion of photons in the material of the detector. All four tracks are required to lie within the geometrical acceptance of the EMC and DIRC and, to ensure the tracks can reach the DIRC, are required to have a transverse momentum of at least 250 MeV/c. If any two photons in the event are identified as coming from $\pi^0 \rightarrow \gamma\gamma$, the event is removed from the sample. The event is divided into hemispheres in the c.m. by a plane perpendicular to the thrust axis [17]. One of the hemispheres, the “leptonside,” is required to contain a $\tau \rightarrow e\nu\bar{\nu}$ or $\tau \rightarrow \mu\nu\bar{\nu}$ decay. If the hemisphere with an electron (muon) has any neutral particles with more than 1.0 (0.5) GeV, the event is not selected. The $e^+ e^- \rightarrow q\bar{q}$ backgrounds are further reduced to $\sim 0.1\%$ of

TABLE I. Diagonal elements of \mathcal{E}_{ij} excluding particle identification, numbers of selected, background, and signal events determined using Eq. (1). The branching-fraction results for this work are presented in the fifth line where the first error is statistical and the second systematic. The PDG average branching fractions or limit [13] are presented in the sixth line.

	$\tau^- \rightarrow \pi^- \pi^- \pi^+ \nu_\tau$	$\tau^- \rightarrow K^- \pi^- \pi^+ \nu_\tau$	$\tau^- \rightarrow K^- \pi^- K^+ \nu_\tau$	$\tau^- \rightarrow K^- K^- K^+ \nu_\tau$
ϵ	0.028	0.031	0.035	0.039
N^{Data}	$(1.5953 \pm 0.0013) \times 10^6$	$(6.956 \pm 0.0026) \times 10^4$	$(1.819 \pm 0.013) \times 10^4$	275 ± 17
N^{Bkg}	$(0.0642 \pm 0.0002) \times 10^6$	$(0.2263 \pm 0.0064) \times 10^4$	$(0.0145 \pm 0.0008) \times 10^4$	2.5 ± 1.5
N^{Sig}	$(55.59 \pm 0.05) \times 10^6$	$(171.5 \pm 1.2) \times 10^4$	$(84.71 \pm 0.66) \times 10^4$	$(9.93 \pm 0.84) \times 10^3$
\mathcal{B} (this work)	$(8.83 \pm 0.01 \pm 0.13)\%$	$(0.273 \pm 0.002 \pm 0.009)\%$	$(0.1346 \pm 0.0010 \pm 0.0036)\%$	$(1.58 \pm 0.13 \pm 0.12) \times 10^{-5}$
\mathcal{B} (PDG average)	$(9.13 \pm 0.05 \pm 0.46)\%$ [4]	$(0.33 \pm 0.05)\%$	$(0.154 \pm 0.009)\%$	$<3.7 \times 10^{-5}$ @ 90% C.L.

the $\tau^- \rightarrow h^- h^- h^+ \nu_\tau$ sample by requiring a thrust magnitude >0.85 .

Backgrounds from Bhabha and $e^+ e^- \rightarrow \mu^+ \mu^-$ events with photon conversions are suppressed by requiring the momentum of the leptonside track to be less than 80% of $\sqrt{s}/2$ and a requirement that none of the three signalside tracks pass an electron identification algorithm. Remaining non- τ background events, including those from two-photon processes, are reduced by requiring the event missing c.m. transverse momentum to be $>0.009\sqrt{s}$.

The remaining background in the sample is predominantly from other τ decays containing π^0 and K_S^0 mesons. Events containing a K_S^0 are identified and removed. Residual backgrounds from decays having a π^0 are reduced by requiring the total energy in the signalside hemisphere deposited in the EMC which is unassociated with the three charged hadron tracks to be <200 MeV. With these requirements, contributions from $\tau^- \rightarrow h^- n \pi^0 \nu$ and $\tau^- \rightarrow h^- h^- h^+ 2\pi^0 \nu$ are negligible.

A track in the $\tau^- \rightarrow h^- h^- h^+ \nu_\tau$ sample is classified as a kaon using a likelihood approach to combine information from the DIRC, DCH, and SVT with a characteristic kaon identification efficiency of $\sim 80\%$ and pion misidentification rate of $\sim 1\%$. If the track fails to be identified as a kaon, it is classified as a pion. Events are selected if the signalside decays are identified as $\tau^- \rightarrow \pi^- \pi^- \pi^+ \nu_\tau$, $\tau^- \rightarrow K^- \pi^- \pi^+ \nu_\tau$, $\tau^- \rightarrow K^- \pi^- K^+ \nu_\tau$, or $\tau^- \rightarrow K^- K^- K^+ \nu_\tau$ with decays having a wrong charge combination removed.

The diagonal elements of \mathcal{E}_{ij} excluding particle identification (ϵ), numbers of selected, background and signal events determined using Eq. (1) are shown in Table I for all modes. The increase in ϵ with number of kaons in the decay is associated primarily with the transverse momentum requirement and the K_S^0 veto. The background fraction from τ decays to $\pi^- \pi^- \pi^+ \pi^0 \nu$ ($K^- \pi^- \pi^+ \pi^0 \nu$, $K^- \pi^- K^+ \pi^0 \nu$, $K^- K^- K^+ \pi^0 \nu$) in the $\pi^- \pi^- \pi^+$ ($K^- \pi^- \pi^+$, $K^- \pi^- K^+$, $K^- K^- K^+$) candidate sample is estimated to be $(3.6 \pm 0.3)\%$ ($(2.3 \pm 0.4)\%$, $(0.4 \pm 0.1)\%$, $<5.0\%$). Non- τ backgrounds comprise less than 0.5% of each channel's final event sample.

The component of \mathcal{E}_{ij} associated with the particle identification, M_{ij} , is shown in the first four rows of Table II.

Note that this matrix includes efficiency losses associated with cross feed of the wrong charge combinations and small factors associated with data control sample corrections to the MC cross-feed efficiencies; therefore, the columns of the table are not expected to sum to 100%.

Systematic uncertainties are assigned for the following: luminosity; cross section; migration matrix elements, which includes MC statistical and systematic errors associated with the efficiency and particle identification; signal-mode modeling; modeling of the EMC and tracking response including scale and resolution uncertainties, the sensitivities of the measurements to the modeling of hadronic and electromagnetic showers in the EMC, and tracking efficiency; modeling of the trigger; and modeling of the backgrounds, including uncertainties on cross-sections and branching fractions. These are summarized in Table III along with the total error correlation matrix. The absolute normalization is a significant source of the correlations.

The branching-fraction results of this analysis are presented in Table I together with the world average values or limit published by the Particle Data Group (PDG) [13], with which they are consistent and significantly more precise. In all four channels, the results where the leptonside has an electron are consistent with those where it has a muon. Our measurement of $\mathcal{B}(\tau^- \rightarrow \pi^- \pi^- \pi^+ \nu_\tau)$ is also consistent with a precision $\mathcal{B}(\tau^- \rightarrow \pi^- \pi^- \pi^+ (\text{ex.}\omega))$ measurement [18] after accounting for the ω . Our measurement of $\mathcal{B}(\tau^- \rightarrow K^- \pi^- \pi^+ \nu_\tau)$ is in agreement with [3] and disagrees by more than 2 standard deviations from

TABLE II. Component of the efficiency migration matrix associated with pion-kaon particle identification, M_{ij} , in percent. The $\tau^- \rightarrow \phi \pi^- \nu_\tau$ and $\tau^- \rightarrow \phi K^- \nu_\tau$ study employs a higher-efficiency-lower-purity kaon selection algorithm.

Candidates	Decay modes			
	$\pi\pi\pi$	$K\pi\pi$	$K\pi K$	KKK
$\pi\pi\pi$	97.40	22.49	4.73	1.02
$K\pi\pi$	1.42	74.87	16.43	6.38
$K\pi K$	0.01	0.49	59.63	25.54
KKK	0.26	50.87
$\phi\pi$			72.54	19.20
ϕK			0.83	66.06

TABLE III. Upper: Systematic uncertainties (%). Lower: Correlation matrix from stat. \oplus syst. covariance matrix.

	$\pi\pi\pi$	$K\pi\pi$	$K\pi K$	KKK
\mathcal{L}	0.9	0.9	0.9	0.9
$\sigma_{e^+e^- \rightarrow \tau^+\tau^-}$	0.3	0.3	0.3	0.3
M_{ij} and particle ID	0.4	3.0	1.9	4.9
signal modeling	0.2	0.2	1.3	2.0
EMC and DCH response	0.8	0.9	0.8	1.2
trigger	0.1	0.1	0.1	0.1
backgrounds	0.4	0.7	0.4	5.5
Total	1.4	3.4	2.7	7.8
$\pi\pi\pi$		0.544	0.390	0.031
$K\pi\pi$			0.177	0.093
$K\pi K$				0.087

[4,5]. We report a first measurement of $\mathcal{B}(\tau^- \rightarrow K^- K^- K^+ \nu_\tau)$ in which no resonance structure is assumed and which has a significance in excess of 8σ .

A $\phi(1020)$ contribution is seen in both the $\tau^- \rightarrow K^- \pi^- K^+ \nu_\tau$ and $\tau^- \rightarrow K^- K^- K^+ \nu_\tau$ decay modes. The use of a kaon selection algorithm with higher efficiency, but less purity, provides significantly higher signal-to-background for $\tau^- \rightarrow \phi K^- \nu_\tau$. The $\tau^- \rightarrow \phi \pi^- \nu_\tau$ ($\tau^- \rightarrow \phi K^- \nu_\tau$) signal has a 5.7σ (9.8σ) level of significance. The last two rows of Table II lists the M_{ij} matrix for this higher efficiency selection. The $K^+ K^-$ invariant mass distributions for the $\tau^- \rightarrow K^- \pi^- K^+ \nu_\tau$ ($\tau^- \rightarrow K^- K^- K^+ \nu_\tau$) mode using this sample is shown in Fig. 1 and 2. Below 1.09 GeV (1.15 GeV), after background subtraction of the non- $K^+ K^-$ events, the $K^+ K^-$ invariant mass distribution




FIG. 1 (color online). $K^+ K^-$ invariant mass in $\tau^- \rightarrow K^- \pi^- K^+ \nu_\tau$ decays. Data are represented by points with error bars, MC of $\tau^- \rightarrow K^- \pi^- K^+ \nu_\tau$ by the open histogram, cross feed from other $\tau^- \rightarrow h^- h^- h^+ \nu_\tau$ channels by the light shaded histogram, and non- τ backgrounds by the dark shaded histogram. The inset shows the background-subtracted data with the fit (solid line) and nonresonant component (dashed line).

from the $\tau^- \rightarrow \phi \pi^- \nu_\tau$ ($\tau^- \rightarrow \phi K^- \nu_\tau$) decay is well described by a Breit-Wigner function convoluted with a Gaussian resolution function for the signal and a third-order polynomial (function in [19]) for the background and is used to fit for the number of events. A binned maximum likelihood fit yields 344 ± 42 (274 ± 16) $\tau^- \rightarrow \phi \pi^- \nu_\tau$ ($\tau^- \rightarrow \phi K^- \nu_\tau$) candidates. MC estimates of the subtracted $q\bar{q}$ background events with a ϕ contribute a 5.2% (0.7%) uncertainty. The $\phi \pi^-$ candidate sample has an additional 4.3% uncertainty arising from potentially peaking $\tau^- \rightarrow \phi \pi^- n\pi^0 \nu_\tau$ background. The fit parameterization contributes a 1.0% (2.0%) error. Accounting for $\mathcal{B}(\phi \rightarrow K^+ K^-)$ [13], $\mathcal{B}(\tau^- \rightarrow \phi \pi^- \nu_\tau) = (3.42 \pm 0.55 \pm 0.25) \times 10^{-5}$ and $\mathcal{B}(\tau^- \rightarrow \phi K^- \nu_\tau) = (3.39 \pm 0.20 \pm 0.28) \times 10^{-5}$ with a correlation of -0.07 . From the fit, we find no evidence for $\tau^- \rightarrow K^- K^- K^+ \nu_\tau$ without a ϕ and set a first upper limit on $\mathcal{B}(\tau^- \rightarrow K^- K^- K^+ \nu_\tau \text{ (ex. } \phi)) < 2.5 \times 10^{-6}$ at 90% C.L.

This is the first measurement of $\mathcal{B}(\tau^- \rightarrow \phi \pi^- \nu_\tau)$. It is consistent with a CLEO limit [20] and larger than the $(1.20 \pm 0.28) \times 10^{-5}$ value predicted by a vector meson dominance model [7]. Our $\mathcal{B}(\tau^- \rightarrow \phi K^- \nu_\tau)$ measurement is consistent with a recent Belle result [21]. Recent calculations using a meson dominance model [22] agree with our $\mathcal{B}(\tau^- \rightarrow \phi \pi^- \nu_\tau)$ and $\mathcal{B}(\tau^- \rightarrow \phi K^- \nu_\tau)$ measurements and with the ratio $\frac{\mathcal{B}(\tau^- \rightarrow \phi \pi^- \nu_\tau)}{\mathcal{B}(\tau^- \rightarrow \phi K^- \nu_\tau)} = 0.99 \pm 0.21$. Our measurements of $\mathcal{B}(\tau^- \rightarrow K^- \pi^- \pi^+ \nu_\tau)$ and $\mathcal{B}(\tau^- \rightarrow K^- K^- K^+ \nu_\tau)$, when combined with improved measurements of the other strange decays, will constrain the CKM element $|V_{us}|$ better than unitarity bounds [6,23].




FIG. 2 (color online). $K^+ K^-$ invariant mass in $\tau^- \rightarrow K^- K^- K^+ \nu_\tau$ decays with two entries per event. Data are represented by points with error bars, the open histogram is the MC calculation of $\tau^- \rightarrow \phi K^- \nu_\tau$, the light shaded histogram is the cross feed from the other $\tau^- \rightarrow h^- h^- h^+ \nu_\tau$ channels, primarily from $\tau^- \rightarrow K^- \pi^- K^+ \nu_\tau$, and non- τ backgrounds by the dark shaded histogram. The inset shows the background-subtracted data with the fit (solid line) and nonresonant component (dashed line).

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MIST (Russia), MEC (Spain), and PPARC (United Kingdom). Individuals have received support from the Marie Curie EIF (European Union) and the A. P. Sloan Foundation.

*Also with Università di Perugia, Dipartimento di Fisica, Perugia, Italy.

[†]Also with Università della Basilicata, Potenza, Italy.

[‡]Also with IPPP, Physics Department, Durham University, Durham DH1 3LE, United Kingdom.

- [1] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963).
- [2] E. Gamiz, M. Jamin, A. Pich, J. Prades, and F. Schwab, Phys. Rev. Lett. **94**, 011803 (2005); S. Chen, M. Davier, E. Gamiz, A. Hocker, A. Pich, and J. Prades, Eur. Phys. J. C **22**, 31 (2001); J. Kambor and K. Maltman, Phys. Rev. D **62**, 093023 (2000); A. Pich and J. Prades, J. High Energy Phys. 10 (1999) 004; S. Narison, Phys. Lett. B **466**, 345 (1999); K. Maltman, Phys. Rev. D **58**, 093015 (1998).
- [3] R. Barate *et al.* (ALEPH Collaboration), Eur. Phys. J. C **1**, 65 (1998).
- [4] R. A. Briere *et al.* (CLEO Collaboration), Phys. Rev. Lett. **90**, 181802 (2003).
- [5] G. Abbiendi *et al.* (OPAL Collaboration), Eur. Phys. J. C **35**, 437 (2004).

- [6] K. Maltman and C. E. Wolfe, Phys. Lett. B **639**, 283 (2006).
- [7] G. Lopez Castro and D. A. Lopez Falcon, Phys. Rev. D **54**, 4400 (1996).
- [8] B. F. Ward, S. Jadach, and Z. Was, Nucl. Phys. B, Proc. Suppl. **116**, 73 (2003).
- [9] S. Banerjee, B. Pietrzyk, J. M. Roney, and Z. Was, arXiv:hep-ph0706.3235.
- [10] B. Aubert *et al.* (*BABAR* Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **479**, 1 (2002).
- [11] S. Jadach, Z. Was, R. Decker, and J. H. Kühn, Comput. Phys. Commun. **76**, 361 (1993).
- [12] E. Barberio and Z. Was, Comput. Phys. Commun. **79**, 291 (1994).
- [13] W.-M. Yao *et al.* (Particle Data Group), J. Phys. G **33**, 1 (2006).
- [14] S. Agostinelli *et al.* (GEANT4 Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **506**, 250 (2003).
- [15] D. J. Lange, Nucl. Instrum. Methods Phys. Res., Sect. A **462**, 152 (2001).
- [16] T. Sjöstrand, Comput. Phys. Commun. **82**, 74 (1994).
- [17] S. Brandt *et al.*, Phys. Lett. **12**, 57 (1964); E. Farhi, Phys. Rev. Lett. **39**, 1587 (1977).
- [18] S. Schael *et al.* (ALEPH Collaboration), Phys. Rep. **421**, 191 (2005).
- [19] H. Albrecht *et al.* (ARGUS Collaboration), Phys. Lett. B **241**, 278 (1990). $f(m) = [m - (2m_K)]e^{b[m^2 - (2m_K)^2]}[m^2 - (2m_K)^2]^a$, $m > 2m_K$; $= 0$, $m < 2m_K$.
- [20] P. Avery *et al.* (CLEO Collaboration), Phys. Rev. D **55**, R1119 (1997).
- [21] K. Abe *et al.* (Belle Collaboration), Phys. Lett. B **643**, 5 (2006).
- [22] A. Flores-Tlalpa and G. Lopez-Castro, arXiv:hep-ph0709.4039.
- [23] R. J. Hill, Phys. Rev. D **74**, 096006 (2006).