

A standard, single dose of inhaled terbutaline attenuates hyperpnea-induced bronchoconstriction and mast cell activation in athletes

A. J. Simpson,¹ J. R. Bood,^{2,3,4} S. D. Anderson,⁵ L. M. Romer,¹ B. Dahlén,^{3,4} S.-E. Dahlén,^{2,4}
and P. Kippelen¹

¹Centre for Human Performance, Exercise and Rehabilitation, College of Health and Life Sciences, Brunel University London, United Kingdom; ²Unit for Experimental Asthma Research, Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden; ³Unit for Clinical Asthma Research, Department of Internal Medicine, Karolinska University Hospital Huddinge, Stockholm, Sweden; ⁴Centre for Allergy Research, Karolinska Institutet, Stockholm, Sweden; and ⁵Department of Respiratory and Sleep Medicine, Royal Prince Alfred Hospital, Sydney, Australia

Submitted 19 August 2015; accepted in final form 28 January 2016

Simpson AJ, Bood JR, Anderson SD, Romer LM, Dahlén B, Dahlén S-E, Kippelen P. A standard, single dose of inhaled terbutaline attenuates hyperpnea-induced bronchoconstriction and mast cell activation in athletes. *J Appl Physiol* 120: 1011–1017, 2016. First published February 4, 2016; doi:10.1152/jappphysiol.00700.2015.—Release of bronchoactive mediators from mast cells during exercise hyperpnea is a key factor in the pathophysiology of exercise-induced bronchoconstriction (EIB). Our aim was to investigate the effect of a standard, single dose of an inhaled β_2 -adrenoceptor agonist on mast cell activation in response to dry air hyperpnea in athletes with EIB. Twenty-seven athletes with EIB completed a randomized, double-blind, placebo-controlled, crossover study. Terbutaline (0.5 mg) or placebo was inhaled 15 min prior to 8 min of eucapnic voluntary hyperpnea (EVH) with dry air. Pre- and postbronchial challenge, urine samples were analyzed by enzyme immunoassay for 11β -prostaglandin $F_{2\alpha}$ (11β -PGF_{2 α}). The maximum fall in forced expiratory volume in 1 s of 14 (12–20)% (median and interquartile range) following placebo was attenuated to 7 (5–9)% with the administration of terbutaline ($P < 0.001$). EVH caused a significant increase in 11β -PGF_{2 α} from 41 (27–57) ng/mmol creatinine at baseline to 58 (43–72) ng/mmol creatinine at its peak post-EVH following placebo ($P = 0.002$). The rise in 11β -PGF_{2 α} was inhibited with administration of terbutaline: 39 (28–44) ng/mmol creatinine at baseline vs. 40 (33–58) ng/mmol creatinine at its peak post-EVH ($P = 0.118$). These data provide novel in vivo evidence of mast cell stabilization following inhalation of a standard dose of terbutaline prior to bronchial provocation with EVH in athletes with EIB.

inhaled β_2 -agonist; exercise-induced bronchoconstriction; eucapnic voluntary hyperpnea; prostaglandin D₂

NEW & NOTEWORTHY

This study provides the first in vivo evidence for a mast cell stabilizing effect of the short-acting inhaled β_2 -adrenoceptor agonist terbutaline, when administered prophylactically at a clinically recommended dose (0.5 mg) before bronchial provocation with dry air. Our data therefore support the proposal that β_2 -adrenoceptor agonist-mediated mast cell stabilization is a major contributor to bronchoprotection in individuals with exercise-induced bronchoconstriction.

EXERCISE-INDUCED BRONCHOCONSTRICTION (EIB) is the transient narrowing of airways that occurs during or shortly after stren-

uous exercise. The majority of individuals with asthma experience EIB (33), and EIB is also prevalent in children (7, 29) and elite athletes (17). The Global Initiative for Asthma guidelines suggest that inhaled short-acting β_2 -adrenoceptor agonists should be prescribed to all individuals with asthma for use “as needed reliever” and recommends their use for the short-term prevention of EIB (27). A recent Cochrane review of 45 studies revealed that inhaled short-acting β_2 -adrenoceptor agonists, when taken prophylactically, reduce postexercise bronchoconstriction by 66% in individuals with EIB (9). The primary mechanism of this protection is believed to be functional antagonism, whereby β_2 -adrenoceptor-induced relaxation of the bronchial smooth muscle opposes the contractile effects of the various mediators of bronchoconstriction (2).

In 1975 and 1976, Anderson and colleagues (4, 5) reported that β_2 -adrenoceptor agonists given by inhalation in low doses were superior to tablets (given in much higher doses) in preventing EIB, even though both formulations induced bronchodilatation. Those authors therefore proposed that the aerosol had an additional site of action to the smooth muscle and prevented EIB by delivering a concentration of drug sufficient to stabilize mast cells in the airways and inhibit the release of “bronchoconstrictor substances.”

Evidence of mast cell activation associated with exercise and hyperpnea of dry air is now well established by the finding of an increase in urinary excretion of 11β -prostaglandin $F_{2\alpha}$ (11β -PGF_{2 α}), i.e., a main metabolite of prostaglandin (PG) D₂, which is produced almost exclusively from mast cells (31, 34, 35, 37, 38, 46). The proposal that inhalation of β_2 -adrenoceptor agonists may prevent EIB by inhibiting release of mast cell mediators is supported by in vitro observations of an inhibition of IgE-dependent release of PGD₂ and histamine from mast cells with the short-acting β_2 -adrenoceptor agonist salbutamol (19). In vivo, the long-acting β_2 -adrenoceptor agonist salmeterol has been shown to inhibit mast cell mediator release following provocation with increasing doses of lysine-aspirin in patients with aspirin-induced asthma (54). Further, a large dose of the long-acting β_2 -adrenoceptor agonist formoterol has been shown to reduce the release of 11β -PGF_{2 α} following bronchial provocation with dry powder of mannitol in patients with asthma (15). Whether a single, standard dose of short-acting β_2 -adrenoceptor agonist—the mainstay treatment for EIB prevention—has a similar stabilizing effect on mast cells during EIB remains to be established.

The aim of this investigation was to test the efficacy of a single, clinically recommended dose of an inhaled short-acting

Address for reprint requests and other correspondence: P. Kippelen, College of Health and Life Sciences, Division of Sport, Health and Exercise Sciences, Brunel Univ. London, Uxbridge, Middlesex, UB8 3PH, U.K. (e-mail: pascale.kippelen@brunel.ac.uk).

β_2 -adrenoceptor agonist at inhibiting mast cell mediator release following hyperpnea with dry air. Our hypothesis was that premedication with 0.5 mg of terbutaline would attenuate the increase in urinary excretion of 11β -PGF_{2 α} following 8 min of eucapnic voluntary hyperpnea (EVH) in athletes with EIB.

METHODS

Subjects. Part of the current methodology has been published elsewhere to address a separate research question (53). Twenty-seven athletes with EIB completed the study. EIB was determined by a fall of $\geq 10\%$ in forced expiratory volume in 1 s (FEV₁) following an 8-min EVH challenge during a screening visit. Participants were nonsmokers, were free from any respiratory infections for 4 wk prior to the study, and had no known chronic medical condition other than asthma and/or EIB. Alcohol, caffeine, and exercise were withheld on the day of testing, and medications were withheld as follows: short-acting β_2 -agonist treatments for a minimum of 8 h, long-acting β_2 -agonist treatments for 24 h, inhaled corticosteroid treatments for 12 h, combination therapies of long-acting β_2 -agonist plus inhaled corticosteroid treatments for 24 h, and nonsteroidal anti-inflammatory medication for 7 days. Participants provided written informed consent after the study protocol, and potential risks were explained. The study was approved by the United Kingdom National Health Service Research Ethics Committee (NHS REC reference number 10/H0716/30).

Experimental design. The study used a randomized, double-blind, placebo-controlled, crossover experimental design with two experimental visits. The experimental visits were completed within 3 wk and separated by at least 2 days. Urine samples were collected at baseline and at 30 and 60 min following EVH with dry air on two separate days: one following treatment with 0.5 mg of terbutaline and one after administration of a placebo. Urine samples were analyzed for 11β -PGF_{2 α} . The primary end points were the change in the concentration of urinary 11β -PGF_{2 α} and the maximum percent fall in FEV₁ following EVH.

To standardize for fluctuations in lung function throughout the day (42), all experimental visits commenced between 8.00 and 11.00 a.m. Upon arrival, participants performed spirometry according to American Thoracic Society/European Respiratory Society guidelines (40). A single 0.5 mg dose of terbutaline was then administered via a dry powder inhaler (Bricanyl Turbohaler, Astra Zeneca, London, U.K.). An empty demonstration Turbohaler was used for administration of the placebo. The active drug (or placebo) was administered by one deep, hard inhalation through the inhaler, followed by a 10-s breath hold. Spirometry was repeated 10 min posttreatment.

The EVH challenge began 15 min after treatment administration. The test consisted of 8 min of dry air hyperpnea at a target ventilation of 85% predicted maximum voluntary ventilation (MVV; calculated as $30 \times$ baseline FEV₁) (1), with the ventilation achieved during the first visit used as the target ventilation for the second visit. The test was performed on a commercially available system (Eucapsys, SMTEC, Nyon, Switzerland) that delivered a dry gas mixture containing 5% CO₂, 20% O₂, and balance N₂. Subjects recovered spontaneously from the EVH challenge, and spirometry was repeated in duplicate at 2, 5, 10, 15, 20, 30, and 60 min postchallenge. The highest of two repeatable FEV₁ values was kept for analysis. The maximum fall in FEV₁ was expressed as a percentage from the posttreatment value. The degree of bronchoprotection afforded by terbutaline was calculated by subtracting the maximum percent fall in FEV₁ on the drug treatment day from the maximum percent fall in FEV₁ on the placebo day, and expressing it as a percentage of the placebo.

The atopic status of the participants was determined by a standard skin prick test (12) conducted 40 min post-EVH during the first experimental visit. The following allergens were tested: cat hair, timothy grass, and house dust mite (ALK-Abello, Reading, U.K.). Histamine and saline were used as positive and negative

controls, respectively. A test was deemed positive if the reaction wheal was ≥ 3 mm.

Ingestion of water was standardized throughout the experimental visits. One hour prior to the visits, subjects were asked to drink 200 ml of water. They were given a further 400 ml upon arrival at the laboratory, and then 200 ml every 30 min. Two baseline urine samples were collected: the first on arrival at the laboratory, which was discarded, and the second 30–60 min later and immediately before administration of the drug, which was used as the baseline. Following the EVH challenge, further urine samples were collected at 30 and 60 min. The urine samples were stored without preservatives at -80°C . The urine samples were analyzed for 11β -PGF_{2 α} using commercially available EIA reagents (Cayman Chemical, Ann Arbor, MI) as described elsewhere (10). Urinary data on excretion of 11β -PGF_{2 α} were normalized in relation to excretion of creatinine using the modified Jaffe colorimetric method [as done previously (10)] and expressed as nanograms of excreted mediator per micromole of creatinine.

Statistics. Sample size was based on a previous study in which the mast cell stabilizing effect of sodium cromoglycate was investigated during EVH in athletes with EIB (34). With a risk alpha of 5%, a risk beta of 95%, and using the effect size from Kippelen and colleagues (34), analysis of 16 urine samples was required [G*Power3 software (26)].

Data were tested for normality using the Shapiro-Wilk test. Baseline FEV₁ (l), changes in FEV₁ following treatment (%), ventilation during EVH (l), level of bronchoprotection (%), and stature of the participants were normally distributed and data are presented as means \pm SD. Differences between conditions for these variables were analyzed using paired sample *t*-tests. All other data were nonnormally distributed and therefore presented as median and interquartile range (Q1–Q3). Differences between conditions for maximum change in FEV₁ (%) and urinary 11β -PGF_{2 α} concentrations postchallenge were analyzed using Wilcoxon signed-rank tests. Differences across time were analyzed using Friedman two-way analysis of variance by ranks. In cases of statistical significance, Wilcoxon signed-rank tests were used to identify where differences occurred. The relationship between the change in FEV₁ (%) and the change in urinary 11β -PGF_{2 α} following EVH was initially assessed via Spearman correlation tests, in each experimental condition, separately; this was then followed by a multiple regression analysis for within-subject repeated measures (8). All statistical analyses were conducted using SPSS 20 (Chicago, IL). The level of significance was set at $P < 0.05$.

RESULTS

Participant characteristics. Age, stature, and body mass of the participants were 23 (19–32) yr, 174 ± 8 cm, and 71 (67–79) kg, respectively. Participants were involved in the following sports: athletics ($n = 13$), cycling ($n = 1$), football ($n = 4$), rugby ($n = 1$), cricket ($n = 1$), netball ($n = 1$), and rowing ($n = 6$). They were training for 8 (5–10) h/wk in aerobic activities, and had 8 (5–12) yr experience in their sport. Medical diagnosis, current treatment, and atopic status are presented in Table 1.

Baseline lung function and ventilation during EVH. Baseline FEV₁ was not significantly different between conditions: 3.68 ± 0.65 and 3.65 ± 0.64 liters in the placebo and terbutaline conditions, respectively ($P = 0.129$). Terbutaline had a small, but statistically significant ($P < 0.001$) bronchodilator effect: FEV₁ increased by $5 \pm 3\%$ with administration of terbutaline. No such bronchodilator effect was noticed after the administration of placebo. Ventilation was slightly but significantly ($P = 0.047$) higher with terbutaline: 102 ± 20 vs. 101 ± 20 l/min (78 ± 7 vs. $80 \pm 8\%$ of predicted MVV).

Table 1. Participant characteristics

| ID | Gender | Atopic Status | Baseline FEV ₁ , % Predicted | Previous Diagnosis | Treatment | Prescribed ICS Dose, µg/day |
|-----|--------|---------------|---|--------------------|-------------------|-----------------------------|
| 1 | M | + | 95 | Asthma+EIB | Combination, SABA | 1,600 |
| 2 | M | + | 92 | Asthma | SABA, ICS | 200 |
| 3 | M | + | 99 | Asthma+EIB | Combination | 400 |
| 4 | F | + | 108 | EIB | SABA, ICS | 200 |
| 5 | M | + | 97 | — | — | — |
| 6 | F | + | 92 | Asthma | SABA, ICS | 200 |
| 7 | M | + | 93 | Asthma+EIB | SABA | — |
| 8 | F | + | 93 | Asthma | Combination | 400 |
| 9 | M | + | 99 | Asthma | SABA, ICS | 600 |
| 10 | M | — | 99 | — | — | — |
| 11 | M | + | 85 | Asthma | — | — |
| 12 | F | + | 103 | Asthma | — | — |
| 13 | M | + | 109 | EIB | SABA | — |
| 14 | F | — | 99 | Asthma+EIB | Combination | 400 |
| 15 | F | — | 109 | — | — | — |
| 16 | M | + | 94 | Asthma | SABA | — |
| 17 | F | — | 96 | Asthma | SABA | — |
| 18 | M | + | 94 | Asthma | SABA | — |
| 19 | M | + | 111 | Asthma | SABA, ICS | 200 |
| 20 | F | + | 102 | EIB | SABA, ICS | 200 |
| 21 | F | + | 86 | Asthma | SABA | — |
| 22 | F | — | 88 | EIB | — | — |
| 23 | F | + | 82 | Asthma+EIB | SABA | — |
| 24 | M | + | 91 | Asthma | SABA | — |
| 25* | M | + | 102 | Asthma | SABA | — |
| 26* | M | — | 85 | Asthma | Combination | 200 |
| 27* | M | + | 109 | Asthma+EIB | ICS | 100 |

Atopic status: +, positive skin prick response to house dust mite, timothy grass, and/or cat hair; FEV₁ % predicted, baseline forced expiratory volume in 1 s expressed relative to the predicted value (50); EIB, exercise-induced bronchoconstriction; SABA, short-acting β_2 -agonist; LABA, long-acting β_2 -agonist; ICS, inhaled corticosteroids; combination, combination therapy of LABA and ICS; *participants who had urinary 11 β -PGF_{2 α} concentrations below the level of detection.

Airway response to EVH. Terbutaline significantly inhibited the airway response to EVH. The maximum fall in FEV₁ was reduced from 14 (12–20)% in the placebo condition to 7 (5–9)% with the administration of terbutaline ($P < 0.001$). The degree of bronchoprotection afforded by terbutaline was $60 \pm 30\%$, with a range of 0–94%. Terbutaline afforded complete bronchoprotection (<10% fall in FEV₁ post-EVH) to 22 of the 27 athletes (81%; Fig. 1). The five athletes who did not receive complete bronchoprotection did not differ from the main cohort in regard to their previous medical diagnosis (i.e., asthma with EIB, asthma only, EIB only, or no previous asthma/EIB diagnosis) or their current asthma treatment (i.e., inhaled β_2 -adrenoceptor agonists, inhaled corticosteroids, or no treatment).

Urinary 11 β -PGF_{2 α} . Three participants were excluded from statistical analysis because their 11 β -PGF_{2 α} concentrations were below the level of detection. Of the 24 remaining participants, 18 had complete data sets (i.e., 11 β -PGF_{2 α} was detectable in all samples), and their data were used to determine the kinetics of urinary 11 β -PGF_{2 α} excretion post-EVH. An additional six participants had acceptable baseline values and at least one acceptable 11 β -PGF_{2 α} value post-EVH and were therefore included in the analysis of peak urinary 11 β -PGF_{2 α} release post-EVH.

Kinetics of 11 β -PGF_{2 α} excretion post-EVH. There was no difference in baseline urinary 11 β -PGF_{2 α} values between conditions ($n = 18$; $P = 0.446$). A significant time effect was noted in the placebo condition ($P = 0.002$), with an increase in urinary 11 β -PGF_{2 α} from 45 (35–71) ng/mmol creatinine at baseline to 58 (41–78) ng/mmol creatinine at 30 min postchallenge ($P = 0.025$; Fig. 2). Terbutaline completely inhibited the

rise in urinary 11 β -PGF_{2 α} post-EVH ($P = 0.446$ vs. baseline). Consequently, urinary 11 β -PGF_{2 α} levels were higher in the placebo vs. the terbutaline condition at 30 min ($P = 0.018$) and 60 min post-EVH ($P = 0.003$; Fig. 2).

Peak of urinary 11 β -PGF_{2 α} post-EVH. Examination of the baseline and peak 11 β -PGF_{2 α} values revealed that EVH caused a significant increase in urinary 11 β -PGF_{2 α} concentration ($n = 24$; $P = 0.002$) that was inhibited with the inhalation of terbutaline ($P = 0.118$, peak vs. baseline; Fig. 3). The inhibition of 11 β -PGF_{2 α} excretion with terbutaline resulted in a significantly lower peak concentration of 11 β -PGF_{2 α} in the terbutaline condition compared with the placebo condition ($P = 0.001$; Fig. 3). Similarly, the magnitude of the change in urinary 11 β -PGF_{2 α} (prechallenge to peak postchallenge) was significantly reduced after pretreatment with terbutaline; from 12 (5–26) ng/mmol creatinine to 3 (–4 to 9) ng/mmol creatinine in the placebo and terbutaline condition, respectively ($P = 0.033$).

Spearman rank correlation coefficient revealed a weak, positive relationship ($r^2 = 0.251$; $P = 0.013$) between the fall in FEV₁ and the change in urinary 11 β -PGF_{2 α} (baseline to peak) post-EVH in the placebo condition. This relationship was absent with the administration of terbutaline ($P = 0.505$). Furthermore, multiple regression analysis revealed no significant relationship between the magnitude of change in urinary 11 β -PGF_{2 α} and the fall in FEV₁ post-EVH ($P = 0.117$).

DISCUSSION

The aim of this study was to determine whether a standard, single dose of inhaled short-acting β_2 -adrenoceptor agonist




Fig. 1. Maximum fall in forced expiratory volume in 1 s (FEV_1) following 8 min of hyperpnea of dry air in athletes with exercise-induced bronchoconstriction pretreated with 0.5 mg terbutaline (closed symbols) or placebo (open symbols). Individual data are shown; box plots represent group median and interquartile range, with whiskers representing the 5th and 95th percentiles. Values under the broken line (10% fall in FEV_1) represent complete broncho-protection ($n = 22$, 81%).

inhibits mast cell mediator release in a model of EIB. Our results demonstrate that 0.5 mg of terbutaline is able to inhibit the rise in urinary 11β -PGF $_{2\alpha}$ excretion following hyperpnea of dry air in athletes with EIB. These results provide the first in vivo evidence of an inhibition of mast cell activation by a low, clinically recommended dose of inhaled short-acting β_2 -adrenoceptor agonist in response to indirect bronchial provocation.

This study supports previous findings of mast cell activation following hyperpnea of dry air in athletes (34, 35). In vivo evidence using 11β -PGF $_{2\alpha}$ to assess the mast cell stabilizing effect of inhaled β_2 -adrenoceptor agonists in a model of EIB (i.e., bronchial provocation with dry powder mannitol) is limited to the use of a large dose of the long-acting β_2 -adrenoceptor agonist formoterol (15). Long-acting β_2 -adrenoceptor agonists are, however, not recommended as monotherapy for the treatment of asthma (27) or EIB (48). We therefore add novel, clinically relevant data, demonstrating that mast cell stabilization can occur following inhalation of a standard, single dose of the short-acting β_2 -adrenoceptor agonist terbutaline.

Mast cell activation in this study was assessed by a change in the urinary concentration of 11β -PGF $_{2\alpha}$. PGD $_2$, the parent molecule of 11β -PGF $_{2\alpha}$, is a major cyclooxygenase metabolite of arachidonic acid. In humans, PGD $_2$ is produced almost exclusively from mast cells (47), rendering the urinary metabolite 11β -PGF $_{2\alpha}$ a reliable and convenient marker of mast cell activation (47). A rise in urinary excretion of 11β -PGF $_{2\alpha}$ has

repeatedly been observed following induced bronchoconstriction in humans, irrespective of the trigger used: allergen (the gold-standard of mast cell dependent bronchoconstriction) (45), exercise (31, 39, 43, 46), EVH (34, 35), or mannitol (14, 15, 37). Further, in vitro work suggests that human mast cells are activated not only by antigen stimulation but also through osmotic stress (28). In the present study, the average ventilation of ~ 100 l/min would have caused water loss from the smaller airways and local hyperosmolarity of the airway surface liquid (22). This is relevant because mast cells are more densely located in the peripheral airways (6, 18). Furthermore, an increase in mast cell density has been noted in individuals with asthma and EIB (36), as well as in competitive swimmers (11). An increase in mast cell density in the peripheral airways, combined with the capability of hyperpnea of dry air to create a hyperosmotic environment, may explain the rise in 11β -PGF $_{2\alpha}$ in the placebo condition in the present study.

Importantly, the deposition of terbutaline from a dry powder inhaler also extends to the small airways (44). This may have facilitated the binding of terbutaline with the β_2 -adrenoceptors on the mast cells infiltrated in the peripheral airways of our study participants. In vitro, mast cells treated with the nonselective β -adrenergic agonist isoprenaline displayed increased levels of cAMP (49). Furthermore, this increase in cAMP correlated with a reduction in the release of the mast cell mediators PGD $_2$, histamine, and leukotriene C $_4$ (49). In vivo, inhaled terbutaline has also been shown to increase plasma cAMP (16). Because pharmacological agents that induce and sustain elevations of intracellular cAMP are well known to suppress mast cell secretions (56), it is likely that the inhibitory




Fig. 2. Median \pm interquartile range (Q1–Q3) urinary concentration of 11β -prostaglandin F $_{2\alpha}$ (11β -PGF $_{2\alpha}$) at baseline and 30 and 60 min post-eucapnic voluntary hyperpnea of dry air following pretreatment with 0.5 mg terbutaline (closed circles) or placebo (open circles) in 18 athletes with exercise-induced bronchoconstriction.




Fig. 3. Urinary concentrations of 11β -PGF $_{2\alpha}$ at baseline and after eucapnic voluntary hyperpnea (EVH) of dry air (peak value) following pretreatment with 0.5 mg terbutaline (closed circles) or placebo (open circles) in 24 athletes with exercise-induced bronchoconstriction. Individual data are shown; box plots represent group median and interquartile range, with whiskers representing the 5th and 95th percentiles.

effect of mast cell mediator release provided by terbutaline was partly mediated by β_2 -adrenoceptor-induced increase of intracellular cAMP.

Our data support the early proposal by Anderson and colleagues (4, 5) that inhaled β_2 -adrenoceptor agonist may attenuate EIB at least partly by inhibiting mast cell activation. We previously showed that inhaled sodium cromoglycate attenuates bronchoconstriction and urinary 11β -PGF $_{2\alpha}$ excretion following EVH (34). Because sodium cromoglycate has no known bronchorelaxant effect on airway smooth muscle (20, 55), we concluded that sodium cromoglycate attenuates hyperpnea-induced bronchoconstriction via mast cell stabilization (34). Similarly here, we propose that the prophylactic effect of terbutaline was partly mediated by inhibition of mast cell mediator release. This new observation in EVH is in keeping with previous data for allergen-induced bronchoconstriction, where albuterol offered significant bronchoprotection in association with the inhibition of mast cell mediator release (32). However, we cannot dismiss other possible modes of action of β_2 -adrenoceptor agonists within the airways. This would be in keeping with the absence of a relationship between bronchoconstriction (i.e., maximum fall in FEV $_1$ post-EVH) and urinary 11β -PGF $_{2\alpha}$ excretion (as observed in our multiple regression analysis), which may be explained by interindividual variation in the contribution of alternative modes of action of terbutaline.

Terbutaline has previously been shown to reduce airway plasma exudation (25). Since vascular engorgement and edema in response to hyperpnea are thought to exaggerate airway obstruction in EIB (3), the antiexudative action of β_2 -adrenoceptor agonists may have contributed to bronchoprotection in our study. Furthermore, there is a possibility that the β_2 -adrenoceptor agonist may have limited excessive and viscous mucosal accumulation by increasing cilia beat frequency (23) and improving clearance of alveolar (51) and/or mucociliary fluids (21, 41). Finally, the β_2 -adrenoceptor agonist may have preserved/improved airway caliber due to its well-known effect on airway smooth muscle tone. However, it has previously

been reported that large doses of orally administered β_2 -adrenoceptor agonists fail to prevent EIB while providing significant bronchodilatation (4, 5). Given these observations and our results showing a marked reduction in PGD $_2$ release, we consider that mast cell stabilization was an important contributor to the bronchoprotection in our experimental set-up.

Mast cell stabilizing treatments are deemed beneficial for the treatment of asthma/airway hyperresponsiveness (AHR) (13). Further, athletes have an increased prevalence of EIB (17), and mast cell activation has been observed in this population during dry air challenge independently to the occurrence of EIB (34, 35). Our findings are therefore of clinical relevance to individuals with asthma/AHR and to athletes. Given that PGD $_2$ may perpetuate inflammation via initiation of the migration of eosinophils to the airways (24), the prevention of mast cell mediator release with β_2 -adrenoceptor agonists may interrupt the chronic inflammatory cycle and improve the management of asthma and/or EIB. Importantly, however, a side effect of regular β_2 -adrenoceptor agonist usage may include the development of tolerance to the medication (30). Therefore, changes to the current guidelines for the management of EIB are not recommended at this time; the use of controller agents, such as inhaled corticosteroids, should be promoted when inhaled β_2 -adrenoceptor agonists have to be used frequently (48). It remains to be determined whether tolerance to β_2 -adrenoceptor agonists is associated in vivo with tolerance to their effects at the level of release of mast cell mediators (52).

To conclude, we have shown that the prophylactic administration of 0.5 mg of inhaled terbutaline not only offers a significant degree of bronchoprotection to athletes with EIB, but is also able to inhibit the release of the mast cell mediator PGD $_2$ following hyperpnea of dry air. These findings are in keeping with the proposal that the superior efficacy of aerosol formulation of β_2 -adrenoceptor agonist compared with tablets for preventing EIB is the inhibition of release of mast cell mediators (4, 5).

GRANTS

This work was performed with financial support from the Swedish Medical Research Council, the Heart Lung Foundation, the Vårdal Foundation, the Stockholm County Council Research Funds (ALF), the Swedish Strategic Research Foundation, the KI-SciLifeLab collaborations on Translational Medicine (ChAMP), the Karolinska Institutet, and the World Anti-Doping Agency.

DISCLOSURES

No conflicts of interest, financial or otherwise, are declared by the author(s).

AUTHOR CONTRIBUTIONS

Author contributions: A.J.S., J.R.B., and P.K. performed experiments; A.J.S. and J.R.B. analyzed data; A.J.S., J.R.B., S.D.A., S.-E.D., and P.K. interpreted results of experiments; A.J.S. prepared figures; A.J.S. drafted manuscript; A.J.S., S.D.A., L.M.R., B.D., S.-E.D., and P.K. edited and revised manuscript; A.J.S., J.R.B., S.D.A., L.M.R., B.D., S.-E.D., and P.K. approved final version of manuscript; S.D.A., L.M.R., B.D., S.-E.D., and P.K. conception and design of research.

REFERENCES

- Anderson SD, Argyros GJ, Magnussen H, Holzer K. Provocation by eucapnic voluntary hyperpnea to identify exercise induced bronchoconstriction. *Br J Sports Med* 35: 344–347, 2001.
- Anderson SD, Caillaud C, Brannan J. Beta2-agonists and exercise-induced asthma. *Clin Rev Allergy Immunol* 31: 163–180, 2006.
- Anderson SD, Daviskas E. The mechanism of exercise-induced asthma is . . . *J Allergy Clin Immunol* 106: 453–459, 2000.
- Anderson SD, Rozea PJ, Dolton R, Lindsay DA. Inhaled and oral bronchodilator therapy in exercise induced asthma. *Aust NZ J Med* 5: 544–550, 1975.
- Anderson SD, Seale JP, Rozea P, Bandler L, Theobald G, Lindsay DA. Inhaled and oral salbutamol in exercise-induced asthma. *Am Rev Respir Dis* 114: 493–500, 1976.
- Andersson CK, Mori M, Bjermer L, Löfdahl CG, Erjefält JS. Novel site-specific mast cell subpopulations in the human lung. *Thorax* 64: 297–305, 2009.
- Bierman EW, Kawabori I, Pierson WE. Incidence of exercise-induced asthma in children. *Pediatrics* 56: 847–850, 1975.
- Bland JM, Altman DG. Calculating correlation coefficients with repeated observations: Part 1—Correlation within subjects. *BMJ* 310: 446, 1995.
- Bonini M, Di Mambro C, Calderon MA, Compalati E, Schünemann H, Durham S, Canonica GW. Beta₂-agonists for exercise-induced asthma. *Cochrane Database Syst Rev* 10: CD003564, 2013.
- Bood JR, Sundblad BM, Delin I, Sjödin M, Larsson K, Anderson SD, Wheelock CE, Dahlén S, Dahlén B. Urinary excretion of lipid mediators in response to repeated eucapnic voluntary hyperpnea in asthmatic subjects. *J Appl Physiol* 119: 272–279, 2015.
- Bougault VV, Loubaki LL, Joubert PP, Turmel JJ, Couture CC, Lavolette MM, Chakir JJ, Boulet LPL. Airway remodeling and inflammation in competitive swimmers training in indoor chlorinated swimming pools. *J Allergy Clin Immunol* 129: 351–3e1, 2012.
- Bousquet J, Heinzlerling L, Bachert C, Papadopoulos NG, Bousquet PJ, Burney PG, Canonica GW, Carlsen KH, Cox L, Haahtela T, Carlsen KCL, Price D, Samolinski B, Simons FER, Wickman M, Annesi-Maesano I, Baena-Cagnani CE, Bergmann KC, Bindsløv-Jensen C, Casale TB, Chiriac A, Cruz AA, Dubakiene R, Durham SR, Fokkens WJ, Gerth-van-Wijk R, Kalayci O, Kowalski ML, Mari A, Mullol J, Nazamova-Baranova L, O'Hehir RE, Ohta K, Panzner P, Passalacqua G, Ring J, Rogala B, Romano A, Ryan D, Schmid-Grendelmeier P, Todo-Bom A, Valenta R, Woehrl S, Yusuf OM, Zuberbier T, Demoly P, Global Allergy and Asthma European Network, Allergic Rhinitis and its Impact on Asthma. Practical guide to skin prick tests in allergy to aeroallergens. *Allergy* 67: 18–24, 2012.
- Bradding P, Walls AF, Holgate ST. The role of the mast cell in the pathophysiology of asthma. *J Allergy Clin Immunol* 117: 1277–1284, 2006.
- Brannan J, Gulliksson M, Anderson SD, Chew N, Kumlin M. Evidence of mast cell activation and leukotriene release after mannitol inhalation. *Eur Respir J* 22: 491–496, 2003.
- Brannan J, Gulliksson M, Anderson SD, Chew N, Seale JP, Kumlin M. Inhibition of mast cell PGD₂ release protects against mannitol-induced airway narrowing. *Eur Respir J* 27: 944–950, 2006.
- Bye PT, Anderson SD, Daviskas E, Marty JJ, Sampson D. Plasma cyclic AMP levels in response to exercise and terbutaline sulphate aerosol in normal and asthmatic subjects. *Eur J Respir Dis* 61: 287–297, 1980.
- Carlsen KH, Anderson SD, Bjermer L, Bonini S, Brusasco V, Canonica W, Cummiskey J, Delgado L, Del Giacco SR, Drobic F, Haahtela T, Larsson K, Palange P, Popov T, van Cauwenberge P. Exercise-induced asthma, respiratory and allergic disorders in elite athletes: epidemiology, mechanisms and diagnosis: part I of the report from the Joint Task Force of the European Respiratory Society (ERS) and the European Academy of Allergy and Clinical Immunology (EAACI) in cooperation with GA 2LEN. *Allergy* 63: 387–403, 2008.
- Carroll NG, Mutavdzic S, James AL. Distribution and degranulation of airway mast cells in normal and asthmatic subjects. *Eur Respir J* 19: 879–885, 2002.
- Church MK, Hiroi J. Inhibition of IgE-dependent histamine release from human dispersed lung mast cells by anti-allergic drugs and salbutamol. *Br J Pharmacol* 90: 421–429, 1987.
- Cox JS. Disodium cromoglycate. Mode of action and its possible relevance to the clinical use of the drug. *Br J Dis Chest* 65: 189–204, 1971.
- Daviskas E, Anderson SD, Eberl S, Chan HK, Young IH, Seale JP. Effects of terbutaline in combination with mannitol on mucociliary clearance. *Eur Respir J* 20: 1423–1429, 2002.
- Daviskas E, Gonda I, Anderson SD. Mathematical modeling of heat and water transport in human respiratory tract. *J Appl Physiol* 69: 362–372, 1990.
- Devalia JL, Sapsford RJ, Rusznak C, Toumbis MJ, Davies RJ. The effects of salmeterol and salbutamol on ciliary beat frequency of cultured human bronchial epithelial cells, in vitro. *Pulm Pharmacol* 5: 257–263, 1992.
- Emery DL, Djokic TD, Graf PD, Nadel JA. Prostaglandin D₂ causes accumulation of eosinophils in the lumen of the dog trachea. *J Appl Physiol* 67: 959–962, 1989.
- Erjefält I, Persson CGA. Pharmacologic control of plasma exudation into tracheobronchial airways. *Am Rev Respir Dis* 143: 1008–1014, 1991.
- Faul F, Erdfelder E, Lang AG, Buchner A. G*Power 3: a flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behav Res Methods* 39: 175–191, 2007.
- Global Initiative for Asthma. *Global Strategy for Asthma Management and Prevention* (Online). Global Initiative for Asthma. http://www.ginasthma.org/local/uploads/files/GINA_Report_2014_Jun11.pdf [7 August, 2014].
- Gulliksson M, Palmberg L, Nilsson G, Ahlstedt S, Kumlin M. Release of prostaglandin D₂ and leukotriene C₄ in response to hyperosmolar stimulation of mast cells. *Allergy* 61: 1473–1479, 2006.
- Haby MM, Peat JK, Mellis CM, Anderson SD, Woolcock AJ. An exercise challenge for epidemiological studies of childhood asthma: validity and repeatability. *Eur Respir J* 8: 729–736, 1995.
- Hancox RJ, Subbarao P, Kamada D, Watson RM, Hargreave FE, Inman MD. Beta2-agonist tolerance and exercise-induced bronchospasm. *Am J Respir Crit Care Med* 165: 1068–1070, 2002.
- Haverkamp HC, Dempsey JA, Miller JD, Romer LM, Pegelow DF, Lovering AT, Eldridge MW. Repeat exercise normalizes the gas-exchange impairment induced by a previous exercise bout in asthmatic subjects. *J Appl Physiol* 99: 1843–1852, 2005.
- Howarth PH, Durham SR, Lee TH, Kay AB, Church MK, Holgate ST. Influence of albuterol, cromolyn sodium and ipratropium bromide on the airway and circulating mediator responses to allergen bronchial provocation in asthma. *Am Rev Respir Dis* 132: 986–992, 1985.
- Karjalainen J. Exercise response in 404 young men with asthma: no evidence for a late asthmatic reaction. *Thorax* 46: 100–104, 1991.
- Kippelen P, Larsson J, Anderson SD, Brannan J, Dahlén B, Dahlén S. Effect of sodium cromoglycate on mast cell mediators during hyperpnea in athletes. *Med Sci Sports Exerc* 42: 1853–1860, 2010.
- Kippelen P, Larsson J, Anderson SD, Brannan J, Delin I, Dahlén B, Dahlén S. Acute effects of beclomethasone on hyperpnea-induced bronchoconstriction. *Med Sci Sports Exerc* 42: 273–280, 2010.
- Lai Y, Altemeier WA, Vandree J, Piliponsky AM, Johnson B, Appel CL, Frevert CW, Hyde DM, Ziegler SF, Smith DE, Henderson WR, Gelb MH, Hallstrand T. Increased density of intraepithelial mast cells in patients with exercise-induced bronchoconstriction regulated through epithelially derived thymic stromal lymphopoietin and IL-33. *J Allergy Clin Immunol* 133: 1448–1455, 2014.
- Larsson J, Perry CP, Anderson SD, Brannan J, Dahlén S, Dahlén B. The occurrence of refractoriness and mast cell mediator release following mannitol-induced bronchoconstriction. *J Appl Physiol* 110: 1029–1035, 2011.
- Mickleborough TD, Murray RL, Ionescu AA, Lindley MR. Fish oil supplementation reduces severity of exercise-induced bronchoconstriction in elite athletes. *Am J Respir Crit Care Med* 168: 1181–1189, 2003.

39. Mickleborough TD, Rundell KW. Dietary polyunsaturated fatty acids in asthma- and exercise-induced bronchoconstriction. *Eur J Clin Nutr* 59: 1335–1346, 2005.
40. Miller MR, Hankinson JL, Brusasco V, Burgos F, Casaburi R, Coates A, Crapo RO, Enright P, van der Grinten CPM, Gustafsson P, Jensen R, Johnson D, MacIntyre N, McKay R, Navajas D, Pedersen OF, Pellegrino R, Viegi G, Wanger J. Standardisation of spirometry. *Eur Respir J* 26: 319–338, 2005.
41. Mortensen J, Groth S, Lange P, Hermansen F. Effect of terbutaline on mucociliary clearance in asthmatic and healthy subjects after inhalation from a pressurised inhaler and a dry powder inhaler. *Thorax* 46: 817–823, 1991.
42. Mortola JP. Breathing around the clock: an overview of the circadian pattern of respiration. *Eur J Appl Physiol* 91: 119–129, 2004.
43. Nagakura T, Obata T, Shichijo K, Matsuda S, Sigimoto H, Yamashita K, Masaki T, Maekawa K. GC/MS analysis of urinary excretion of 9alpha,11beta-PGF2 in acute and exercise-induced asthma in children. *Clin Exp Allergy* 28: 181–186, 1998.
44. Newman SP, Morén F, Trofast E, Talaee N, Clarke SW. Deposition and clinical efficacy of terbutaline sulphate from Turbuhaler, a new multi-dose powder inhaler. *Eur Respir J* 2: 247–252, 1989.
45. O’Sullivan S, Roquet A, Dahlén B, Dahlén S, Kumlin M. Urinary excretion of inflammatory mediators during allergen-induced early and late phase asthmatic reactions. *Clin Exp Allergy* 28: 1332–1339, 1998.
46. O’Sullivan S, Roquet A, Dahlén B, Larsen F, Eklund A, Kumlin M, O’Byrne PM, Dahlén S. Evidence for mast cell activation during exercise-induced bronchoconstriction. *Eur Respir J* 12: 345–350, 1998.
47. O’Sullivan S. On the role of PGD2 metabolites as markers of mast cell activation in asthma. *Acta Physiol Scand Suppl* 644: 1–74, 1999.
48. Parsons JP, Hallstrand T, Rundell KW, Anderson SD, Mastronarde JG, Kaminsky DA, Hull J, Storms WW, Weiler JM, Cheek FM, Wilson KC, American Thoracic Society Subcommittee on Exercise-induced Bronchoconstriction. An official American Thoracic Society clinical practice guideline: exercise-induced bronchoconstriction. *Am J Respir Crit Care Med* 187: 1016–1027, 2013.
49. Peachell PT, MacGlashan DW, Lichtenstein LM, Schleimer RP. Regulation of human basophil and lung mast cell function by cyclic adenosine monophosphate. *J Immunol* 140: 571–579, 1988.
50. Quanjer P, Tammeling G, Cotes J, Pedersen O, Peslin R, Yernault J. Lung volumes and forced ventilatory flows. Report working party standardization of lung function tests, European community for steel and coal Official statement of the European Respiratory Society. *Eur Respir J* 16: 5–35, 1993.
51. Sakuma T, Suzuki S, Usuda K, Handa M, Okaniwa G, Nakada T, Fujimura S, Matthey MA. Preservation of alveolar epithelial fluid transport mechanisms in rewarmed human lung after severe hypothermia. *J Appl Physiol* 80: 1681–1686, 1996.
52. Scola AM, Chong LK, Chess-Williams R, Peachell PT. Influence of agonist intrinsic activity on the desensitisation of beta2-adrenoceptor-mediated responses in mast cells. *Br J Pharmacol* 143: 71–80, 2004.
53. Simpson AJ, Tufvesson E, Anderson SD, Romer LM, Bjermer L, Kippelen P. Effect of terbutaline on hyperpnoea-induced bronchoconstriction and urinary club cell protein 16 in athletes. *J Appl Physiol* 115: 1450–1456, 2013.
54. Szczeklik A, Dworski R, Mastalerz L, Prokop A, Sheller JR, Nizankowska E, Cmiel A, Oates JA. Salmeterol prevents aspirin-induced attacks of asthma and interferes with eicosanoid metabolism. *Am J Respir Crit Care Med* 158: 1168–1172, 1998.
55. Thomson NC, Clark CJ, Boyd G, Moran F. Effect of sodium cromoglycate on bronchial smooth muscle. *Br J Clin Pharmacol* 12: 440–441, 1981.
56. Weston MC, Peachell PT. Regulation of human mast cell and basophil function by cAMP. *Gen Pharmacol* 31: 715–719, 1998.

