

The Arts in Psychotherapy

What makes us the same? What makes us different? Development of a shared model and manual of group therapy practice across art therapy, dance movement therapy and music therapy within community mental health care

--Manuscript Draft--

Manuscript Number:	AIP-D-20-00116R2
Article Type:	Full length article
Keywords:	Arts Therapies; Practice; mental health; manual; group; intervention development
Corresponding Author:	Catherine Carr, Ph.D. UNITED KINGDOM
First Author:	Catherine Elizabeth Carr, PhD, PGDip Music Therapy, BA(Hons)
Order of Authors:	Catherine Elizabeth Carr, PhD, PGDip Music Therapy, BA(Hons) Barbara Feldtkeller, PGCE, MSc, PGDip Dance Movement Psychotherapy Jennifer French, PGDip Music Therapy Dominik Havsteen-Franklin, PhD, MSc, PGDip Art Therapy Val Huet, PhD, MA, PGDip, BA(Hons) Vicky Karkou, MD, PhD, PGDip, MEd, BEd Stefan Priebe, FRCPsych, Dr med. habil, MD Stephen Sandford, MBA, PGDip Music Therapy, BMus (Hons)
Abstract:	Arts therapies draw upon many theories and models of practice, but lack consensus in shared factors of their provision. In preparation for a randomised controlled trial involving art therapy, dance movement therapy and music therapy, we used experiential and practice-based methods to develop a model and manual for mixed diagnosis arts therapies groups in community mental health care. Six arts therapists met over the course of a year to explore commonalities and differences in art, dance movement and music therapy groups. Arts-based and consensus methods were used to develop practice principles, which were expanded through workshops with arts therapists working in mental health services. A model, manual, training and adherence guide were produced, which incorporated role-play and video-based reflection. The model is underpinned by transdiagnostic and contextual psychotherapy theories. The manual outlines ten core principles, alongside 19 practice principles. Fourteen scenarios where therapists commonly intervene informed ongoing therapist training and development. This is the first model to describe practice across three types of arts therapies for mixed diagnosis groups in community mental health care. Whilst overall manual utility (including adherence) and effectiveness is yet to be assessed, the development methods may be informative for wider arts therapies groups and clientele.
Response to Reviewers:	We have unblinded the manuscript and edited the paragraphs so that there are fewer long stretches of text. We have also included a brief explanation in the introduction to make clear that the trial involves only three of the arts therapies and why we were unable to include dramatherapy (page 2).

Highlights

- We present a model of arts therapies practice for mixed diagnosis community mental health groups.
- This was developed for a trial involving art therapy, dance movement therapy and music therapy.
- The model considers the specific role of the arts and common group factors.
- Arts based workshops were used to develop a manual and practice principles for the trial.
- The training and development methods may be of interest to wider arts therapists and researchers.

Abstract

Arts therapies draw upon many theories and models of practice, but lack consensus in shared factors of their provision. In preparation for a randomised controlled trial involving art therapy, dance movement therapy and music therapy, we used experiential and practice-based methods to develop a model and manual for mixed diagnosis arts therapies groups in community mental health care. Six arts therapists met over the course of a year to explore commonalities and differences in art, dance movement and music therapy groups. Arts-based and consensus methods were used to develop practice principles, which were expanded through workshops with arts therapists working in mental health services. A model, manual, training and adherence guide were produced, which incorporated role-play and video-based reflection. The model is underpinned by transdiagnostic and contextual psychotherapy theories. The manual outlines ten core principles, alongside 19 practice principles. Fourteen scenarios where therapists commonly intervene informed ongoing therapist training and development. This is the first model to describe practice across three types of arts therapies for mixed diagnosis groups in community mental health care. Whilst overall manual utility (including adherence) and effectiveness is yet to be assessed, the development methods may be informative for wider arts therapies groups and clientele.

What makes us the same? What makes us different? Development of a shared model and manual of group therapy practice across art therapy, dance movement therapy and music therapy within community mental health care

Catherine Carr^{a,b*}, Barbara Feldtkeller^c, Jennifer French^b, Dominik Havsteen-Franklin^{d,e}, Val Huet^f, Vicky Karkou^g, Stefan Priebe^{a,b}, Stephen Sandford^b

^a Unit for Social and Community Psychiatry, Newham Centre for Mental Health, Glen Road, London E13 8SP United Kingdom. c.e.carr@gmul.ac.uk; s.priebe@gmul.ac.uk

^b East London Foundation NHS Trust, Robert Dolan House, 9 Alie Street, London E1 8DE United Kingdom. Jennifer.French1@nhs.net; stephensandford@nhs.net

^c Avon and Wiltshire Mental Health Partnership NHS Trust, Callington Road Hospital, Bristol BS4 5BJ United Kingdom. Barbara.Feldtkeller@nhs.net

^d Department of Arts and Humanities, Gaskell Building, Brunel University London, Uxbridge UB8 3PH United Kingdom. Dominik.Havsteen-Franklin@brunel.ac.uk

^e Central and North West London Foundation NHS Trust, 7a Woodfield Road, London W9 2BA United Kingdom.

^f British Association of Art Therapists, 24-27 White Lion Street, London N1 9PD United Kingdom. val@baat.org

^g Faculty of Arts and Sciences and Faculty of Health and Social Care, Edge Hill University, St Helens Road, Ormskirk L39 4QP United Kingdom. karkouv@edghill.ac.uk

* Corresponding author: c.e.carr@gmul.ac.uk Telephone: 07720685698


Declarations of interest: None

Acknowledgements:

We are grateful to all the arts therapists and service user who contributed their experience and expertise to the manual development workshops, Cornelia Bent and Emma Windle for their assistance in the set-up and running of these and Jessica Cardona and Lauren Hounsell for their careful proof-reading of citations and references.

Funding: This work was supported by East London NHS Foundation Trust and Avon and Wiltshire Partnership Mental Health Trust. The ERA study is funded by the National Institute for Health Research (NIHR) Health Technology Assessment programme [grant number 17/29/01]. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

Figure



The arts therapies are forms of psychotherapy that involve active participation in an art-form with an arts therapist to facilitate the therapeutic process through creative, non-verbal and verbal expression. Although in many countries arts therapies often have separate and varying training programmes, professional registration and recognition, they can also be regarded as sharing enough common characteristics to be part of joint clinical and research tasks (Karkou & Sanderson, 2006; Zubala, MacIntyre, Gleeson, & Karkou, 2014). Within the United Kingdom, art therapy, music therapy and dramatherapy are regulated by the Health and Care Professions Council (HCPC, 2013), while dance movement psychotherapists can achieve registration with the UK Council for Psychotherapy (UKCP, 2012). As with other forms of therapy, the arts therapies can be provided on an individual or group basis and there are many models of practice that have evolved from, and are tailored for different clinical groups and settings (Karkou & Sanderson, 2006). This diversity of practice enables the arts therapies to flexibly meet a range of health and care needs, but can make wider assessment and generalisation of their effectiveness challenging.

There is a rich body of arts therapies practice in mental health (Brooke, 2006; Hogan, 2001; Jones, 2007; McCaffrey, 2016; Nemetz, 2006). However literature tends to focus upon a single arts modality and in effectiveness studies, a specific diagnosis, which does not reflect actual provision (Fenner, Abdelazim, Bräuninger, Strehlow, & Seifert, 2017). Until relatively recently, there was also little consideration of the commonalities and differences of practice (Jones, 2020; Karkou & Sanderson, 2006) which limits our understanding of the factors shared across arts therapies and the practice principles that are important to apply in our work. This may be to some extent, historical. For example, within the United Kingdom, there were initial concerns of individual professions being grouped under one umbrella term of 'arts therapies' (Barrington, 2005) due to feared loss of identity and the

notion that approaches were too different or poorly understood to be considered together (Karkou & Sanderson, 2006; Waller, 1991). Conversely, there are many examples of arts therapists collaborating and building a shared understanding across modalities (Colbert & Bent, 2018; Havsteen-Franklin, Jovanovic, Reed, Charles, & Lucas, 2017; Huet, 1997; Knill, 1994; Landy, 1995; Purdie, 1993; Sajnani, Marxen & Zarate, 2017; Samaritter, 2018).

This paper presents developmental work to produce a model of group arts therapies practice for patients with different diagnoses (diagnostically heterogeneous) in community mental health care. The aim was to sufficiently identify, agree and describe core features of group arts therapies practice to produce a manual in preparation for a randomised controlled trial involving art therapy, dance movement therapy and music therapy (the ERA study, ISRCTN:88805048). Whilst the trial does not involve dramatherapy due to the logistics of the study design and local resources, we were keen to ensure the model could represent the full spectrum of arts therapies practice. Dramatherapists were therefore involved in the development process. In line with guidance for the development of complex interventions (Bleijenberg et al., 2018; Cotterill et al., 2018; Craig et al., 2019; Hoffmann et al., 2014; O’Cathain et al., 2019; Rousseau et al., 2019; Turner et al., 2019), our objectives were to (a) identify and agree upon theories that underpinned the intervention; (b) identify commonalities in the provision of art, dance movement and music therapy groups; and (c) develop a manual and set of principles to enable adherence rating in practice.

Considerations in Manualising Arts Therapies

Treatment manuals can be thought of as a means to describe the core “conceptual and structural boundaries” of how a treatment is provided and works (Addis, Cardemil, Duncan, & Miller 2006, p.132). Initially developed for research (Luborskey & DeRubeis, 1984),

manuals are also adopted or developed for clinical practice (Addis et al., 2006). Within randomised controlled trials, the purpose of a manual is to sufficiently describe the aims, methods and hypothesised mechanisms of the treatment(s) under investigation (Grant et al., 2018).

Manuals enable therapists to understand what is important to happen and how, and to monitor how far their practice aligns with this (Addis et al., 2006), whilst they enable researchers to check whether what is hypothesised to be of importance is actually provided across all participants (treatment fidelity) (Borrelli, 2011; Nelson, Cordray, Hulleman, Darrow, & Sommer, 2012). This enhances internal validity of the study, and enables future replication and comparison with different treatments (Nelson et al., 2012). The relative importance of practice elements to outcomes can also be better specified and understood (Kraemer, Wilson, Fairburn, & Agras, 2002; Nelson et al., 2012). As Rolvsjord, Gold and Stige (2005) note, use of a manual does not “guarantee valid results, nor will all results of studies without therapy manuals necessarily lack validity” (p.22). Manualisation does not aim for or guarantee superiority over non-manualised treatments (Truijens, Zühlke-van Hulzen, & Vanheule, 2019; Tshuschke et al., 2015); nor are novel manualised procedures easily adopted in routine practice (Bitsika & Sharpley, 2006; Carroll & Nuro, 2002; Chorpita, 2006).

Most arts therapies research manuals draw upon existing psychotherapy approaches, or wider theories to develop new interventions for specific populations, with a focus on one single arts modality (Azoulay & Orkibi, 2015; Behrends, Müller & Dziobek, 2012; Bryl & Goodill, 2019; Hackett, Taylor, et al., 2017; Parkinson & Whiter, 2016; Rolvsjord et al., 2005; Story & Beck, 2017; Zubala, 2013). Others have used intervention mapping (Bartholomew, Parcel, Kok, Gottlieb & Fernández, 2011) as a development method (Aalbers et al. 2019; Haeyen & Heijman, 2020; Haeyen, van Hooren, Dehue & Hutschemaekers, 2018). Those

manuals that have incorporated practices across the arts therapies focus on a multi-modal arts approach with a distinct client population such as depression (Parsons et al., 2020), or dementia (Lyons, 2019). One intervention has been developed across different arts therapies with children in mainstream schools (Moula, Karkou, & Powell, 2019). However, none so far have developed manuals across different arts therapies modalities for diagnostically heterogeneous groups in mental health care.

Within psychotherapy, there are a number of manuals for diagnostically heterogeneous groups. Notably, Barlow, Allen & Choate (2004) developed a transdiagnostic “uniform treatment protocol” for a range of emotional disorders, including adaptations for group treatment (Bullis et al., 2015; Laposka, Mancuso, Abraham, & Loli-Dano, 2017). Karterud (2016) similarly published a manual for mentalization-based group therapy. Both share a focus on common factors of group treatment, whilst also highlighting specific aspects of the approach.

Beyond research, a well-designed manual can also support therapists’ self-monitoring, professional development (Addis et al., 2006; Luborsky & DeRubeis, 1984) and act as a map or toolkit for therapists (Bryl & Goodill, 2019; Taylor-Buck & Dent-Brown, 2015). As such, some arts therapies professional bodies have developed clinical guidelines using consensus based methods, such as nominal group or Delphi technique (Eyre, 2013; Hackett, Ashby, Parker, Goody, & Power, 2017; Havsteen-Franklin, 2014; Springham, Dunne, Noyse & Swearingen, 2012; Taylor Buck, Dent-brown, Parry, & Boote, 2014; Taylor Buck & Hendry, 2016; Wright & Holttum, 2020).

Despite the many manualised forms of psychotherapy, studies of how such manualisation is delivered in practice highlight the challenges in doing so. Manuals are not always accepted or understood by therapists, who may perceive them as challenging their

professional identity, limiting therapeutic autonomy and creativity (Addis et al., 2006; Bergström, Ladd, Jones, Rosso, & Michael, 2019; Carroll & Nuro, 2002; Forbat, Black, & Dulgar, 2015; Truijens et al., 2019; Tschuschke et al., 2015). Implementation requires significant resource commitment (Addis et al., 2006; Chorpita, 2006) and if developed outside of the clinical context, may be limited in their external validity (Bitsika & Sharpley, 2006; Carroll & Nuro, 2002; Chorpita, 2006; Westen, 2002). Focus upon specific or novel factors of the manualised treatment can be to the detriment of known essential factors of therapy (e.g., therapeutic relationship) (Carroll & Nuro, 2002; Addis et al., 2006; Herman-Smith, Pearson, Cordiano, & Aguirre-McLaughlin, 2008); ignore the unique contribution of client and therapist (Addis et al., 2006; Rolvsjord et al., 2005) and focus upon technical execution to the detriment of training and supervision (Bein et al., 2000; Chorpita, 2006). This can have unintended and potentially negative consequences for both client outcomes (Henry et al., 1993; Vakoch & Strupp, 2000) and therapists' practice (Forbat et al., 2015).

Manuals can also fail to recognise the variation in therapist training, theoretical allegiance(s), experience, and competence (Cramer, Tschuschke, Koemeda, Schulthess, & von Wyl, 2020; Najavits, 1997; Tschuschke et al., 2015). For process-oriented therapies, the idea of following a predefined formula can be at odds with responding to the needs of the patient (Rolvsjord et al., 2005; Taylor Buck & Dent-brown, 2014; Vakoch & Strupp, 2000). Manuals may also challenge long-standing practice and require integration, building of confidence and skills (Najavits, 1997; Sholomskas et al., 2005). Research procedures and restrictions may also impact upon therapists' performance and confidence (Gold, Erkkilä & Crawford, 2012; Robb, Burns, Docherty, & Haas, 2011).

Manuals therefore require a fine balance of specification and flexibility. This includes careful consideration of common therapeutic factors (Beutler, 2006; Bradt, 2012; Rolvsjord

et al., 2005; Waltz, Addis, Koerner, & Jacobsen, 1993; Westen, 2002), practitioner involvement, clinical context (Bryl & Goodill, 2019; Carroll & Nuro, 2002; Chorpita, 2006; Forbat et al., 2015; Sidani & Braden, 2011; Westen, 2002) and therapist training and support (Butler, Henry, & Strupp, 1995; Carroll & Nuro, 2002; Gold et al., 2012; Robb et al., 2011). Studies of manual implementation have highlighted that therapists tend to work ‘tacitly’ or ‘implicitly’, responding to the person in the moment without thinking explicitly about the action being performed (Lee & Vakoch, 1996; Najavits, 1997; Vakoch & Strupp, 2000). Reber notes that within implicit learning “...not only is the learner unaware of the process of learning, the very knowledge itself is highly resistant to explication” (2001, p.15433). This may be even more important for arts therapists, given their frequent work within a non-verbal, implicit domain and the association of such processes with creativity (Gerge, Hawes, Eklöf, & Pedersen, 2019; McAleer, Bowler, Bowler, & Schoemann, 2020).

Studies have also demonstrated that overarching principles are more likely to be understood and implemented rather than listing predefined techniques (Lee & Vakoch, 1996; Vakoch & Strupp, 2000). Accordingly, some arts therapists have based manuals on Waltz, Addis and Koerner’s (1993) framework of principles for practice, rather than rules or procedures (Bryl & Goodill, 2019; Rolvsjord et al., 2005). In order to have an effective learning experience, Lee & Vakoch (1996) suggest that implicit knowledge first needs to be made explicit. Therefore, techniques to explore the tacit/implicit way in which therapists work with groups are important to draw out in practice. Given their non-verbal and creative possibilities, the arts are an effective medium to achieve this. We therefore used a range of arts and practice-based methods in the manual development of this study.

Method of Manual Development

Manual development began with monthly meetings between six arts therapists: two each of art therapy, dance movement therapy and music therapy. The arts therapists each had backgrounds covering arts therapies practice, academic research and strategic or managerial positions within a range of NHS mental health trusts. Our initial meetings highlighted challenges in learning one another's separate arts therapies languages and finding a shared understanding in the theories, tools and techniques we used. These meetings informed development of the underpinning theory and model which enabled the group to explore practice. Subsequent meetings focused upon defining principles of practice according to Waltz et al.'s framework (1993). Each arts therapist listed principles for practice according to their discipline and the techniques that would be used to achieve this. These were discussed as a group and formed the basis of 10 core principles of the manual, 18 principles for practice and 15 proscribed principles (Table 1).

We then used nominal group technique (Boddy, 2012; Harvey & Holmes, 2012; Havsteen-Franklin, 2014; Taylor Buck & Hendry, 2016), facilitated by DHF and CC to explore common clinical scenarios encountered within community mental health arts therapies groups. The focus of this technique was to describe common scenarios in group arts therapy where the therapist would make an intervention (defined as an observable action made by the therapist). It was highlighted that interventions could be both to address a clinical need or deficit, or to promote, encourage or highlight strengths and resources. Each group member listed as many scenarios as possible, over the course of 20 minutes. These were written up and displayed on flipchart paper around the room. From a list of 90 scenarios, these were then grouped thematically, resulting in a list of 14 scenarios (Table 2). In a final step, group members individually ranked the importance of each scenario based on their experience in practice. Individual ranking scores were combined to produce an overall group ranking.

Wider Arts Therapist Involvement and Feedback

Once the ten core principles, principles for practice and clinical scenarios were developed, we ran a 1-day workshop in two NHS Trusts, with 19 arts therapists in total. Attendees were arts therapists who had expressed an interest in running arts therapies groups in the forthcoming ERA study. One person with lived experience of mental health problems advising on the study and two dramatherapists also participated.

Arts-based methods were used to explore therapists' stances towards the level of directiveness used in sessions and group boundaries (Brabender & Fallon, 2018; Dies, 1985). For this, we used a movement-based exercise, where therapists explored the space between two polarities (e.g., active vs passive) in the room and then discussed the reasons for positioning themselves where they did. This led to reflections on instances where greater activity from the therapist, or less might be appropriate. Therapists explored how they established safety in early sessions by working with a range of arts objects and materials (e.g., scarves, wooden blocks, paper and pens), and core principles from the ensuing discussion were recorded on a flipchart. Group processes were explored through a music drawing narrative exercise (Booth, 2005) facilitated by CC, resulting in artistic representations of common group processes and a written reflection by each therapist. The experience of this exercise was discussed as a group and core themes from this discussion were written down. The group then discussed the aims of group arts therapy, contributing drawings and written words to large pieces of paper. The group reviewed 14 clinical scenarios developed to date and broke into small groups to role-play one of these scenarios as they had experienced in their own practice. This role play was specific to each arts modality with participants taking turns in the role of therapist. As a final step, the arts

therapists were asked to rank the importance of the 14 clinical scenarios, to provide additional feedback on the relevance of each scenario. Following these workshops, thematic analysis led to development of five core group aims and additional principles for practice (Table 1). The 10 core principles were unchanged, but were elaborated from workshop discussions to illustrate ‘What this is’ and ‘What this isn’t’ (following Rolvsjord et al., 2005).

Refinement and Selection of Principles for Practice

To refine the principles for practice, each principle was listed (20 from the main principles, 20 from promoted practices and 27 proscribed practices) and grouped thematically, forming 19 core areas. Core areas were then compared against existing group art, music and dance movement therapy literature in mental health and a group mentalization therapy adherence scale (Folmo et al., 2017; Karterud, 2016). In grouping the items, a number of principles shared amongst group therapies were identified. Where this was the case, the item naming, as entitled by Karterud (2016) was retained (items 1, 2, 6, 7, 8, 10). This formed the final structure for core principles and adherence rating.

Results

Underpinning Theory

Transdiagnostic Theory and Contextual Model of Psychotherapy

We took as our starting point, transdiagnostic theories of mental illness (Bakker, 2019; Casey, Olivery, & Insel, 2014; Sanislow, 2016). Whilst no single model exists, current theory suggests concepts of negative valence (e.g., acute threat/fear, loss), positive valence (e.g., motivation, reward), cognition, social processes, arousal and regulation are relevant to understanding shared features of mental illness (Cuthbert & Insel, 2013). There is particular interest in the transdiagnostic role of emotion regulation, with difficulties in using adaptive

regulation strategies, intolerance of uncertainty and experiential avoidance as three potential mechanisms (Aldao, 2012; Khakpoor, Bytamar, & Saed, 2019). Whilst the course and experience of mental illness is unique to each individual, within secondary mental health care, these transdiagnostic concepts are reflected in common challenges faced by patients in terms of managing and regulating mood and anxiety, maintaining motivation in daily life and becoming stuck in patterns of thinking, feeling or relating, with associated emotional and relational distress (Herrman, Saxena & Moodie, 2005; World Health Organisation, 2001). Relationships can be further affected through challenges in communicating experiences, thoughts and feelings making them difficult to develop or sustain (Brüne et al., 2009; France & Kramer, 2001; Kerns, 2007; Perlini et al., 2012; Sánchez et al., 2019). Many experience social isolation, stigmatisation and lack confidence to socialise and wider effects of disempowerment, loss of identity, role and self-esteem (Baker, Jodrey, & Intagliata, 1992; Goldberg, Rollins, & Lehman, 2003; Halford & Hayes, 1995; Thornicroft et al., 2016; Trompenaars, Masthoff, Van Heck, de Vries, & Hodiament, 2007).

We then drew upon the contextual model of psychotherapy (Wampold, 2015; Wampold & Imel, 2015), which suggests that features shared across different types of therapy are the most important factors that account for clinical change. Determinants include the client's preference for, understanding and expectations of therapy and the therapist's experience, commitment and belief to the model of therapy provided, along with a positive working relationship (Wampold, 2015; Wampold & Imel, 2015). In line with the contextual model, a meta-analysis showed that for mental health patients, receiving a preferred treatment is associated with lower dropout from treatment and higher therapeutic alliance (Windle et al., 2019). A further meta-analysis across group therapies has demonstrated that common shared factors account for the greatest amount of change in therapy, rather than unique

factors (Orfanos, Banks, & Priebe, 2015). These common group factors have been studied in various combinations, but include, at a meta-level; supportive factors (instillation of hope, acceptance, altruism, universality and cohesion); self-reflection (including self-disclosure, catharsis and secure emotional expression), psychological work (including interpersonal learning, self-understanding and awareness of relational impact) and social learning (including modelling, vicarious learning, guidance and education) (Mac Nair-Semands, Ogrodniczuk, & Joyce, 2010; MacKenzie, 1990; Yalom & Leszcz, 2005). On the basis of the above, we can assume that many of the important therapeutic factors in group arts therapies are those shared with wider forms of group therapy as a whole.

More recent research (Orfanos & Priebe, 2017) suggests that within closed community therapy groups, the first few sessions are most important in determining outcomes. Sessions where group members showed high cohesion in the first sessions predicted future engagement and greatest therapeutic gains. Reporting similar findings for different types of group psychotherapy, Tchuschke and Dies (1994) proposed that such early group integration promotes capacity for self-disclosure, which increases interpersonal feedback thus increasing opportunities for positive feedback from the group. Burlingame, Fuhriman and Johnson (2001) highlight the importance of certain practices in developing cohesion, namely: Thorough preparation of members for the group experience, identification of useful processes early on in therapy, consistency in temporal and spatial features of the group, fostering member to member interaction, feedback to the group, a warm and caring therapeutic stance and ensuring members relate to each other in positive ways. Within the arts therapies, a positive association has also been identified between therapeutic alliance and symptom reduction (Heynen, Roest, Willemars, & van Hooren 2017).

The above suggests that important factors within group therapy are patient

understanding and clear expectations of what will happen in therapy, patient and therapist shared commitment to working in a particular way and time to develop and establish a good working alliance. Within groups this highlights the importance of patient treatment preference, clear information on what to expect, fostering group cohesion in the initial phases and promoting active engagement quickly and early on in the therapeutic process.

The Additional Contribution of the Arts in Therapy

The defining factor of the arts therapies is emphasis upon the use of, and active participation in the arts during therapy. Koch (2017) suggests five factors specific to the arts therapies: aesthetics (beauty and authentic expression), hedonism (pleasure and play), meaning making (through non-verbal communication/symbolisation/metaphor), enactive transitional support in times of change, and generativity (creativity and creation). Inclusion of an active and creative arts-based process provides opportunities for embodied self-expression, creativity and a non-verbal means of relating with other people (Samaritter, 2018). The art form brings a third object- concrete (such as a recording or piece of art work), or experienced (such as movement or heard sound)- into the matrix of relations within the group (Karkou & Sanderson, 2006). Each person brings their own unique creative and cultural identity. This can be explored as part of the therapy with the potential to link and expand upon their personal relationship with, and use of the art-form as a 'helping resource' in their day-to-day lives (Ansdell & Meehan, 2010).

Springham and Huet (2018) suggest art-work in art therapy may be considered a second order representation within psychological processes (representing alternative 'as if' scenarios, rather than reality). This provides a means of trying out different ways of relating to oneself and others in an imaginative and safe way. Arts therapies are therefore well

placed to help patients identify difficulties and strengths through the use of the art-form in the varying group interactions. Use of the art-form can facilitate the experience and expression of emotions and discovery of personal meaning in creative activities, allowing experiences and learning through non-verbal and verbal communication (Gerge et al., 2019).

In other words, through creativity, imagination and play, patients are helped to explore new or different emotional and cognitive experiences with support of both the group and the therapist. The production of a piece of art is often cited as a means of strengthening self-esteem by clients (Buck & Havsteen-Franklin, 2013; Grocke et al., 2014) and can provide access to personal and interpersonal resources which may be continued into wider daily and creative life (Ansdell & Meehan, 2010; McCaffrey, Carr, Solli, & Hense, 2018; McCaffrey, Edwards, & Fannon, 2011).

Historically, arts therapists have integrated wider psychotherapy theories including expressive, psychodynamic, person-centred, attachment and group-interactive models (Karkou & Sanderson, 2006; Odell-Miller, 2013; Springham & Huet, 2018; Zubala, MacIntyre, Gleeson, & Karkou, 2013). Principles from each of these informed our understanding, along with principles of resource orientation (Priebe, Omer, Giacco, & Slade, 2014; Rolvsjord, 2010; Rolvsjord, 2015) and recovery (Leamy, Bird, Le Boutillier, Williams, & Slade, 2011; McCaffrey, 2016; McCaffrey et al., 2011, 2018; Solli, Rolvsjord, & Borg, 2013).

The principles of person-centred therapy (Rogers, 2011/1967) are especially relevant to how the therapist models and interacts with members of the group involving a non-directive, empathic stance. The psychodynamic model (Montgomery, 2002) provides a means of understanding patterns of relating and emotional responses across unconscious and conscious levels, which facilitates understanding of non-verbal and implicit levels of

communication. Resource and recovery-oriented approaches (Ansdell & Meehan, 2010; Buck & Havsteen-Franklin, 2013; McCaffrey et al., 2018; Rolvsjord, 2010) are informed by a move towards a collaborative, contextual and health-promotion focus rather than a deficit or treatment-based one. Principles include a shared, equal, collaborative process; focus on patients' strengths and potentials; acknowledging the patients' creative identity; being emotionally involved in the creative process and fostering positive emotions (McCaffrey et al., 2018).

Implications for the different arts modalities:

Whilst we hypothesise that it is the shared, or common factors of group therapy that are associated with eventual change of health outcomes, each arts modality has specific multi-sensory and aesthetic properties which can provide unique opportunities within a group context (Malchiodi, 2005). Whilst in simple terms a single sensory modality is implied (art, the eye; music, the ear; dance movement, the body), a range of senses are activated, offering many different sensory and embodied modes of expression and experience (Gerge et al., 2019; McNiff, 1981). Stern (2010) and Malloch and Trevarthen's (2009) studies of human communication show how timing, shape and intensity ("vitality affects") are used to non-verbally communicate intention and regulate interaction from birth. The multi-sensory and multi-modal features of arts-based work can facilitate attunement and regulation of basic physical and behavioural experiences, such as relaxation, self-soothing, and building of healthy attachments (Behrends et al., 2012; Metzner et al., 2018; Springham & Huet, 2018).

In our model (Figure 1), the importance of the art form is the appeal to the patient (their preferred type of arts therapy), the facilitation of active participation and emotional engagement in the use of the art form, the introduction of creativity, and the support of

exchange and interactions. The final therapeutically effective processes however are non-specific to the art form and will benefit from the diagnostic heterogeneity of the group. Figure 1 outlines how transdiagnostic theories of mental illness (orange), informed choice (yellow), common factors to group therapies (green), resource and recovery oriented theories (pink) and specific arts therapies processes (purple) relate within this model. We contextualise the arts-based processes within arts therapies groups as being an additional process to those that already occur within talking-based group therapy.

Manual

The manual encompasses five group aims, 10 overarching core principles, 14 clinical scenarios and 19 principles for therapist practice. The group aims relate directly to shared features of mental illness, whilst the overarching principles describe core features of how the groups should be managed over time. The clinical scenarios are used as a springboard for therapists to reflect on common features of running arts therapies groups. We purposefully did not define interventions for these scenarios as (a) many scenarios can unfold simultaneously and are not necessarily distinct; and (b) there are a multitude of ways that such scenarios might be addressed. Rather, we defined 19 principles for therapist practice which enable therapists to observe themselves and others when in action and form the basis of adherence rating for the ERA trial.

Group Aims

Our discussions identified five core group aims of arts therapies groups for community mental health settings. Importantly, these aims incorporate both implicit (embodied, felt or lived) and explicit (verbal and reflective) relational experiences and build from basic pre-requisites for therapy and the individual, to wider relational and community goals:

1. To provide a safe space to be with others

2. To promote engagement with the art form to

explore ways of using the art form and its relevance to one's own life;

explore inner creativity;

explore opportunities to extend and use the arts in daily life.

3. To provide regulation of arousal and affect to

explore ways of using the arts to aid self-regulation;

become aware of and find new ways of expressing and transforming emotions.

4. To offer opportunities to build upon existing relational capacity to

build awareness of self and others;

facilitate non-verbal engagement and expression with others;

explore new and different ways of relating to others;

break isolation and offer opportunities for support and socialisation.

5. Through the above, use the arts to build confidence, self-esteem, self-agency and autonomy in interactions with others and to further develop connectedness, hope, identity, meaning and empowerment (recovery) (Leamy et al., 2011).

Ten Core Principles

The ten core principles set the overall structure and format of groups over time. Items one to three highlight important actions to be taken by the therapist in preparation and support for the groups (such as non-attendance and team communication). Item four defines the general group structure for a single session, whilst items seven to nine describe the overall group task at beginning, middle and end stages of therapy. Items five, six and ten highlight the importance of active arts participation, the therapist's role in supporting the

group to become familiar with this and the importance of maintaining a collaborative and strengths-based approach, which is in keeping with a recovery focus.

1. The arts therapies groups are fully explained with information and video material so that patients can make an informed choice as to their preference

What this means: For the ERA study, potential members are shown video clips to aid decision making for the art-form that they most prefer. The videos explain what happens during an art therapy, dance movement therapy and music therapy session. Through the voices of service users, the videos cover what to expect, examples of activities during sessions and helpful and challenging features. Prior to the group start, an individual meeting is arranged with the therapists in the group therapy room, to provide a sense of what the space will look like. The meeting focuses upon introductions, the reasons for coming to therapy (including hopes and expectations) and explanation of the general aim, structure and process of group therapy. There is an opportunity to ask questions, agree ground rules and expectations, including the level of communication the person wishes to have.

What this is not: Insisting upon a particular modality choice that negates the patients' preference. Declining to discuss arts therapies options or concerns.

2. Attendance is encouraged and supported through maintaining active contact with patients throughout the treatment phase

What this is: Strategies to support attendance are discussed and agreed in the initial meeting and can include text reminders, and agreed modes of communication for non-attendance. Within the group, therapists acknowledge group absences and ensure time to welcome and re-orient returning members and late arrivals.

What this is not: Persistent calls to demand why the person is unable to attend. Ignoring members' preferences for attendance support. Ignoring group absences and their impact.

3. The therapists ensure communication is maintained and held with the multi-disciplinary care team throughout.

What this means: Therapists ensure ongoing communication with the wider care team to inform them of arising issues or concerns and to ensure high quality of support and care. The level of communication is outlined and agreed in the pre-group meeting, including the nature of confidentiality and boundaries of what will be shared with the wider care team.

What this is not: Ignoring group members' preferences for team communication. Not sharing important information regarding risk, safeguarding or wider care support.

4. Group structure always contains: (a) an opening warm-up; (b) use of the art-form, with space for verbal reflection; (c) a closing circle to reflect upon the group experience.

What this means: Sessions have an element of predictability, giving space for individuals to share at the beginning and end of the group. The middle section is flexible and the amount of alternation between using the art-form and talking may vary across modalities. Opening and closing sections should be led by the therapist to start, but may be developed or expanded upon by the group as sessions progress. The amount of time given to each section will vary depending on the stage of the group (early/late on in the process) and how group members are on any given day.

What this is not: Rigid reliance on exactly the same opening and closing activity each week. Not following the lead of the group in when to transition from and to activities.

5. Active participation, exploration and use of the art-form is encouraged.

What this means: Therapists aim to begin using the art-form with group members as quickly as possible in sessions, particularly where members are reluctant, reticent or avoidant. Use of the art-form should be introduced sensitively, with a level of activity and degree of structure that fits the presentation of the group. The therapist should ensure that the group does not go for long periods of talking or silence, without any use of the art-form.

What this is not: Forcing members to use an art-form when they are not ready to do so – a single group member might not directly use the art-form themselves, but may watch attentively to the rest of the group's creations. Ignoring the current situation of a group member which may make use of the art-form difficult on that day.

6. Therapists always offer structure within the art-form to promote group cohesion and only let this become free when there is space and safety to do so.

What this means: Structure is always offered to begin to establish an atmosphere of safety and cohesion between members within the group. Time should be allowed for group members to explore the art-form within the safety of such structures so that self-agency and competency in using the art-form are established.

What this is not: Limiting the level of freedom and activity in sessions to rigid or predefined structures. Not varying the level of structure according to presenting needs.

7. Within the early phase of therapy (weeks 1-5), the therapist focuses upon building trust and active engagement with the art-form, allowing plenty of time for this to happen.

8. Middle to late sessions (weeks 6-17): Developing content and arts-based activities in collaboration with service users to build variety (as opposed to doing the same arts-activity each session).

9. End phase (weeks 18-20): The group reflects on the process as a whole and review to prepare for ending. Therapists signpost patients to wider arts-based opportunities in the community.

What this means: Within the ERA study, sessions will run for up to 90 minutes, twice per week over 20 weeks. The above phases (in weeks) are a guide and it is acknowledged that each group will have its own unique early, middle and ending processes. What is important is for the therapist to be aware of the phase of therapy in relation to overall group process, so that time is allowed for each of these stages whilst balancing group cohesion and offering opportunities to challenge or extend group activities.

What this is not: Ignoring the stage of the group and propensity to use the art form. Persisting in activities without collaborating with group members. Not returning to an earlier stage if the group requires. Ignoring the interests and needs of the group.

10. Sessions are recovery-oriented: Therapists work collaboratively with patients so that their strengths and preferences in using the art-form are acknowledged to support their future recovery and signposting.

What this is: Therapists work collaboratively with patients with a here and now focus, so that their strengths and preferences in using the art-form are acknowledged. Group members are supported to explore issues raised, with challenges acknowledged. The therapist models curiosity in understanding members' experiences but does not immediately interpret or suggest meanings for the group. Throughout the sessions, the therapists acknowledge these strengths and preferences in terms of supporting future recovery. Therapists signpost group members where appropriate to wider community services which may meet the strengths and preferences expressed in the group.

What this is not: Ignoring risks, problems or clinical issues. Minimising or ignoring the impact of events from the past upon group members. Referring group members onto services without exploring preferences and wishes to do so together first.

Clinical Scenarios Where a Therapist Would Make an Intervention

Table 2 summarises 14 scenarios where arts therapists would make an intervention or observable action within the group. The items are listed in overall ranked order with 1, being the scenario ranked as most important and 14, the least important.

Principles for Practice and Adherence Rating

The 19 principles for practice (Table 3) encompass qualities the therapist displays in the group (items 1, 2, 10), and actions the therapist should take. Many are shared with other forms of group therapy but have the potential to also take an arts-based focus. For example, items 3 and 4 are especially important within arts therapies, where the art-form can be used jointly by the patient and therapist to aid connection to others (attunement) and change arousal state. Here, the role of the therapist is to support self-regulation through containment and transformation of expression. Items 11, 12, 13 and 18 are solely arts-based, whilst items 15 and 17 combine both common and arts group processes. Items 1, 2, 6, 7, 8 and 10 relate directly to common factors outlined by Karterud, and thus retained the original names (Folmo et al., 2017; Karterud, 2016). Items 4 and 15, whilst similar to Karterud (Regulation of tension level; Focus on emotions), were renamed to better align with the transdiagnostic focus, and acknowledge the wider range of possible implicit arts therapies experiences (bodily sensation, emotion, or as represented in the art form).

Discussion

Strengths and Limitations

The manual addresses many of the challenges faced in defining a single model of group arts therapy for diagnostically heterogeneous patients. The model has a clear theoretical basis and considers the pragmatic challenges of providing a standardised framework for three different arts modalities. Most importantly, we were able to reach agreement upon core aims and practice principles between therapists from different arts modalities and training backgrounds. Our development process followed current guidance for the development of complex interventions (Bleijenberg et al., 2018; Cotterill et al., 2018; Craig et al., 2019; Hoffmann et al., 2014; O’Cathain et al., 2019; Rousseau et al., 2019; Turner et al., 2019) and incorporated feedback from arts therapists. Experiential and practice-based methods enabled direct exploration and representation of diverse but context-specific practice. Core principles are clear and simple, whilst the practice principles accommodate a range of training backgrounds. The clinical scenarios provide the first comprehensive synthesis of situations considered important for arts therapists to act within a group.

Whilst the model is pluralistic in its wider approaches, this may pose challenges for arts therapists who have trained and practise in a defined therapy paradigm. For example, arts therapists working from a formal psychoanalytic model may be challenged by the more active and collaborative stance. Therapists with a strong person-centred leaning may be challenged by greater therapist directiveness early on in sessions. Whilst we believe that the manual encompasses current group arts therapy practice in community health care, it remains to be seen whether it has utility beyond the life of the study and within regular community-based practice. Feedback from arts therapists in the trial and further review by external bodies (for example, training providers) will assist further validation.

As the forthcoming ERA study encompasses only art, dance movement and music therapy, dramatherapists had limited involvement and thus, this model may not fully reflect aspects of dramatherapy practice. However, we hope that this may be examined separately from the ERA trial in collaboration with dramatherapists in the future. There was also limited service-user involvement, with only one service user attending the workshops. Further collaboration with and feedback from service users (both study participants and from a lived experience advisory panel) during the ERA trial will be important to understand the relevance of this way of working, including helpful and unhelpful factors.

The manual is necessarily transdiagnostic (Aldao, 2012; Cuthbert, 2014; Cuthbert & Insel, 2013; Khakpoor et al., 2019), and does not define diagnosis-specific actions or principles. However, it also does not contradict existing guidelines (Havsteen-Franklin, 2014; NICE, 2014; Springham et al., 2012; Wright & Holttum, 2020). Further work will be necessary to align where diagnosis-specific actions are indicated and required (Beutler, 2006).

Comparison to Existing Arts Therapies Practice

This manual differs to others developed so far, in that it seeks to identify shared aims and elements of practice across different arts therapy modalities and diagnoses. However, there is strong congruence between our manual and those practices outlined by specific arts therapists and professional bodies (Eyre, 2013; Hackett, Ashby, et al., 2017; Havsteen-Franklin, 2014; Parsons et al., 2020; Springham et al., 2012; Taylor Buck et al., 2014; Taylor Buck & Hendry, 2016; Wright & Holttum, 2020). In their study of therapist descriptions of in-session therapeutic constructs, Havsteen-Franklin et al. (2017), identified ten symptomatic events and 14 bipolar constructs. These events, whilst focusing upon only symptoms, have some similarities to the 14 scenarios identified by this group including: high and low affect,

acute psychosis, ambivalent responses, concrete solutions to interpersonal problems and insight. Similarly, the bipolar constructs cover similar areas of attunement, boundaries, affect regulation, directiveness, curiosity and use of the arts. Samaritter's (2018) Delphi study of music and dance movement therapists suggests five core themes of embodied presence, somato-sensory engagement, emotional engagement, non-verbal communication and intercultural involvement. The 12 steps identified as arts related activities and procedures, follow a similar pattern to those described in our 19 principles for practice; from invitation and engagement to exploration, play and performance (Samaritter, 2018). Parsons et al. (2020) and Parkinson and Whiter (2016) also describe similar aims, group structure and principles for depression and early intervention for psychosis.

The role of Experiential and Practice-Based Arts Methods in Training

Arts-based methods are gaining interest as a tool for creative exploration in clinical practice and research and were central to the manual development (Beer, 2016; Borgdorff, 2010; Geiger, Shpigelman, & Feniger-Schaal, 2020; Gerber et al., 2020; Ledger & McCaffrey, 2015; McCaffrey et al., 2020; McCaffrey & Edwards, 2015; Meltzer, 2020; Priebe et al., 2019; Sajnani, Mayor, & Tillberg-Webb, 2020; Springborg, 2020). As arts therapists, we recognised the value in working creatively with our colleagues to recognise the tensions and congruence across training traditions and art-forms. The workshops were lively and inclusive, enabling us to capture the essence of practice across arts modalities through our group reflections on the creative arts process.

The workshops informed how we provide training of therapists for the trial itself. Initial feedback suggested that the arts therapists valued opportunities to reflect upon and explore their practice. We therefore built in a strong ethos for the training to promote therapist self-

exploration of how their practice aligns with the model and manual principles, alongside regular opportunities for therapists to provide their ideas and feedback. Training is provided in three whole day workshops spread over a few months.

Day one provides an overview of the ERA trial and the theory underpinning the intervention. Three core principles (7-9) are explored by therapist pairs leading an experiential session to demonstrate how these principles would be met in their arts modality. Therapists are asked at the end of this day to begin peer observation to become familiar in being observed for research and discussing practice with another.

Day two involves feedback from therapists' experiences of peer observation and role plays based on one of the 14 clinical scenarios. Role plays are video recorded to give therapists a sense of how recording will feel when conducting sessions and for review in training day 3. Therapists are introduced to the adherence rating scale and encouraged to begin using this to assess their current practice. Introductory interviews and study procedures are also discussed.

The final training day covers role play of introductory meetings, how to report adverse events and adherence rating. Therapists watch excerpts of videos from training day 2. Therapists from the modality in the video rate this excerpt, with the remaining group observing and later commenting upon the overall process. Excerpts are selected to ensure coverage of all 19 principles for practice and discussions focus upon the ways in which the principles can be observed in the videos and put into practice.

Conclusion

This is the first model of arts therapies group practice developed across three arts modalities for mixed diagnosis groups in community mental health care. Our emphasis on

direct arts-based experience led to development of a rich corpus of practice. Most importantly, we were able to reach agreement of core principles between three different arts modalities. Such methods may be of wider value to arts therapists in clinical supervision and reflective practice, as well as wider clinicians and researchers. Whilst manual utility (including adherence) and model effectiveness is yet to be assessed, the methods may prove useful in developing models of practice for wider arts therapies groups and clientele.

Key words: arts therapies; practice; mental health; manual; group; intervention development

Table legend:

Table 1: Development of Practice Principles Through Initial Meetings and Therapist Workshops

Table 2: Clinical Scenarios Where Therapists Make an Observable Therapeutic Action

Table 3: Therapist Principles for Practice

Figure legend: *Figure to be printed in colour*

Figure 1: Logic model summarising underpinning theory, group aims, core principles, therapist practice principles and related scenarios.

References

- Aalbers, S., Vink, A., Freeman, R. E., Pattiselanno, K., Spreen, M., & van Hooren, S. (2019). Development of an improvisational music therapy intervention for young adults with depressive symptoms: An intervention mapping study. *Arts in Psychotherapy, 65*, 101584. <https://doi.org/10.1016/j.aip.2019.101584>
- Addis, M. E., Cardemil, E. V., Duncan, B. L., & Miller, S. D. (2006). Does manualization improve therapy outcomes? In J. C. Norcross, L. E. Beutler, & R. F. Levant (Eds.), *Evidence-based practices in mental health: Debate and dialogue on the fundamental questions* (pp. 131–160). Washington, DC: American Psychological Association. <https://doi.org/10.1037/11265-003>
- Aldao, A. (2012). Emotion regulation strategies as transdiagnostic processes: A closer look at the invariance of their form and function. *Revista de Psicopatología y Psicología Clínica, 17*(3), 261–277. <https://doi.org/10.5944/rppc.vol.17.num.3.2012.11843>
- Ansdell, G., & Meehan, J. (2010). “Some light at the end of the tunnel”: Exploring users’ evidence for the effectiveness of music therapy in adult mental health settings. *Music and Medicine, 2*, 29–40. <https://doi.org/10.1177/1943862109352482>
- Azoulay, B., & Orkibi, H. (2015). The four-phase CBN psychodrama model: A manualized approach for practice and research. *Arts in Psychotherapy, 42*, 10–18. <https://doi.org/10.1016/j.aip.2014.12.012>
- Baker, F., Jodrey, D., & Intagliata, J. (1992). Social support and quality of life of community support clients. *Community Mental Health Journal, 28*(5), 397–411.

<https://doi.org/10.1007/BF00761058>

- Bakker, G. M. (2019). A new conception and subsequent taxonomy of clinical psychological problems. *BMC Psychology*, 7(1), 46. <https://doi.org/10.1186/s40359-019-0318-8>
- Barlow, D. H., Allen, L. B., & Choate, M. L. (2004). Toward a unified treatment for emotional disorders. *Behavior Therapy*, 35(2), 205–230. [https://doi.org/10.1016/S0005-7894\(04\)80036-4](https://doi.org/10.1016/S0005-7894(04)80036-4)
- Barrington, K. A. (2005). *Music Therapy: A study in professionalisation* (Doctoral dissertation). Retrieved from etheses.dur.ac.uk/2791/
- Bartholomew, L. K., Parcel, G. S., Kok, G., Gottlieb, N. H., & Fernández, M. E. (2011). *Planning health promotion programs: An intervention mapping approach* (3rd ed.). San Francisco, CA: Jossey-Bass.
- Beer, L. E. (2016). From embedded to embodied: including music in arts-based music therapy research. *Music Therapy Perspectives*, 34(1), 33–40. <https://doi.org/10.1093/mtp/miv006>
- Behrends, A., Müller, S., & Dziobek, I. (2012). Moving in and out of synchrony: A concept for a new intervention fostering empathy through interactional movement and dance. *Arts in Psychotherapy*, 39(2), 107–116. <https://doi.org/10.1016/j.aip.2012.02.003>
- Bein, E., Anderson, T., Strupp, H. H., Henry, W. P., Schacht, T. E., Binder, J. L., & Butler, S. F. (2000). The effects of training in time-limited dynamic psychotherapy: Changes in therapeutic outcome. *Psychotherapy Research*, 10(2), 119–132. <https://doi.org/10.1080/713663669>
- Bergström, B., Ladd, A., Jones, A., Rosso, L., & Michael, P. (2019). A national survey of clinical

- psychology graduate student attitudes toward psychotherapy treatment manuals: A replication study. *International Journal of Business, Human and Social Sciences*, 12(6), 923–928. <http://doi.org/10.5281/zenodo.3299863>
- Beutler, L. E. (2006). It isn't the size, but the fit. *Clinical Psychology: Science and Practice*, 9(4), 434–438. <https://doi.org/10.1093/clipsy.9.4.434>
- Bitsika, V., & Sharpley, C. F. (2006). Treating the client rather than the symptoms: Moving beyond manualised treatments in psychotherapy. *Australian Journal of Guidance and Counselling*, 16(2), 159–175. <https://doi.org/10.1375/ajgc.16.2.159>
- Bleijenberg, N., de Man-van Ginkel, J. M., Trappenburg, J. C. A., Ettema, R. G. A., Sino, C. G., Heim, N., ... Schuurmans, M. J. (2018). Increasing value and reducing waste by optimizing the development of complex interventions: Enriching the development phase of the Medical Research Council (MRC) Framework. *International Journal of Nursing Studies*, 79, 86–93. <https://doi.org/10.1016/j.ijnurstu.2017.12.001>
- Boddy, C. (2012). The nominal group technique: An aid to brainstorming ideas in research. *Qualitative Market Research: An International Journal*, 15(1), 6–18. <https://doi.org/10.1108/13522751211191964>
- Booth, J. M. (2005). Music, drawing and narrative: An adaptation of the Bonny method of guided imagery and music. *Journal of the Association for Music and Imagery*, 10, 55–74. Retrieved from <https://ami-bonnymethod.org/resources/journals>
- Borgdorff, H. (2010). The production of knowledge in artistic research. In M. Biggs & H. Karlsson (Eds.), *The Routledge Companion to Research in the Arts* (pp. 44–63). New York, NY: Routledge. Retrieved from <https://doi.org/10.4324/9780203841327.ch3>

- Borrelli, B. (2011). The assessment, monitoring, and enhancement of treatment fidelity in public health clinical trials. *Journal of Public Health Dentistry*, 71(s1), S52–S63.
<https://doi.org/10.1111/j.1752-7325.2011.00233.x>
- Brabender, V., & Fallon, A. (2018). *Group psychotherapy in inpatient, partial hospital, and residential care settings*. Washington, DC: American Psychological Association.
Retrieved from <https://doi.org/10.1037/0000113-000>
- Bradt, J. (2012). Randomized controlled trials in music therapy: Guidelines for design. *Journal of Music Therapy*, 49(2), 120–149. <https://doi.org/10.1093/jmt/49.2.120>
- Brooke, S. L. (Ed.) (2006). *Creative arts therapies manual: A guide to the history, theoretical approaches, assessment, and work with special populations of art, play, dance, music, drama and poetry therapies*. Springfield, IL: Charles C Thomas.
- Brüne, M., Abdel-Hamid, M., Sonntag, C., Lehmkämpfer, C., & Langdon, R. (2009). Linking social cognition with social interaction: Non-verbal expressivity, social competence and “mentalising” in patients with schizophrenia spectrum disorders. *Behavioral and Brain Functions*, 5(6). <https://doi.org/10.1186/1744-9081-5-6>
- Bryl, K., & Goodill, S. (2019). Development, execution and acceptance of a manualized dance/movement therapy treatment protocol for the clinical trial in the treatment of negative symptoms and psychosocial functioning in schizophrenia. *American Journal of Dance Therapy*, 1–26. <https://doi.org/10.1007/s10465-019-09312-8>
- Buck, E. T., & Havsteen-Franklin, D. (2013). Connecting with the image: How art psychotherapy can help to re-establish a sense of epistemic trust. *ATOL: Art Therapy Online*, 4(1). <https://doi.org/10.25602/GOLD.atol.v4i1.310>

- Bullis, J. R., Sauer-Zavala, S., Bentley, K. H., Thompson-Hollands, J., Carl, J. R., & Barlow, D. H. (2015). The unified protocol for transdiagnostic treatment of emotional disorders: preliminary exploration of effectiveness for group delivery. *Behavior Modification, 39*(2), 295–321. <https://doi.org/10.1177/0145445514553094>
- Burlingame, G. M., Fuhriman, A., & Johnson, J. E. (2001). Cohesion in group psychotherapy. *Psychotherapy, 38*(4), 373–379. <https://doi.org/10.1037/0033-3204.38.4.373>
- Butler, S. F., Henry, W. P., & Strupp, H. H. (1995). Measuring adherence in time-limited dynamic psychotherapy. *Psychotherapy: Theory, Research, Practice, Training, 32*(4), 629–638. <https://doi.org/10.1037/0033-3204.32.4.629>
- Carroll, K. M., & Nuro, K. F. (2002). One size cannot fit all: A stage model for psychotherapy manual development. *Clinical Psychology: Science and Practice, 9*(4), 396–406. <https://doi.org/10.1093/clipsy/9.4.396>
- Casey, B. J., Oliveri, M. E., & Insel, T. (2014). A neurodevelopmental perspective on the research domain criteria (RDoC) framework. *Biological Psychiatry, 76*(5), 350–353. <https://doi.org/10.1016/j.biopsych.2014.01.006>
- Chorpita, B. F. (2006). Treatment manuals for the real world: Where do we build them? *Clinical Psychology: Science and Practice, 9*(4), 431–433. <https://doi.org/10.1093/clipsy.9.4.431>
- Colbert, T. & Bent, C. (Eds.) (2018). *Working across modalities in the arts therapies: Creative collaborations*. Abingdon: Routledge.
- Cotterill, S., Knowles, S., Martindale, A. M., Elvey, R., Howard, S., Coupe, N., ... Spence, M. (2018). Getting messier with TIDieR: Embracing context and complexity in intervention

reporting. *BMC Medical Research Methodology*, 18(1).

<https://doi.org/10.1186/s12874-017-0461-y>

Craig, P., Dieppe, P., Macintyre, S., Michie, S., Nazareth, I., & Petticrew, M. (2006).

Developing and evaluating complex interventions. Swindon: Medical Research Council.

Retrieved from <https://mrc.ukri.org/documents/pdf/complex-interventions-guidance/>

Cramer, A., Tschuschke, V., Koemeda, M., Schulthess, P., & von Wyl, A. (2020). The

therapists' training and their attitudes towards therapy as predictors of therapeutic interventions. *Journal of Contemporary Psychotherapy*, 50(1), 67–76.

<https://doi.org/10.1007/s10879-019-09421-y>

Cuthbert, B. N. (2014). The RDoC framework: Facilitating transition from ICD/DSM to

dimensional approaches that integrate neuroscience and psychopathology. *World*

Psychiatry, 13(1), 28–35. <https://doi.org/10.1002/wps.20087>

Cuthbert, B. N., & Insel, T. R. (2013). Toward the future of psychiatric diagnosis: The seven

pillars of RDoC. *BMC Medicine*, 11(126), 1–8.

<https://doi.org/10.1186/1741-7015-11-126>

Dies, R. R. (1985). Leadership in short-term group therapy: Manipulation or facilitation?

International Journal of Group Psychotherapy, 35(3), 435–455.

<https://doi.org/10.1080/00207284.1985.11491426>

Eyre, L. (2013). Adults in a recovery model setting. In L. Eyre (Ed.), *Guidelines for music*

therapy practice in mental health (pp. 115–167). Gilsum, NH: Barcelona.

Feder, E., & Feder, B. (1981). *The expressive arts therapies*. Upper Saddle River, NJ: Prentice

Hall.

- Fenner, P., Abdelazim, R. S., Bräuninger, I., Strehlow, G., & Seifert, K. (2017). Provision of arts therapies for people with severe mental illness. *Current Opinion in Psychiatry*, *30*(4), 306–311. <https://doi.org/10.1016/j.biopsych.2014.01.006>
- Folmo, E. J., Karterud, S. W., Bremer, K., Walther, K. L., Kvarstein, E. H., & Pedersen, G. A. F. (2017). The design of the MBT-G adherence and quality scale. *Scandinavian Journal of Psychology*, *58*(4), 341–349. <https://doi.org/10.1111/sjop.12375>
- Forbat, L., Black, L., & Dulgar, K. (2015). What clinicians think of manualized psychotherapy interventions: Findings from a systematic review. *Journal of Family Therapy*, *37*(4), 409–428. <https://doi.org/10.1111/1467-6427.12036>
- France, J. & Kramer, S. (Eds.) (2001). *Communication and mental illness. Theoretical and practical approaches*. London, UK: Jessica Kingsley
- Geiger, A., Shpigelman, C. N., & Feniger-Schaal, R. (2020). The socio-emotional world of adolescents with intellectual disability: A drama therapy-based participatory action research. *Arts in Psychotherapy*, *70*, 101679. <https://doi.org/10.1016/j.aip.2020.101679>
- Gerber, N., Biffi, E., Biondo, J., Gemignani, M., Hannes, K., & Siegesmund, R. (2020). Arts-based research in the social and health sciences: Pushing for change with an interdisciplinary global arts-based research initiative. *Forum Qualitative Sozialforschung*, *21*(2). <https://doi.org/10.17169/fqs-21.2.3496>
- Gerge, A., Hawes, J., Eklöf, L., & Pedersen, I. (2019). Proposed mechanisms of change in arts-based psychotherapies. *Voices: A World Forum for Music Therapy*, *19*(2), 31. <https://doi.org/10.15845/voices.v19i2.2564>

- Gold, C., Erkkilä, J., & Crawford, M. J. (2012). Shifting effects in randomised controlled trials of complex interventions: A new kind of performance bias? *Acta Psychiatrica Scandinavica*, *125*(5), 307–314. <https://doi.org/10.1111/j.1600-0447.2012.01922.x>
- Goldberg, R. W., Rollins, A. L., & Lehman, A. F. (2003). Social network correlates among people with psychiatric disabilities. *Psychiatric Rehabilitation Journal*, *26*(4), 393–403. <https://doi.apa.org/doi/10.2975/26.2003.393.402>
- Grant, S., Mayo-Wilson, E., Montgomery, P., Macdonald, G., Michie, S., Hopewell, S., & Moher, D. (2018). CONSORT-SPI 2018. Explanation and elaboration: Guidance for reporting social and psychological intervention trials. *Trials*, *19*, 406. <https://doi.org/10.1186/s13063-018-2735-z>
- Grocke, D., Bloch, S., Castle, D., Thompson, G., Newton, R., Stewart, S., ... Gold, C. (2014). Group music therapy for severe mental illness: A randomized embedded- experimental mixed methods study. *Acta Psychiatrica Scandinavica*, *130*, 144–153. <https://doi.org/10.1111/acps.12224>
- Hackett, S. S., Ashby, L., Parker, K., Goody, S., & Power, N. (2017). UK art therapy practice-based guidelines for children and adults with learning disabilities. *International Journal of Art Therapy*, *22*(2), 84–94. <https://doi.org/10.1080/17454832.2017.1319870>
- Hackett, S. S., Taylor, J. L., Freeston, M., Jahoda, A., McColl, E., Pennington, L., & Kaner, E. (2017). Interpersonal art psychotherapy for the treatment of aggression in people with learning disabilities in secure care: A protocol for a randomised controlled feasibility study. *Pilot and Feasibility Studies*, *3*, 42. <https://doi.org/10.1186/s40814-017-0186-z>
- Haeyen, S., & Heijman, J. (2020). Compassion focused art therapy for people diagnosed with

a cluster b/c personality disorder: An intervention mapping study. *Arts in Psychotherapy*, 69. <https://doi.org/10.1016/j.aip.2020.101663>

Haeyen, S., van Hooren, S., Dehue, F., & Hutschemaekers, G. (2018). Development of an art-therapy intervention for patients with personality disorders: an intervention mapping study. *International Journal of Art Therapy: Inscape*, 23(3), 125–135. <https://doi.org/10.1080/17454832.2017.1403458>

Halford, W., & Hayes, R. (1995). Social skills in schizophrenia: Assessing the relationship between social skills, psychopathology and community functioning. *Social Psychiatry and Psychiatric Epidemiology*, 30(1), 14–19. <https://doi-org.ezproxy.library.qmul.ac.uk/10.1007/BF00784429>

Harvey, N., & Holmes, C. A. (2012). Nominal group technique: An effective method for obtaining group consensus. *International Journal of Nursing Practice*, 18(2), 188–194. <https://doi.org/10.1111/j.1440-172X.2012.02017.x>

Havsteen-Franklin, D. (2014). Consensus for using an arts-based response in art therapy. *International Journal of Art Therapy: Inscape*, 19(3), 107–113. <https://doi.org/10.1080/17454832.2014.968598>

Havsteen-Franklin, D., Jovanovic, N., Reed, N., Charles, M., & Lucas, C. (2017). Developing a shared language within arts psychotherapies: A personal construct psychology approach to understanding clinical change. *Arts in Psychotherapy*, 55, 103–110. <https://doi.org/10.1016/j.aip.2017.05.002>

Health and Care Professions Council. (2013). *Standards of proficiency- Arts Therapists*. Retrieved from <https://www.hcpc-uk.org/resources/standards/standards-of->

proficiency-arts-therapists/

Henry, W. P., Strupp, H. H., Butler, S. F., Schacht, T. E., & Binder, J. L. (1993). Effects of training in time-limited dynamic psychotherapy: Changes in therapist behavior. *Journal of Consulting and Clinical Psychology, 61*(3), 434–440. <https://doi.org/10.1037/0022-006X.61.3.434>

Herman-Smith, R., Pearson, B., Cordiano, T. S., & Aguirre-McLaughlin, A. M. (2008). Addressing individual client needs in manualized treatment: Case comparisons. *Clinical Case Studies, 7*(5), 377–396. <https://doi.org/10.1177/1534650108315867>

Herrman, H., Saxena, S., & Moodie, R. (2005). *Promoting mental health: Concepts, emerging evidence, practice*. Geneva: World Health Organisation. Retrieved from http://whqlibdoc.who.int/publications/2005/9241562943_eng.pdf?ua=1

Heynen, E., Roest, J., Willemars, G., & van Hooren, S. (2017). Therapeutic alliance is a factor of change in arts therapies and psychomotor therapy with adults who have mental health problems. *Arts in Psychotherapy, 55*, 111–115. <https://doi.org/10.1016/j.aip.2017.05.006>

Hoffmann, T. C., Glasziou, P. P., Boutron, I., Milne, R., Perera, R., Moher, D., ... Michie, S. (2014). Better reporting of interventions: Template for intervention description and replication (TIDieR) checklist and guide. *BMJ, 348*. <https://doi.org/10.1136/bmj.g1687>

Hogan, S. (2001). *Healing arts: The history of art therapy*. London: Jessica Kingsley.

Huet, V. (1997). Challenging professional confidence in arts therapies and psychiatric rehabilitation. *Inscape, 2*(1), 14–19. <https://doi.org/10.1080/17454839708413040>

Jones, D. P. (2020). *The arts therapies: A revolution in healthcare* (2nd Ed.). London, UK:

Routledge. <https://doi.org/10.4324/9780203308066>

Jones, P. (2007). *Drama as therapy: Theory, practice and research* (2nd ed.). London: Routledge.

Karkou, V., & Sanderson, P. (2006). *The arts therapies: A research based map of the field*. Edinburgh: Elsevier. <https://doi.org/10.1016/B978-0-443-07256-7.X5001-3>

Karterud, S. (2016). *Mentalization-based group therapy (MBT-G): A theoretical, clinical, and research manual*. Oxford, UK: Oxford University Press.
<https://doi.org/10.1093/med:psych/9780198753742.001.0001>

Kerns, J. (2007). Verbal communication impairments and cognitive control components in people with schizophrenia. *Journal of Abnormal Psychology, 116*(2), 279–289.
<https://doi.org/10.1037/0021-843X.116.2.279>

Khakpoor, S., Bytamar, J. M., & Saed, O. (2019). Reductions in transdiagnostic factors as the potential mechanisms of change in treatment outcomes in the Unified Protocol: A randomized clinical trial. *Research in Psychotherapy: Psychopathology, Process and Outcome, 22*(3), 402–412. <https://doi.org/10.4081/ripppo.2019.379>

Koch, S. C. (2017). Arts and health: Active factors and a theory framework of embodied aesthetics. *Arts in Psychotherapy, 54*, 85–91. <https://doi.org/10.1016/j.aip.2017.02.002>

Knill, P. J. (1994). Multiplicity as a tradition: Theories for interdisciplinary arts therapies- an overview. *Arts in Psychotherapy, 21*(5), 319–328.
[https://doi.org/10.1016/0197-4556\(94\)90059-0](https://doi.org/10.1016/0197-4556(94)90059-0)

Kraemer, H. C., Wilson, T., Fairburn, C. G., & Agras, S. (2002). Mediators and moderators of treatment effects in randomized clinical trials. *Archives of General Psychiatry, 59*(10),

877–883. <https://doi.org/10.1001/archpsyc.59.10.877>

Landy, R. J. (1995). Isolation and collaboration in the creative arts therapies- the implications of crossing borders. *Arts in Psychotherapy*, *22*(2), 83–86.

[https://doi.org/10.1016/0197-4556\(95\)00010-3](https://doi.org/10.1016/0197-4556(95)00010-3)

Laposa, J. M., Mancuso, E., Abraham, G., & Loli-Dano, L. (2017). Unified protocol transdiagnostic treatment in group format: A preliminary investigation with anxious individuals. *Behavior Modification*, *41*(2), 253–268.

<https://doi.org/10.1177/0145445516667664>

Leamy, M., Bird, V., Le Boutillier, C., Williams, J., & Slade, M. (2011). Conceptual framework for personal recovery in mental health: Systematic review and narrative synthesis. In *British Journal of Psychiatry*, *199*(6), 445–452.

<https://doi.org/10.1192/bjp.bp.110.083733>

Ledger, A., & McCaffrey, T. (2015). Performative, arts-based, or arts-informed? Reflections on the development of arts-based research in music therapy. *Journal of Music Therapy*, *52*(4), 441–456. <https://doi.org/10.1093/JMT/THV013>

Lee, Y., & Vakoch, D. A. (1996). Transfer and retention of implicit and explicit learning.

British Journal of Psychology, *87*(4), 637–651.

<https://doi.org/10.1111/j.2044-8295.1996.tb02613.x>

Lyons, S. (2019). *Arts therapies for dementia: A systematic review and community-based case study on the value of music therapy and dance movement therapy* (Unpublished doctoral dissertation). Liverpool, UK: Edge Hill University.

Luborksy, L., & DeRubeis, R. (1984). The use of psychotherapy treatment manuals: A small

revolution in psychotherapy research style. *Clinical Psychology Review*, 4(1), 5–14.

[https://doi.org/10.1016/0272-7358\(84\)90034-5](https://doi.org/10.1016/0272-7358(84)90034-5)

Mac Nair-Semands, R. R., Ogradniczuk, J. S., & Joyce, A. S. (2010). Structure and initial validation of a short form of the therapeutic factors inventory. *International Journal of Group Psychotherapy*, 60(2), 245–281. <https://doi.org/10.1521/ijgp.2010.60.2.245>

MacKenzie, K. R. (1990). *Introduction to time-limited group psychotherapy*. Washington, DC: American Psychiatric Press.

Malchiodi, C. A. (Ed.) (2005). *Expressive therapies*. New York, NY: Guilford Press.

Malloch, S., & Trevarthen, C. (2009). Musicality: Communicating the vitality and interests of life. In S. Malloch & C. Trevarthen (Eds.), *Communicative musicality: Exploring the basis of human companionship* (pp. 1–11). Oxford, UK: Oxford University Press.

McAleer, J. T., Bowler, J. L., Bowler, M. C., & Schoemann, A. M. (2020). Implicit and explicit creativity: Further evidence of the integrative model. *Personality and Individual Differences*, 154, 109643. <https://doi.org/10.1016/j.paid.2019.109643>

McCaffrey, T. (2016). Music therapy in mental health care for adults. In J. Edwards (Ed.) *The Oxford handbook of music therapy* (pp.249–267). Oxford, UK: Oxford University Press. <https://doi.org/10.1093/oxfordhb/9780199639755.013.29>

McCaffrey, T., Carr, C., Solli, H. P., & Hense, C. (2018). Music therapy and recovery in mental health: seeking a way forward. *Voices: A World Forum for Music Therapy*, 18(1). <https://doi.org/10.15845/voices.v18i1.918>

McCaffrey, T., & Edwards, J. (2015). Meeting art with art: Arts-based methods enhance researcher reflexivity in research with mental health service users. *Journal of Music*

Therapy, 52(4), 515–532. <https://doi.org/10.1093/JMT/THV016>

McCaffrey, T., Edwards, J., & Fannon, D. (2011). Is there a role for music therapy in the recovery approach in mental health? *Arts in Psychotherapy*, 38(3), 185–189.

<https://doi.org/10.1016/j.aip.2011.04.006>

McCaffrey, T., Higgins, P., Monahan, C., Moloney, S., Nelligan, S., Clancy, A., & Cheung, P. S. (2020). Exploring the role and impact of group songwriting with multiple stakeholders in recovery-oriented mental health services. *Nordic Journal of Music Therapy*, 1–20.

<https://doi.org/10.1080/08098131.2020.1771755>

McNiff, S. (1981). *The Arts and Psychotherapy*. Springfield, IL: Charles C Thomas.

Meltzer, C. (2020). The reflexive practitioner. Using arts-based methods and research for professional development. In L. Mackay, G. Barton, S. Garvis, & V. Sappa (Eds.), *Arts-based research: Resilience and well-being across the lifespan* (pp. 209–231). Cham:

Palgrave Macmillan. https://doi.org/10.1007/978-3-030-26053-8_12

Metzner, S., Jaeger, U., Masuhr, O., Olschewski, U., Gräfe, E., Böske, A. C., & Dümpelmann, M. (2018). Forms of attunement during the initial stages of music therapy for patients with acute psychosis - a multicentre clinical study. *Nordic Journal of Music Therapy*,

27(5), 360–380. <https://doi.org/10.1080/08098131.2018.1478879>

Montgomery, C. (2002). Role of dynamic group therapy in psychiatry. *Advances in Psychiatric Treatment*, 8(1), 34–41. [https://doi.org/DOI: 10.1192/apt.8.1.34](https://doi.org/DOI:10.1192/apt.8.1.34)

Moula, Z., Karkou, V., & Powell, J. (2019). A pilot cross-over randomised controlled trial of child-focused process and outcome evaluation of arts therapies at primary mainstream schools: study protocol. *Arts and Health*.

<https://doi.org/10.1080/17533015.2019.1703198>

Najavits, L. M. (1997). Psychotherapists' implicit theories of therapy. *Journal of Psychotherapy Integration*, 7(1), 1–16. <https://doi.org/10.1037/h0101137>

National Institute of Clinical Excellence. (2014). *Psychosis and schizophrenia in adults: treatment and management. Guidance and guidelines*. National Institute of Clinical Excellence. Retrieved from <https://www.nice.org.uk/guidance/cg178>

Nelson, M. C., Cordray, D. S., Hulleman, C. S., Darrow, C. L., & Sommer, E. C. (2012). A procedure for assessing intervention fidelity in experiments testing educational and behavioral interventions. *Journal of Behavioral Health Services and Research*, 39(4), 374–396. <https://doi.org/10.1007/s11414-012-9295-x>

Nemetz, L. D. (2006). Moving with meaning: The historical progression of dance/movement therapy. In S. L. Brooke (Ed.), *Creative arts therapies manual: A guide to the history, theoretical approaches, assessment, and work with special populations of art, play, dance, music, drama and poetry therapies* (pp. 95–108). Springfield, IL: Charles C Thomas.

O’Cathain, A., Croot, L., Duncan, E., Rousseau, N., Sworn, K., Turner, K. M., ... Hoddinott, P. (2019). Guidance on how to develop complex interventions to improve health and healthcare. *BMJ Open*, 9(8). <https://doi.org/10.1136/bmjopen-2019-029954>

Odell-Miller, H. (2013). The development of clinical music therapy in adult mental health practice: Music, health and therapy. Case study. In V. Bates, A. Bleakley, & S. Goodman (Eds.), *Medicine health and the arts: Approaches to the medical humanities* (pp. 264–280). London, UK: Routledge. <https://doi.org/10.4324/9780203079614>

Orfanos, S., Banks, C., & Priebe, S. (2015). Are group psychotherapeutic treatments effective for patients with schizophrenia? A systematic review and meta-analysis. *Psychotherapy and Psychosomatics*, *84*(4), 241–249.

<https://doi.org/10.1159/000377705>

Orfanos, S., & Priebe, S. (2017). Group therapies for schizophrenia: initial group climate predicts changes in negative symptoms. *Psychosis*, *9*(3), 225–234.

<https://doi.org/10.1080/17522439.2017.1311360>

Parkinson, S., & Whiter, C. (2016). Exploring art therapy group practice in early intervention psychosis. *International Journal of Art Therapy: Inscape*, *21*(3), 116–127.

<https://doi.org/10.1080/17454832.2016.1175492>

Parsons, A., Omylinska-Thurston, J., Karkou, V., Harlow, J., Haslam, S., Hobson, J., ... Griffin, J. (2020). Arts for the blues – a new creative psychological therapy for depression. *British Journal of Guidance & Counselling*, *48*(1), 5–20.

<https://doi.org/10.1080/03069885.2019.1633459>

Perlini, C., Marini, A., Garzitto, M., Isola, M., Cerruti, S., Marinelli, V., ... Brambilla, P. (2012). Linguistic production and syntactic comprehension in schizophrenia and bipolar disorder. *Acta Psychiatrica Scandinavica*, *126*(5), 363–376.

<https://doi.org/10.1111/j.1600-0447.2012.01864.x>

Priebe, S., Arenas Borrero, Á., Bird, V., Džubur Kulenović, A., Giacco, D., Gómez Restrepo, C., ... Uribe Restrepo, J. M. (2019). Possibilities for the future of global mental health: A scenario planning approach. *BMC Psychiatry*, *19*(1). <https://doi.org/10.1186/s12888-019-2381-3>

Priebe, S., Omer, S., Giacco, D., & Slade, M. (2014). Resource-oriented therapeutic models in psychiatry: Conceptual review. *British Journal of Psychiatry*, *204*(4), 256–261.

<https://doi.org/10.1192/bjp.bp.113.135038>

Purdie, H. (1993). Music therapy and its relationship with creative arts therapies. *British Journal of Music Therapy*, *7*(2), 23–25.

<https://doi.org/10.1177%2F135945759300700204>

Reber, A. S. (2001). Tacit knowledge, psychology of. In N. J. Smelser, & P. B. Baltes (Eds.), *International Encyclopedia of the Social & Behavioral Sciences* (pp.15431–15435).

Oxford: Elsevier.

<https://doi.org/10.1016/B0-08-043076-7/01492-3>

Robb, S. L., Burns, D. S., Docherty, S. L., & Haase, J. E. (2011). Ensuring treatment fidelity in a multi-site behavioral intervention study: Implementing NIH behavior change consortium recommendations in the SMART trial. *Psycho-Oncology*, *20*(11), 1193–1201. <https://doi.org/10.1002/pon.1845>

Rogers, C. R. (2011). *On Becoming a Person. A therapist's view of psychotherapy*. London: Constable & Robinson. (Original work published in 1967)

Rolvsjord, R. (2010). *Resource-oriented music therapy in mental health care*. Gilsum, NH: Barcelona. Retrieved from <https://www.barcelonapublishers.com/resource-oriented-music-therapy-mental-health-care>

Rolvsjord, R. (2015). What clients do to make music therapy work: A qualitative multiple case study in adult mental health care. *Nordic Journal of Music Therapy*, *24*(4), 296–321. <https://doi.org/10.1080/08098131.2014.964753>

- Rolvjord, R., Gold, C., & Stige, B. (2005). Research rigour and therapeutic flexibility: Rationale for a therapy manual developed for a randomised controlled trial. *Nordic Journal of Music Therapy*, *14*(1), 15–32. <https://doi.org/10.1080/08098130509478122>
- Rousseau, N., Turner, K. M., Duncan, E., O’Cathain, A., Croot, L., Yardley, L., & Hoddinott, P. (2019). Attending to design when developing complex health interventions: A qualitative interview study with intervention developers and associated stakeholders. *PLoS ONE*, *14*(10). <https://doi.org/10.1371/journal.pone.0223615>
- Sajnani, N., Marxen, E. & Zarate, R. (2017). Critical perspectives in the arts therapies: Response/ability across a continuum of practice. *Arts in Psychotherapy*, *54*, 28–37. <https://doi.org/10.1016/j.aip.2017.01.007>
- Sajnani, N., Mayor, C., & Tillberg-Webb, H. (2020). Aesthetic presence: The role of the arts in the education of creative arts therapists in the classroom and online. *Arts in Psychotherapy*, *69*, 101668. <https://doi.org/10.1016/j.aip.2020.101668>
- Samaritter, R. (2018). The aesthetic turn in mental health: reflections on an explorative study into practices in the arts therapies. *Behavioral Sciences*, *8*(4), 41. <https://doi.org/10.3390/bs8040041>
- Sánchez, J., Sung, C., Phillips, B. N., Tschopp, M. K., Muller, V., Lee, H. –L., & Chan, F. (2019). Predictors of perceived social effectiveness of individuals with serious mental illness. *Psychiatric Rehabilitation Journal*, *42*(1), 88–99. <https://doi.apa.org/doi/10.1037/prj0000321>
- Sanislow, C. (2016). Connecting psychopathology meta-structure and mechanisms. *Journal of Abnormal Psychology*, *125*(8), 1158–1165.

<https://content.apa.org/doi/10.1037/abn0000207>

Sholomskas, D. E., Syracuse-Siewert, G., Rounsaville, B. J., Ball, S. A., Nuro, K. F., & Carroll, K. M. (2005). We don't train in vain: A dissemination trial of three strategies of training clinicians in cognitive-behavioral therapy. *Journal of Consulting and Clinical Psychology, 73*(1), 106–115. <https://doi.org/10.1037/0022-006X.73.1.106>

Sidani, S. & Braden, C. J. (2011). *Design, evaluation and translation of nursing interventions*. Chichester, UK: John Wiley & Sons.

Solli, H. P., Rolvsjord, R., & Borg, M. (2013). Toward understanding music therapy as a recovery-oriented practice within mental health care: A meta-synthesis of service users' experiences. *Journal of Music Therapy, 50*(4), 244–273. <https://doi.org/10.1093/jmt/50.4.244>

Springborg, C. (2020). Designing dance into qualitative research. In J. Ward, & H. Shortt (Eds.), *Using arts-based research methods* (pp. 41–74). Cham: Palgrave Macmillan. https://doi.org/10.1007/978-3-030-33069-9_3

Springham, N., Dunne, K., Noyse, S., & Swearingen, K. (2012). Art therapy for personality disorder: 2012 UK professional consensus guidelines, development process and outcome. *International Journal of Art Therapy, 17*(3), 130–134. <https://doi.org/10.1080/17454832.2012.734834>

Springham, N., & Huet, V. (2018). Art as relational encounter: An ostensive communication theory of art therapy. *Art Therapy, 35*(1), 4–10. <https://doi.org/10.1080/07421656.2018.1460103>

Stern, D. N. (2010). *Forms of vitality: Exploring dynamic experience in psychology, the arts,*

psychotherapy and development. Oxford, UK: Oxford University Press.

Story, K. M., & Beck, B. D. (2017). Guided imagery and music with female military veterans:

An intervention development study. *Arts in Psychotherapy, 55*, 93–102.

<https://doi.org/10.1016/j.aip.2017.05.003>

Taylor Buck, E., & Dent-brown, K. (2014). Creative manuals. *International Journal of Art*

Therapy, 19(2), 82–87. <https://doi.org/10.1080/17454832.2014.906475>

Taylor Buck, E., Dent-Brown, K., Parry, G., & Boote, J. (2014). Dyadic art psychotherapy: Key principles, practices and competences. *Arts in Psychotherapy, 41*(2), 163–173.

<https://doi.org/10.1016/j.aip.2014.01.004>

Taylor Buck, E., & Hendry, A. (2016). Developing principles of best practice for art therapists working with children and families. *International Journal of Art Therapy, 21*(2), 56–65.

<https://doi.org/10.1080/17454832.2016.1170056>

Thornicroft, G., Mehta, N., Clement, S., Evans-Lacko, S., Doherty, M., Rose, D., ... Henderson, C. (2016). Evidence for effective interventions to reduce mental-health-related stigma and discrimination. *Lancet, 387*(10023), 12–18.

[https://doi.org/10.1016/S0140-6736\(15\)00298-6](https://doi.org/10.1016/S0140-6736(15)00298-6)

Trompenaars, F. J., Masthoff, E. D., Van Heck, G. L., de Vries, J., & Hodiament, P. P. (2007).

Relationships between social functioning and quality of life in a population of Dutch adult psychiatric outpatients. *International Journal of Social Psychiatry, 53*(1), 36–47.

<https://doi.org/10.1177/0020764006074281>

Truijens, F., Zühlke-van Hulzen, L., & Vanheule, S. (2019). To manualize, or not to manualize:

Is that still the question? A systematic review of empirical evidence for manual

superiority in psychological treatment. *Journal of Clinical Psychology*, 75(3), 329–343.

<https://doi.org/10.1002/jclp.22712>

Tschuschke, V., & Dies, R. (1994). Intensive analysis of therapeutic factors and outcome in long-term inpatient groups. *International Journal of Group Psychotherapy*, 44(2), 185–208. <https://doi.org/10.1080/00207284.1994.11490742>

Tschuschke, V., Cramer, A., Koehler, M., Berglar, J., Muth, K., Staczan, P., ... Koemeda-Lutz, M. (2015). The role of therapists' treatment adherence, professional experience, therapeutic alliance, and clients' severity of psychological problems: Prediction of treatment outcome in eight different psychotherapy approaches. Preliminary results of a naturalistic study. *Psychotherapy Research*, 25(4), 420–434.

<https://doi.org/10.1080/10503307.2014.896055>

Turner, K. M., Rousseau, N., Croot, L., Duncan, E., Yardley, L., O' Cathain, A., & Hoddinott, P. (2019). Understanding successful development of complex health and healthcare interventions and its drivers from the perspective of developers and wider stakeholders: An international qualitative interview study. *BMJ Open*, 9(5).

<https://doi.org/10.1136/bmjopen-2018-028756>

UKCP. (2012). *UKCP standards of education and training (2012) - the minimum core criteria psychotherapy with adults*.

Retrieved from <http://www.psychotherapy.org.uk/index.php?id=428>

Vakoch, D. A., & Strupp, H. H. (2000). The evolution of psychotherapy training: Reflections on manual-based learning and future alternatives. *Journal of Clinical Psychology*, 56(3), 309–318.

[https://doi.org/10.1002/\(SICI\)1097-4679\(200003\)56:3<309::AID-JCLP7>3.0.CO;2-7](https://doi.org/10.1002/(SICI)1097-4679(200003)56:3<309::AID-JCLP7>3.0.CO;2-7)

Waller, D. (1991). *Becoming a profession: The history of art therapy in Britain 1940-1982*

(*Psychology revivals*). London: Routledge. <https://doi.org/10.4324/9780203750209>

Waltz, J., Addis, M. E., Koerner, K., & Jacobson, N. S. (1993). Testing the integrity of a

psychotherapy protocol: Assessment of adherence and competence. *Journal of*

Consulting and Clinical Psychology, 61(4), 620–630. [https://doi.org/10.1037/0022-](https://doi.org/10.1037/0022-006X.61.4.620)

006X.61.4.620

Wampold, B. E. (2015). How important are the common factors in psychotherapy? An

update. *World Psychiatry*, 14(3), 270–277. <https://doi.org/10.1002/wps.20238>

Wampold, B. E., & Imel, Z. E. (2015). *The great psychotherapy debate: The evidence for what*

makes psychotherapy work (2nd ed.). New York, NY: Routledge.

<https://doi.org/10.4324/9780203582015>

Westen, D. (2002). Manualizing manual development. *Clinical Psychology: Science and*

Practice, 9(4), 416–418. <https://doi.org/10.1093/clipsy.9.4.416>

Windle, E., Tee, H., Sabitova, A., Jovanovic, N., Priebe, S., & Carr, C. (2020). Association of

patient treatment preference with dropout and clinical outcomes in adult psychosocial

mental health interventions: a systematic review and meta-analysis. *JAMA Psychiatry*,

77(3), 294–302. <https://doi.org/10.1001/jamapsychiatry.2019.3750>

World Health Organization (2001). *Mental Health: New understanding, new hope*. Geneva:

World Health Organization.

Retrieved from https://www.who.int/whr/2001/en/whr01_en.pdf?ua=1

Wright, T., & Holttum, S. (2020). *BAAT Guidelines on Art Therapy for People with a*

Psychosis-Related Diagnosis. Retrieved from

[https://www.baat.org/Assets/Docs/General/BAAT Guidelines AT Psychosis PART 1-2-3.pdf](https://www.baat.org/Assets/Docs/General/BAAT%20Guidelines%20AT%20Psychosis%20PART%201-2-3.pdf)

Yalom, I. D., & Leszcz M. (2005). *The theory and practice of group psychotherapy* (5th ed.). New York, NY: Basic Books.

Zubala, A. (2013). *Description and evaluation of arts therapies practice with adults suffering from depression in the UK* (Doctoral dissertation). Retrieved from <https://eresearch.qmu.ac.uk/handle/20.500.12289/7400>

Zubala, A., MacIntyre, D. J., Gleeson, N., & Karkou, V. (2013). Description of arts therapies practice with adults suffering from depression in the UK: Quantitative results from the nationwide survey. *Arts in Psychotherapy, 40*(5), 458–464. <https://doi.org/10.1016/j.aip.2013.09.003>

Zubala, A., MacIntyre, D. J., Gleeson, N., & Karkou, V. (2014). Description of arts therapies practice with adults suffering from depression in the UK: Qualitative findings from the nationwide survey. *Arts in Psychotherapy, 41*(5), 535–544. <https://doi.org/10.1016/j.aip.2014.10.005>

Table 1*Development of Practice Principles Through Initial Meetings and Therapist Workshops*

	Initial meetings	Workshops	Final grouping
Practice principles	Proscribed principles		
Non-judgemental, validating attitude Empathy, unconditional positive regard, congruence of therapist, modelling for the group	Over-sharing. Therapist as a 'blank slate'.	Use of therapist self-disclosure as a means of building trust and sense of being understood	1. Engagement, interest and warmth
Modelling curiosity and interest in clients' experience of the modality (as opposed to interpreting)	Direct interpretation without checking client's understanding of events or ignoring client's communicated understanding of events	Being with what is. Cultivating curiosity.	2. Exploration, curiosity, not-knowing stance
Use of the art form and stance to meet the person/group where they are at.	Not intervening when expression is uncontained	Letting go of tension.	3. Active and dynamic attunement
Supporting self-regulation through containment and transformation of expression		Safe space for feelings to be expressed.	4. Arousal regulation
The therapist adapts their stance and involvement with the group depending on whether the group is in early, middle or late stages.		Therapist stance is more active in early stages. Therapist becomes less directive but is ready to resume directions if the group requires.	5. Developmentally informed group leadership

	Initial meetings	Workshops	Final grouping
	Practice principles	Proscribed principles	
Importance of supportive and effective co-working.			6. Cooperation between therapists
Modelling reliability and trustworthiness through actions	Rigid adherence to boundaries without collaborating or seeking mutual understanding with clients	Promoting safety. Elastic/flexible boundaries to be negotiated as a group. Fluidity. Boundaries as a group container and structure. Therapist responsibility.	7. Managing group boundaries
Early stages focus on building trust, safety and cohesion. Later stages, group autonomy is promoted, with the therapist becoming more directive to prepare for group ending.	Not intervening/ being completely non-directive in the group Therapist directs- does not give group opportunity to lead or direct	Make changes led by client.	8. Regulating group phases
Open and equal relationship to model safety and ways of relating in the group.	Once cohesion and trust are established, not challenging group to try out new things or take risks	Group cohesion. Commonality of experience. Arts promote a sense of belonging – community and friendship.	9. Active promotion of group cohesion
Encouraging sharing and helping between group members	Contributions ignored or dismissed by therapist		10. Caring for the group and each member
Encouraging active participation, exploration and use of the art form	Not assisting or intervening when the client is unable to use the art form	Creativity. Enabling participation. Potential space, possibilities.	11. Modelling use and potentials of art-form

	Initial meetings		Workshops	Final grouping
	Practice principles	Proscribed principles		
Encourage self-expression and creativity		Allowing the group to avoid or not use the art form	Sharing of different cultures Play, Enabling participation.	12. Encouraging engagement and exploration of the art-form
Encourage exploration and new ways of using the art form.		Persisting in an activity which is not appealing or helpful to the group Allowing long periods of talking or silence without any use of the art form.	Not dogmatic but coherent. Practice evolves with group life.	13. Developing and expanding use of the art-form
Highlighting group interactions- encouraging clients to make their own meaning from this and try out new ways of relating		Problem solving for the group	Noticing patterns → changing patterns. Acknowledging trauma and working through. Exploration of differences.	14. Focus on personal and group issues
Emotional involvement in the creative process		Therapist over-involvement in emotional processes	Working with what is within Experiencing self differently.	15. Focus on implicit experiences (arts-based, embodied, emotional)
Relating all interactions in the sessions back to the here and now Develop relationships that are shared, equal, collaborative, respectful, mutually trusting Use of transference to understanding evolving relationships within the group		Focus upon past events without connection to here and now of the group	Making connections Breaking isolation Social contact Peer support Practising interactions Shared experiences Communication. Acknowledging diversity and difference.	16. Focus on relationship between therapists and patients
Help service users to discover personal meaning and understanding from the arts process			Therapist is dynamic and moves fluidly between verbal and nonverbal ways of relating	17. Encouraging integration of arts, implicit and relational experiences in the group

Initial meetings		Workshops	Final grouping
Practice principles	Proscribed principles		
		<p>A place where emotions are ok.</p> <p>Integration</p> <p>Connecting with different parts of self, integrating using creative means.</p> <p>Other ways of seeing self/others.</p>	
Support service users to build on useful and helpful experiences of using the art-form within the group by identifying ways this can be continued in their day-to-day life.		Use of arts in own life outside of therapy for resilience/enjoyment.	18. Integrating use of art-form inside and outside of sessions
Offering encouragement and facilitating/fostering positive experiences	<p>Focus only on deficits or problems</p> <p>Avoiding problems</p>	<p>Building confidence and self-esteem.</p> <p>Resilience building.</p>	19. Acknowledging and encouraging strengths, preferences and positive experiences
Collaborating with group members regarding goals and methods of working			
Focusing upon group members' strengths and potentials, recognising clients' competence in the creative process and unique creative identity			

Table 2*Clinical Scenarios Where Therapists Make an Observable Therapeutic Action*

Item	Definition
1 Unmanageable affect	High / low affect that impacts upon ability to relate to or be with the art form/others and is difficult to self-regulate
2 Power dynamics	Interpersonal dynamics, criticism, scapegoating, blame, dominance, boundaries, them and us
3 Collaboration	Empathy, attunement, relating, working with others, co-production of the art form
4 Ambivalence about using arts media	Difficulty using art form due eg. to lack of experience, current feeling state, self-consciousness in front of others
5 Using the arts for retreat/distraction/avoidance	Avoiding contact with others by focusing solely on the arts medium, avoiding connection to feelings
6 Identity	Artistic biography, narratives, cultural associations, rites of passage, rituals
7 Linking arts process with personal relationships and community	Making connections between use of the art form and relationships within/outside the group.
8 Improvisation	Authentic, new artistic element, expression, play, creativity in use of art form in the moment
9 Incongruence between art form and feelings	Expression within the art-form does not match the presentation or stated affect of the group/group member
10 Concrete, rigid, stuck	Set ways of using the art-form, lack of flexibility, lack of connection to feelings and interpersonal process.
11 Sub-groups	Pairing, trios, quartets
12 Diverse uses of the arts forms	New ways of using the arts media eg. playing a chair with drumsticks, finding a different movement, different styles and genres of the art form introduced.
13 Over-compliance / need to 'fit in'	Taking a background role in the group, using the arts to hide from/avoid challenge by others. Working to please the therapist.
14 Pre-occupation with an 'aesthetic' or competence	Valuing the artistic product over and above the arts/group process. Focusing on ability/competence to use the arts without relating back to self and situation.

Table 3

Therapist Principles for Practice

Therapist principles for practice

<p>1. Engagement, interest and warmth The therapist conveys empathy, authenticity, positive regard, offers encouragement and is emotionally involved both during and outside the creative process.</p>
<p>2. Exploration, curiosity, not-knowing stance Overall stance of the therapist in terms of active listening, modelling curiosity and interest and conveying a stance of 'not-knowing'.</p>
<p>3. Active and dynamic attunement The therapist actively meets and attunes to the presenting internal state of each individual and the group as a whole through use of the art-form, body language and words.</p>
<p>4. Arousal regulation The therapist uses attunement and interactions within relationships, primarily with the art-form, to help clients to regulate arousal levels, paying attention to under/over-arousal to establish a calm and reflective holding presence. The therapist later supports clients to carefully expand upon and return to different arousal states. The therapist only challenges the client/group once safety and cohesion are established.</p>
<p>5. Developmentally informed group leadership The therapist's style of group leadership changes depending on the developmental level of the group with greater activity at the beginning and end of the group process.</p>
<p>6. Cooperation between therapists The therapist and co-facilitator work together to establish a safe space, model relational capacity and cooperation through confident, transparent and open communication.</p>
<p>7. Managing group boundaries Active and dynamic management of the physical space and group boundaries. Supporting and encouraging attendance, maintaining contact, modelling reliability and trustworthiness through actions.</p>
<p>8. Regulating group phases The therapist is active in helping the group establish and move between beginning, middle and end stages of the session and process as a whole.</p>
<p>9. Active promotion of group cohesion The therapist works actively to promote a sense of group wholeness and belonging through arts-based activities and discussion especially in early stages of the group.</p>
<p>10. Caring for the group and each member The therapist provides a calm, reflective and holding presence; promotes shared, equal, collaborative and mutually trusting relationships. Actively addresses destructive behaviour or threats towards the group.</p>
<p>11. Modelling use and potentials of art-form The therapist demonstrates ways to begin, and use the art-form.</p>
<p>12. Encouraging engagement and exploration of the art-form The therapist supports members to overcome anxiety in use of arts materials and develop curiosity as to what is possible.</p>
<p>13. Developing and expanding use of the art-form</p>

The therapist supports and begins to challenge members to play creatively to develop and expand their use of the art-form.

14. Focus on personal and group issues

The therapist supports group members to actively explore personal and group issues raised in the here and now of the group.

15. Focus on implicit experiences (arts-based, embodied, emotional)

The therapist draws attention to the embodied or felt qualities of using the art-form through arts-based or verbal responses.

16. Focus on relationship between therapists and patients

The therapist seeks to develop shared, equal, mutually trusting relationships, highlights group interactions and encourages clients to make their own meaning/try out new ways of relating; encourages clients to learn from one another.

17. Encouraging integration of arts, implicit and relational experiences in the group

The therapist encourages connection of bodily and arising emotional/felt states to the process of engaging with the art form and helps to make connections from the use of the arts to relationships within the group.

18. Integrating use of art-form inside and outside of sessions

The therapist promotes and explores connections between arts, group experiences and everyday life.

19. Acknowledging and encouraging strengths, preferences and positive experiences

The therapist nurtures positive experiences through collaboration and acknowledgement of the group's existing strengths and preferences.

The arts therapies are forms of psychotherapy that involve active participation in an art-form with an arts therapist to facilitate the therapeutic process through creative, non-verbal and verbal expression. Although in many countries arts therapies often have separate and varying training programmes, professional registration and recognition, they can also be regarded as sharing enough common characteristics to be part of joint clinical and research tasks (Karkou & Sanderson, 2006; Zubala, MacIntyre, Gleeson, & Karkou, 2014). Within the United Kingdom, art therapy, music therapy and dramatherapy are regulated by the Health and Care Professions Council (HCPC, 2013), while dance movement psychotherapists can achieve registration with the UK Council for Psychotherapy (UKCP, 2012). As with other forms of therapy, the arts therapies can be provided on an individual or group basis and there are many models of practice that have evolved from, and are tailored for different clinical groups and settings (Karkou & Sanderson, 2006). This diversity of practice enables the arts therapies to flexibly meet a range of health and care needs, but can make wider assessment and generalisation of their effectiveness challenging.

There is a rich body of arts therapies practice in mental health (Brooke, 2006; Hogan, 2001; Jones, 2007; McCaffrey, 2016; Nemetz, 2006). However literature tends to focus upon a single arts modality and in effectiveness studies, a specific diagnosis, which does not reflect actual provision (Fenner, Abdelazim, Bräuninger, Strehlow, & Seifert, 2017). Until relatively recently, there was also little consideration of the commonalities and differences of practice (Jones, 2020; Karkou & Sanderson, 2006) which limits our understanding of the factors shared across arts therapies and the practice principles that are important to apply in our work. This may be to some extent, historical. For example, within the United Kingdom, there were initial concerns of individual professions being grouped under one umbrella term of 'arts therapies' (Barrington, 2005) due to feared loss of identity and the

notion that approaches were too different or poorly understood to be considered together (Karkou & Sanderson, 2006; Waller, 1991). Conversely, there are many examples of arts therapists collaborating and building a shared understanding across modalities (Colbert & Bent, 2018; Havsteen-Franklin, Jovanovic, Reed, Charles, & Lucas, 2017; Huet, 1997; Knill, 1994; Landy, 1995; Purdie, 1993; Sajnani, Marxen & Zarate, 2017; Samaritter, 2018).

This paper presents developmental work to produce a model of group arts therapies practice for patients with different diagnoses (diagnostically heterogeneous) in community mental health care. The aim was to sufficiently identify, agree and describe core features of group arts therapies practice to produce a manual in preparation for a randomised controlled trial involving art therapy, dance movement therapy and music therapy (the ~~xxERA~~ study, ISRCTN:88805048-xxxxx). Whilst the trial does not involve dramatherapy due to the logistics of the study design and local resources, we were keen to ensure the model could represent the full spectrum of arts therapies practice. Dramatherapists were therefore involved in the development process. In line with guidance for the development of complex interventions (Bleijenberg et al., 2018; Cotterill et al., 2018; Craig et al., 2019; Hoffmann et al., 2014; O’Cathain et al., 2019; Rousseau et al., 2019; Turner et al., 2019), our objectives were to (a) identify and agree upon theories that underpinned the intervention; (b) identify commonalities in the provision of art, dance movement and music therapy groups; and (c) develop a manual and set of principles to enable adherence rating in practice.

Considerations in Manualising Arts Therapies

Treatment manuals can be thought of as a means to describe the core “conceptual and structural boundaries” of how a treatment is provided and works (Addis, Cardemil, Duncan,

& Miller 2006, p.132). Initially developed for research (Luborskey & DeRubeis, 1984), manuals are also adopted or developed for clinical practice (Addis et al., 2006). Within randomised controlled trials, the purpose of a manual is to sufficiently describe the aims, methods and hypothesised mechanisms of the treatment(s) under investigation (Grant et al., 2018).

Manuals enable therapists to understand what is important to happen and how, and to monitor how far their practice aligns with this (Addis et al., 2006), whilst they enable researchers to check whether what is hypothesised to be of importance is actually provided across all participants (treatment fidelity) (Borrelli, 2011; Nelson, Cordray, Hulleman, Darrow, & Sommer, 2012). This enhances internal validity of the study, and enables future replication and comparison with different treatments (Nelson et al., 2012). The relative importance of practice elements to outcomes can also be better specified and understood (Kraemer, Wilson, Fairburn, & Agras, 2002; Nelson et al., 2012). As Rolvsjord, Gold and Stige (2005) note, use of a manual does not “guarantee valid results, nor will all results of studies without therapy manuals necessarily lack validity” (p.22). Manualisation does not aim for or guarantee superiority over non-manualised treatments (Truijens, Zühlke-van Hulzen, & Vanheule, 2019; Tshuschke et al., 2015); nor are novel manualised procedures easily adopted in routine practice (Bitsika & Sharpley, 2006; Carroll & Nuro, 2002; Chorpita, 2006).

Most arts therapies research manuals draw upon existing psychotherapy approaches, or wider theories to develop new interventions for specific populations, with a focus on one single arts modality (Azoulay & Orkibi, 2015; Behrends, Müller & Dziobek, 2012; Bryl & Goodill, 2019; Hackett, Taylor, et al., 2017; Parkinson & Whiter, 2016; Rolvsjord et al., 2005; Story & Beck, 2017; Zubala, 2013). Others have used intervention mapping (Bartholomew, Parcel, Kok, Gottlieb & Fernández, 2011) as a development method (Aalbers et al. 2019;

Haeyen & Heijman, 2020; Haeyen, van Hooren, Dehue & Hutschemaekers, 2018). Those manuals that have incorporated practices across the arts therapies focus on a multi-modal arts approach with a distinct client population such as depression (Parsons et al., 2020), or dementia (Lyons, 2019). One intervention has been developed across different arts therapies with children in mainstream schools (Moula, Karkou, & Powell, 2019). However, None so far have developed manuals across different arts therapies modalities for diagnostically heterogeneous groups in mental health care.

However, Within a number of psychotherapy, there are a number of manuals ~~exist~~ for diagnostically heterogeneous groups. Notably, Barlow, Allen & Choate (2004) developed a transdiagnostic “uniform treatment protocol” for a range of emotional disorders, including adaptations for group treatment (Bullis et al., 2015; Lapos, Mancuso, Abraham, & Lolidano, 2017). Karterud (2016) similarly published a manual for mentalization-based group therapy. Both share a focus on common factors of group treatment, whilst also highlighting specific aspects of the approach.

Beyond research, a well-designed manual can also support therapists’ self-monitoring, professional development (Addis et al., 2006; Luborsky & DeRubeis, 1984) and act as a map or toolkit for therapists (Bryl & Goodill, 2019; Taylor-Buck & Dent-Brown, 2015). As such, some arts therapies professional bodies have developed clinical guidelines using consensus based methods, such as nominal group or Delphi technique (Eyre, 2013; Hackett, Ashby, Parker, Goody, & Power, 2017; Havsteen-Franklin, 2014; Springham, Dunne, Noyse & Swearingen, 2012; Taylor Buck, Dent-brown, Parry, & Boote, 2014; Taylor Buck & Hendry, 2016; Wright & Holttum, 2020).

Despite the many manualised forms of psychotherapy, studies of how such manualisation is delivered in practice highlight the challenges in doing so. Manuals are not always

accepted or understood by therapists, who may perceive them as challenging their professional identity, limiting therapeutic autonomy and creativity (Addis et al., 2006; Bergström, Ladd, Jones, Rosso, & Michael, 2019; Carroll & Nuro, 2002; Forbat, Black, & Dulgar, 2015; Truijens et al., 2019; Tschuschke et al., 2015). Implementation requires significant resource commitment (Addis et al., 2006; Chorpita, 2006) and if developed outside of the clinical context, may be limited in their external validity (Bitsika & Sharpley, 2006; Carroll & Nuro, 2002; Chorpita, 2006; Westen, 2002). Focus upon specific or novel factors of the manualised treatment can be to the detriment of known essential factors of therapy (e.g., therapeutic relationship) (Carroll & Nuro, 2002; Addis et al., 2006; Herman-Smith, Pearson, Cordiano, & Aguirre-McLaughlin, 2008); ignore the unique contribution of client and therapist (Addis et al., 2006; Rolvsjord et al., 2005) and focus upon technical execution to the detriment of training and supervision (Bein et al., 2000; Chorpita, 2006). This can have unintended and potentially negative consequences for both client outcomes (Henry et al., 1993; Vakoch & Strupp, 2000) and therapists' practice (Forbat et al., 2015).

Manuals can also fail to recognise the variation in therapist training, theoretical allegiance(s), experience, and competence (Crameri, Tschuschke, Koemeda, Schulthess, & von Wyl, 2020; Najavits, 1997; Tschuschke et al., 2015). For process-oriented therapies, the idea of following a predefined formula can be at odds with responding to the needs of the patient (Rolvsjord et al., 2005; Taylor Buck & Dent-brown, 2014; Vakoch & Strupp, 2000). Manuals may also challenge long-standing practice and require integration, building of confidence and skills (Najavits, 1997; Sholomskas et al., 2005). Research procedures and restrictions may also impact upon therapists' performance and confidence (Gold, Erkkilä & Crawford, 2012; Robb, Burns, Docherty, & Haas, 2011).

Manuals therefore require a fine balance of specification and flexibility. This includes careful consideration of common therapeutic factors (Beutler, 2006; Bradt, 2012; Rolvsjord et al., 2005; Waltz, Addis, Koerner, & Jacobsen, 1993; Westen, 2002), practitioner involvement, clinical context (Bryl & Goodill, 2019; Carroll & Nuro, 2002; Chorpita, 2006; Forbat et al., 2015; Sidani & Braden, 2011; Westen, 2002) and therapist training and support (Butler, Henry, & Strupp, 1995; Carroll & Nuro, 2002; Gold et al., 2012; Robb et al., 2011). Studies of manual implementation have highlighted that therapists tend to work ‘tacitly’ or ‘implicitly’, responding to the person in the moment without thinking explicitly about the action being performed (Lee & Vakoch, 1996; Najavits, 1997; Vakoch & Strupp, 2000). Reber notes that within implicit learning “...not only is the learner unaware of the process of learning, the very knowledge itself is highly resistant to explication” (2001, p.15433). This may be even more important for arts therapists, given their frequent work within a non-verbal, implicit domain and the association of such processes with creativity (Gerge, Hawes, Eklöf, & Pedersen, 2019; McAleer, Bowler, Bowler, & Schoemann, 2020).

___Studies have also demonstrated that overarching principles are more likely to be understood and implemented rather than listing predefined techniques (Lee & Vakoch, 1996; Vakoch & Strupp, 2000). Accordingly, some arts therapists have based manuals on Waltz, Addis and Koerner’s (1993) framework of principles for practice, rather than rules or procedures (Bryl & Goodill, 2019; Rolvsjord et al., 2005). In order to have an effective learning experience, Lee & Vakoch (1996) suggest that implicit knowledge first needs to be made explicit. Therefore, techniques to explore the tacit/implicit way in which therapists work with groups are important to draw out in practice. Given their non-verbal and creative possibilities, the arts are an effective medium to achieve this. We therefore used a range of arts and practice-based methods in the manual development of this study.

Method of Manual Development

Manual development began with monthly meetings between six arts therapists: two each of art therapy, dance movement therapy and music therapy. The arts therapists each had backgrounds covering arts therapies practice, academic research and strategic or managerial positions within a range of NHS mental health trusts. Our initial meetings highlighted challenges in learning one another's separate arts therapies languages and finding a shared understanding in the theories, tools and techniques we used. These meetings informed development of the underpinning theory and model which enabled the group to explore practice. Subsequent meetings focused upon defining principles of practice according to Waltz et al.'s framework (1993). Each arts therapist listed principles for practice according to their discipline and the techniques that would be used to achieve this. These were discussed as a group and formed the basis of 10 core principles of the manual, 18 principles for practice and 15 proscribed principles (Table 1).

We then used nominal group technique (Boddy, 2012; Harvey & Holmes, 2012; Havsteen-Franklin, 2014; Taylor Buck & Hendry, 2016), facilitated by DHF[xx] and CC[xx] to explore common clinical scenarios encountered within community mental health arts therapies groups. The focus of this technique was to describe common scenarios in group arts therapy where the therapist would make an intervention (defined as an observable action made by the therapist). It was highlighted that interventions could be both to address a clinical need or deficit, or to promote, encourage or highlight strengths and resources. Each group member listed as many scenarios as possible, over the course of 20 minutes. These were written up and displayed on flipchart paper around the room. From a list of 90 scenarios, these were then grouped thematically, resulting in a list of 14 scenarios (Table 2). In a final

step, group members individually ranked the importance of each scenario based on their experience in practice. Individual ranking scores were combined to produce an overall group ranking.

Wider Arts Therapist Involvement and Feedback

Once the ten core principles, principles for practice and clinical scenarios were developed, we ran a 1-day workshop in two NHS Trusts, with 19 arts therapists in total. Attendees were arts therapists who had expressed an interest in running arts therapies groups in the forthcoming [ERA\[xxxx\]](#) study. One person with lived experience of mental health problems advising on the study and two dramatherapists also participated.

Arts-based methods were used to explore therapists' stances towards the level of directiveness used in sessions and group boundaries (Brabender & Fallon, 2018; Dies, 1985). For this, we used a movement-based exercise, where therapists explored the space between two polarities (e.g., active vs passive) in the room and then discussed the reasons for positioning themselves where they did. This led to reflections on instances where greater activity from the therapist, or less might be appropriate. Therapists explored how they established safety in early sessions by working with a range of arts objects and materials (e.g., scarves, wooden blocks, paper and pens), and core principles from the ensuing discussion were recorded on a flipchart. Group processes were explored through a music drawing narrative exercise (Booth, 2005) facilitated by [CC\[xx\]](#), resulting in artistic representations of common group processes and a written reflection by each therapist. The experience of this exercise was discussed as a group and core themes from this discussion were written down. The group then discussed the aims of group arts therapy, contributing drawings and written words to large pieces of paper. The group reviewed 14 clinical

scenarios developed to date and broke into small groups to role-play one of these scenarios as they had experienced in their own practice. This role play was specific to each arts modality with participants taking turns in the role of therapist. As a final step, the arts therapists were asked to rank the importance of the 14 clinical scenarios, to provide additional feedback on the relevance of each scenario. Following these workshops, thematic analysis led to development of five core group aims and additional principles for practice (Table 1). The 10 core principles were unchanged, but were elaborated from workshop discussions to illustrate ‘What this is’ and ‘What this isn’t’ (following Rolvsjord et al., 2005).

Refinement and Selection of Principles for Practice

To refine the principles for practice, each principle was listed (20 from the main principles, 20 from promoted practices and 27 proscribed practices) and grouped thematically, forming 19 core areas. Core areas were then compared against existing group art, music and dance movement therapy literature in mental health and a group mentalization therapy adherence scale (Folmo et al., 2017; Karterud, 2016). In grouping the items, a number of principles shared amongst group therapies were identified. Where this was the case, the item naming, as entitled by Karterud (2016) was retained (items 1, 2, 6, 7, 8, 10). This formed the final structure for core principles and adherence rating.

Results

Underpinning Theory

Transdiagnostic Theory and Contextual Model of Psychotherapy

We took as our starting point, transdiagnostic theories of mental illness (Bakker, 2019; Casey, Olivery, & Insel, 2014; Sanislow, 2016). Whilst no single model exists, current theory suggests concepts of negative valence (e.g., acute threat/fear, loss), positive valence (e.g.,

motivation, reward), cognition, social processes, arousal and regulation are relevant to understanding shared features of mental illness (Cuthbert & Insel, 2013). There is particular interest in the transdiagnostic role of emotion regulation, with difficulties in using adaptive regulation strategies, intolerance of uncertainty and experiential avoidance as three potential mechanisms (Aldao, 2012; Khakpoor, Bytamar, & Saed, 2019). Whilst the course and experience of mental illness is unique to each individual, within secondary mental health care, these transdiagnostic concepts are reflected in common challenges faced by patients in terms of managing and regulating mood and anxiety, maintaining motivation in daily life and becoming stuck in patterns of thinking, feeling or relating, with associated emotional and relational distress (Herrman, Saxena & Moodie, 2005; World Health Organisation, 2001). Relationships can be further affected through challenges in communicating experiences, thoughts and feelings making them difficult to develop or sustain (Brüne et al., 2009; France & Kramer, 2001; Kerns, 2007; Perlini et al., 2012; Sánchez et al., 2019). Many experience social isolation, stigmatisation and lack confidence to socialise and wider effects of disempowerment, loss of identity, role and self-esteem (Baker, Jodrey, & Intagliata, 1992; Goldberg, Rollins, & Lehman, 2003; Halford & Hayes, 1995; Thornicroft et al., 2016; Trompenaars, Masthoff, Van Heck, de Vries, & Hodiament, 2007).

We then drew upon the contextual model of psychotherapy (Wampold, 2015; Wampold & Imel, 2015), which suggests that features shared across different types of therapy are the most important factors that account for clinical change. Determinants include the client's preference for, understanding and expectations of therapy and the therapist's experience, commitment and belief to the model of therapy provided, along with a positive working relationship (Wampold, 2015; Wampold & Imel, 2015). In line with the contextual model, a meta-analysis showed that for mental health patients, receiving a preferred treatment is

associated with lower dropout from treatment and higher therapeutic alliance (Windle et al., 2019). A further meta-analysis across group therapies has demonstrated that common shared factors account for the greatest amount of change in therapy, rather than unique factors (Orfanos, Banks, & Priebe, 2015). These common group factors have been studied in various combinations, but include, at a meta-level; supportive factors (instillation of hope, acceptance, altruism, universality and cohesion); self-reflection (including self-disclosure, catharsis and secure emotional expression), psychological work (including interpersonal learning, self-understanding and awareness of relational impact) and social learning (including modelling, vicarious learning, guidance and education) (Mac Nair-Semands, Ogrodniczuk, & Joyce, 2010; MacKenzie, 1990; Yalom & Leszcz, 2005). On the basis of the above, we can assume that many of the important therapeutic factors in group arts therapies are those shared with wider forms of group therapy as a whole.

More recent research (Orfanos & Priebe, 2017) suggests that within closed community therapy groups, the first few sessions are most important in determining outcomes. Sessions where group members showed high cohesion in the first sessions predicted future engagement and greatest therapeutic gains. Reporting similar findings for different types of group psychotherapy, Tchuschke and Dies (1994) proposed that such early group integration promotes capacity for self-disclosure, which increases interpersonal feedback thus increasing opportunities for positive feedback from the group. Burlingame, Fuhriman and Johnson (2001) highlight the importance of certain practices in developing cohesion, namely: Thorough preparation of members for the group experience, identification of useful processes early on in therapy, consistency in temporal and spatial features of the group, fostering member to member interaction, feedback to the group, a warm and caring therapeutic stance and ensuring members relate to each other in positive ways. Within the

arts therapies, a positive association has also been identified between therapeutic alliance and symptom reduction (Heynen, Roest, Willemars, & van Hooren 2017).

___The above suggests that important factors within group therapy are patient understanding and clear expectations of what will happen in therapy, patient and therapist shared commitment to working in a particular way and time to develop and establish a good working alliance. Within groups this highlights the importance of patient treatment preference, clear information on what to expect, fostering group cohesion in the initial phases and promoting active engagement quickly and early on in the therapeutic process.

The Additional Contribution of the Arts in Therapy

The defining factor of the arts therapies is emphasis upon the use of, and active participation in the arts during therapy. Koch (2017) suggests five factors specific to the arts therapies: aesthetics (beauty and authentic expression), hedonism (pleasure and play), meaning making (through non-verbal communication/symbolisation/metaphor), enactive transitional support in times of change, and generativity (creativity and creation). Inclusion of an active and creative arts-based process provides opportunities for embodied self-expression, creativity and a non-verbal means of relating with other people (Samaritter, 2018). The art form brings a third object- concrete (such as a recording or piece of art work), or experienced (such as movement or heard sound)- into the matrix of relations within the group (Karkou & Sanderson, 2006). Each person brings their own unique creative and cultural identity. This can be explored as part of the therapy with the potential to link and expand upon their personal relationship with, and use of the art-form as a 'helping resource' in their day-to-day lives (Ansdell & Meehan, 2010).

___Springham and Huet (2018) suggest art-work in art therapy may be considered a second

order representation within psychological processes (representing alternative ‘as if’ scenarios, rather than reality). This provides a means of trying out different ways of relating to oneself and others in an imaginative and safe way. Arts therapies are therefore well placed to help patients identify difficulties and strengths through the use of the art-form in the varying group interactions. Use of the art-form can facilitate the experience and expression of emotions and discovery of personal meaning in creative activities, allowing experiences and learning through non-verbal and verbal communication (Gerge et al., 2019).

___ In other words, through creativity, imagination and play, patients are helped to explore new or different emotional and cognitive experiences with support of both the group and the therapist. The production of a piece of art is often cited as a means of strengthening self-esteem by clients (Buck & Havsteen-Franklin, 2013; Grocke et al., 2014) and can provide access to personal and interpersonal resources which may be continued into wider daily and creative life (Ansdell & Meehan, 2010; McCaffrey, Carr, Solli, & Hense, 2018; McCaffrey, Edwards, & Fannon, 2011).

Historically, arts therapists have integrated wider psychotherapy theories including expressive, psychodynamic, person-centred, attachment and group-interactive models (Karkou & Sanderson, 2006; Odell-Miller, 2013; Springham & Huet, 2018; Zubala, MacIntyre, Gleeson, & Karkou, 2013). Principles from each of these informed our understanding, along with principles of resource orientation (Priebe, Omer, Giacco, & Slade, 2014; Rolvsjord, 2010; Rolvsjord, 2015) and recovery (Leamy, Bird, Le Boutillier, Williams, & Slade, 2011; McCaffrey, 2016; McCaffrey et al., 2011, 2018; Solli, Rolvsjord, & Borg, 2013).

___ The principles of person-centred therapy (Rogers, 2011/1967) are especially relevant to how the therapist models and interacts with members of the group involving a non-

directive, empathic stance. The psychodynamic model (Montgomery, 2002) provides a means of understanding patterns of relating and emotional responses across unconscious and conscious levels, which facilitates understanding of non-verbal and implicit levels of communication. Resource and recovery-oriented approaches (Ansdell & Meehan, 2010; Buck & Havsteen-Franklin, 2013; McCaffrey et al., 2018; Rolvsjord, 2010) are informed by a move towards a collaborative, contextual and health-promotion focus rather than a deficit or treatment-based one. Principles include a shared, equal, collaborative process; focus on patients' strengths and potentials; acknowledging the patients' creative identity; being emotionally involved in the creative process and fostering positive emotions (McCaffrey et al., 2018).

Implications for the different arts modalities:

Whilst we hypothesise that it is the shared, or common factors of group therapy that are associated with eventual change of health outcomes, each arts modality has specific multi-sensory and aesthetic properties which can provide unique opportunities within a group context (Malchiodi, 2005). Whilst in simple terms a single sensory modality is implied (art, the eye; music, the ear; dance movement, the body), a range of senses are activated, offering many different sensory and embodied modes of expression and experience (Gerge et al., 2019; McNiff, 1981). Stern (2010) and Malloch and Trevarthen's (2009) studies of human communication show how timing, shape and intensity ("vitality affects") are used to non-verbally communicate intention and regulate interaction from birth. The multi-sensory and multi-modal features of arts-based work can facilitate attunement and regulation of basic physical and behavioural experiences, such as relaxation, self-soothing, and building of healthy attachments (Behrends et al., 2012; Metzner et al., 2018; Springham & Huet, 2018).

In our model (Figure 1), the importance of the art form is the appeal to the patient (their preferred type of arts therapy), the facilitation of active participation and emotional engagement in the use of the art form, the introduction of creativity, and the support of exchange and interactions. The final therapeutically effective processes however are non-specific to the art form and will benefit from the diagnostic heterogeneity of the group. Figure 1 outlines how transdiagnostic theories of mental illness (orange), informed choice (yellow), common factors to group therapies (green), resource and recovery oriented theories (pink) and specific arts therapies processes (purple) relate within this model. We contextualise the arts-based processes within arts therapies groups as being an additional process to those that already occur within talking-based group therapy.

Manual

The manual encompasses five group aims, 10 overarching core principles, 14 clinical scenarios and 19 principles for therapist practice. The group aims relate directly to shared features of mental illness, whilst the overarching principles describe core features of how the groups should be managed over time. The clinical scenarios are used as a springboard for therapists to reflect on common features of running arts therapies groups. We purposefully did not define interventions for these scenarios as (a) many scenarios can unfold simultaneously and are not necessarily distinct; and (b) there are a multitude of ways that such scenarios might be addressed. Rather, we defined 19 principles for therapist practice which enable therapists to observe themselves and others when in action and form the basis of adherence rating for the [ERA\[xxxxx\]](#) trial.

Group Aims

Our discussions identified five core group aims of arts therapies groups for community mental health settings. Importantly, these aims incorporate both implicit (embodied, felt or lived) and explicit (verbal and reflective) relational experiences and build from basic pre-requisites for therapy and the individual, to wider relational and community goals:

1. To provide a safe space to be with others

2. To promote engagement with the art form to

explore ways of using the art form and its relevance to one's own life;

explore inner creativity;

explore opportunities to extend and use the arts in daily life.

3. To provide regulation of arousal and affect to

explore ways of using the arts to aid self-regulation;

become aware of and find new ways of expressing and transforming emotions.

4. To offer opportunities to build upon existing relational capacity to

build awareness of self and others;

facilitate non-verbal engagement and expression with others;

explore new and different ways of relating to others;

break isolation and offer opportunities for support and socialisation.

5. Through the above, use the arts to build confidence, self-esteem, self-agency and autonomy in interactions with others and to further develop connectedness, hope, identity, meaning and empowerment (recovery) (Leamy et al., 2011).

Ten Core Principles

The ten core principles set the overall structure and format of groups over time. Items

one to three highlight important actions to be taken by the therapist in preparation and support for the groups (such as non-attendance and team communication). Item four defines the general group structure for a single session, whilst items seven to nine describe the overall group task at beginning, middle and end stages of therapy. Items five, six and ten highlight the importance of active arts participation, the therapist's role in supporting the group to become familiar with this and the importance of maintaining a collaborative and strengths-based approach, which is in keeping with a recovery focus.

1. The arts therapies groups are fully explained with information and video material so that patients can make an informed choice as to their preference

What this means: For the [ERA~~xxxxx~~](#) study, potential members are shown video clips to aid decision making for the art-form that they most prefer. The videos explain what happens during an art therapy, dance movement therapy and music therapy session. Through the voices of service users, the videos cover what to expect, examples of activities during sessions and helpful and challenging features. Prior to the group start, an individual meeting is arranged with the therapists in the group therapy room, to provide a sense of what the space will look like. The meeting focuses upon introductions, the reasons for coming to therapy (including hopes and expectations) and explanation of the general aim, structure and process of group therapy. There is an opportunity to ask questions, agree ground rules and expectations, including the level of communication the person wishes to have.

What this is not: Insisting upon a particular modality choice that negates the patients' preference. Declining to discuss arts therapies options or concerns.

2. Attendance is encouraged and supported through maintaining active contact with patients throughout the treatment phase

What this is: Strategies to support attendance are discussed and agreed in the initial meeting and can include text reminders, and agreed modes of communication for non-attendance. Within the group, therapists acknowledge group absences and ensure time to welcome and re-orient returning members and late arrivals.

What this is not: Persistent calls to demand why the person is unable to attend. Ignoring members' preferences for attendance support. Ignoring group absences and their impact.

3. The therapists ensure communication is maintained and held with the multi-disciplinary care team throughout.

What this means: Therapists ensure ongoing communication with the wider care team to inform them of arising issues or concerns and to ensure high quality of support and care. The level of communication is outlined and agreed in the pre-group meeting, including the nature of confidentiality and boundaries of what will be shared with the wider care team.

What this is not: Ignoring group members' preferences for team communication. Not sharing important information regarding risk, safeguarding or wider care support.

4. Group structure always contains: (a) an opening warm-up; (b) use of the art-form, with space for verbal reflection; (c) a closing circle to reflect upon the group experience.

What this means: Sessions have an element of predictability, giving space for individuals to share at the beginning and end of the group. The middle section is flexible and the amount of alternation between using the art-form and talking may vary across modalities. Opening and closing sections should be led by the therapist to start, but may be developed or expanded upon by the group as sessions progress. The amount of time given to each

section will vary depending on the stage of the group (early/later on in the process) and how group members are on any given day.

What this is not: Rigid reliance on exactly the same opening and closing activity each week. Not following the lead of the group in when to transition from and to activities.

5. Active participation, exploration and use of the art-form is encouraged.

What this means: Therapists aim to begin using the art-form with group members as quickly as possible in sessions, particularly where members are reluctant, reticent or avoidant. Use of the art-form should be introduced sensitively, with a level of activity and degree of structure that fits the presentation of the group. The therapist should ensure that the group does not go for long periods of talking or silence, without any use of the art-form.

What this is not: Forcing members to use an art-form when they are not ready to do so – a single group member might not directly use the art-form themselves, but may watch attentively to the rest of the group’s creations. Ignoring the current situation of a group member which may make use of the art-form difficult on that day.

6. Therapists always offer structure within the art-form to promote group cohesion and only let this become free when there is space and safety to do so.

What this means: Structure is always offered to begin to establish an atmosphere of safety and cohesion between members within the group. Time should be allowed for group members to explore the art-form within the safety of such structures so that self-agency and competency in using the art-form are established.

What this is not: Limiting the level of freedom and activity in sessions to rigid or predefined structures. Not varying the level of structure according to presenting needs.

7. Within the early phase of therapy (weeks 1-5), the therapist focuses upon building trust and active engagement with the art-form, allowing plenty of time for this to happen.

8. Middle to late sessions (weeks 6-17): Developing content and arts-based activities in collaboration with service users to build variety (as opposed to doing the same arts-activity each session).

9. End phase (weeks 18-20): The group reflects on the process as a whole and review to prepare for ending. Therapists signpost patients to wider arts-based opportunities in the community.

What this means: Within the [ERA~~xxxx~~](#) study, sessions will run for up to 90 minutes, twice per week over 20 weeks. The above phases (in weeks) are a guide and it is acknowledged that each group will have its own unique early, middle and ending processes. What is important is for the therapist to be aware of the phase of therapy in relation to overall group process, so that time is allowed for each of these stages whilst balancing group cohesion and offering opportunities to challenge or extend group activities.

What this is not: Ignoring the stage of the group and propensity to use the art form. Persisting in activities without collaborating with group members. Not returning to an earlier stage if the group requires. Ignoring the interests and needs of the group.

10. Sessions are recovery-oriented: Therapists work collaboratively with patients so that their strengths and preferences in using the art-form are acknowledged to support their future recovery and signposting.

What this is: Therapists work collaboratively with patients with a here and now focus, so that their strengths and preferences in using the art-form are acknowledged. Group

members are supported to explore issues raised, with challenges acknowledged. The therapist models curiosity in understanding members' experiences but does not immediately interpret or suggest meanings for the group. Throughout the sessions, the therapists acknowledge these strengths and preferences in terms of supporting future recovery. Therapists signpost group members where appropriate to wider community services which may meet the strengths and preferences expressed in the group.

What this is not: Ignoring risks, problems or clinical issues. Minimising or ignoring the impact of events from the past upon group members. Referring group members onto services without exploring preferences and wishes to do so together first.

Clinical Scenarios Where a Therapist Would Make an Intervention

Table 2 summarises 14 scenarios where arts therapists would make an intervention or observable action within the group. The items are listed in overall ranked order with 1, being the scenario ranked as most important and 14, the least important.

Principles for Practice and Adherence Rating

The 19 principles for practice (Table 3) encompass qualities the therapist displays in the group (items 1, 2, 10), and actions the therapist should take. Many are shared with other forms of group therapy but have the potential to also take an arts-based focus. For example, items 3 and 4 are especially important within arts therapies, where the art-form can be used jointly by the patient and therapist to aid connection to others (attunement) and change arousal state. Here, the role of the therapist is to support self-regulation through containment and transformation of expression. Items 11, 12, 13 and 18 are solely arts-based, whilst items 15 and 17 combine both common and arts group processes. Items 1, 2,

6, 7, 8 and 10 relate directly to common factors outlined by Karterud, and thus retained the original names (Folmo et al., 2017; Karterud, 2016). Items 4 and 15, whilst similar to Karterud (Regulation of tension level; Focus on emotions), were renamed to better align with the transdiagnostic focus, and acknowledge the wider range of possible implicit arts therapies experiences (bodily sensation, emotion, or as represented in the art form).

Discussion

Strengths and Limitations

The manual addresses many of the challenges faced in defining a single model of group arts therapy for diagnostically heterogeneous patients. The model has a clear theoretical basis and considers the pragmatic challenges of providing a standardised framework for three different arts modalities. Most importantly, we were able to reach agreement upon core aims and practice principles between therapists from different arts modalities and training backgrounds. Our development process followed current guidance for the development of complex interventions (Bleijenberg et al., 2018; Cotterill et al., 2018; Craig et al., 2019; Hoffmann et al., 2014; O’Cathain et al., 2019; Rousseau et al., 2019; Turner et al., 2019) and incorporated feedback from arts therapists. Experiential and practice-based methods enabled direct exploration and representation of diverse but context-specific practice. Core principles are clear and simple, whilst the practice principles accommodate a range of training backgrounds. The clinical scenarios provide the first comprehensive synthesis of situations considered important for arts therapists to act within a group.

Whilst the model is pluralistic in its wider approaches, this may pose challenges for arts therapists who have trained and practise in a defined therapy paradigm. For example, arts therapists working from a formal psychoanalytic model may be challenged by the more

active and collaborative stance. Therapists with a strong person-centred leaning may be challenged by greater therapist directiveness early on in sessions. Whilst we believe that the manual encompasses current group arts therapy practice in community health care, it remains to be seen whether it has utility beyond the life of the study and within regular community-based practice. Feedback from arts therapists in the trial and further review by external bodies (for example, training providers) will assist further validation.

As the forthcoming [ERA\[xxxx\]](#) study encompasses only art, dance movement and music therapy, dramatherapists had limited involvement and thus, this model may not fully reflect aspects of dramatherapy practice. However, we hope that this may be examined separately from the [ERA\[xxxx\]](#) trial in collaboration with dramatherapists in the future. There was also limited service-user involvement, with only one service user attending the workshops. Further collaboration with and feedback from service users (both study participants and from a lived experience advisory panel) during the [ERA\[xxxx\]](#) trial will be important to understand the relevance of this way of working, including helpful and unhelpful factors.

The manual is necessarily transdiagnostic (Aldao, 2012; Cuthbert, 2014; Cuthbert & Insel, 2013; Khakpoor et al., 2019), and does not define diagnosis-specific actions or principles. However, it also does not contradict existing guidelines (Havsteen-Franklin, 2014; NICE, 2014; Springham et al., 2012; Wright & Holttum, 2020). Further work will be necessary to align where diagnosis-specific actions are indicated and required (Beutler, 2006).

Comparison to Existing Arts Therapies Practice

This manual differs to others developed so far, in that it seeks to identify shared aims and elements of practice across different arts therapy modalities and diagnoses. However, there is strong congruence between our manual and those practices outlined by specific arts

therapists and professional bodies (Eyre, 2013; Hackett, Ashby, et al., 2017; Havsteen-Franklin, 2014; Parsons et al., 2020; Springham et al., 2012; Taylor Buck et al., 2014; Taylor Buck & Hendry, 2016; Wright & Holttum, 2020). In their study of therapist descriptions of in-session therapeutic constructs, Havsteen-Franklin et al. (2017), identified ten symptomatic events and 14 bipolar constructs. These events, whilst focusing upon only symptoms, have some similarities to the 14 scenarios identified by this group including: high and low affect, acute psychosis, ambivalent responses, concrete solutions to interpersonal problems and insight. Similarly, the bipolar constructs cover similar areas of attunement, boundaries, affect regulation, directiveness, curiosity and use of the arts. Samaritter's (2018) Delphi study of music and dance movement therapists suggests five core themes of embodied presence, somato-sensory engagement, emotional engagement, non-verbal communication and intercultural involvement. The 12 steps identified as arts related activities and procedures, follow a similar pattern to those described in our 19 principles for practice; from invitation and engagement to exploration, play and performance (Samaritter, 2018). Parsons et al. (2020) and Parkinson and Whiter (2016) also describe similar aims, group structure and principles for depression and early intervention for psychosis.

The role of Experiential and Practice-Based Arts Methods in Training

Arts-based methods are gaining interest as a tool for creative exploration in clinical practice and research and were central to the manual development (Beer, 2016; Borgdorff, 2010; Geiger, Shpigelman, & Feniger-Schaal, 2020; Gerber et al., 2020; Ledger & McCaffrey, 2015; McCaffrey et al., 2020; McCaffrey & Edwards, 2015; Meltzer, 2020; Priebe et al., 2019; Sajjani, Mayor, & Tillberg-Webb, 2020; Springborg, 2020). As arts therapists, we recognised the value in working creatively with our colleagues to recognise the tensions and

congruence across training traditions and art-forms. The workshops were lively and inclusive, enabling us to capture the essence of practice across arts modalities through our group reflections on the creative arts process.

___The workshop_s informed how we provide training of therapists for the trial itself. Initial feedback suggested that the arts therapists valued opportunities to reflect upon and explore their practice. We therefore built in a strong ethos for the training to promote therapist self-exploration of how their practice aligns with the model and manual principles, alongside regular opportunities for therapists to provide their ideas and feedback. Training is provided in three whole day workshops spread over a few months.

___Day one provides an overview of the ERA~~[xxxxx]~~ trial and the theory underpinning the intervention. Three core principles (7-9) are explored by therapist pairs leading an experiential session to demonstrate how these principles would be met in their arts modality. Therapists are asked at the end of this day to begin peer observation to become familiar in being observed for research and discussing practice with another.

___Day two involves feedback from therapists' experiences of peer observation and role plays based on one of the 14 clinical scenarios. Role plays are video recorded to give therapists a sense of how recording will feel when conducting sessions and for review in training day 3. Therapists are introduced to the adherence rating scale and encouraged to begin using this to assess their current practice. Introductory interviews and study procedures are also discussed.

___The final training day covers role play of introductory meetings, how to report adverse events and adherence rating. Therapists watch excerpts of videos from training day 2. Therapists from the modality in the video rate this excerpt, with the remaining group observing and later commenting upon the overall process. Excerpts are selected to ensure

coverage of all 19 principles for practice and discussions focus upon the ways in which the principles can be observed in the videos and put into practice.

Conclusion

This is the first model of arts therapies group practice developed across three arts modalities for mixed diagnosis groups in community mental health care. Our emphasis on direct arts-based experience led to development of a rich corpus of practice. Most importantly, we were able to reach agreement of core principles between three different arts modalities. Such methods may be of wider value to arts therapists in clinical supervision and reflective practice, as well as wider clinicians and researchers. Whilst manual utility (including adherence) and model effectiveness is yet to be assessed, the methods may prove useful in developing models of practice for wider arts therapies groups and clientele.

Key words: arts therapies; practice; mental health; manual; group; intervention development

Table legend:

Table 1: Development of Practice Principles Through Initial Meetings and Therapist Workshops

Table 2: Clinical Scenarios Where Therapists Make an Observable Therapeutic Action

Table 3: Therapist Principles for Practice

Figure legend: *Figure to be printed in colour*

Figure 1: Logic model summarising underpinning theory, group aims, core principles, therapist practice principles and related scenarios.

References

- Aalbers, S., Vink, A., Freeman, R. E., Pattiselanno, K., Spreen, M., & van Hooren, S. (2019). Development of an improvisational music therapy intervention for young adults with depressive symptoms: An intervention mapping study. *Arts in Psychotherapy, 65*, 101584. <https://doi.org/10.1016/j.aip.2019.101584>
- Addis, M. E., Cardemil, E. V., Duncan, B. L., & Miller, S. D. (2006). Does manualization improve therapy outcomes? In J. C. Norcross, L. E. Beutler, & R. F. Levant (Eds.), *Evidence-based practices in mental health: Debate and dialogue on the fundamental questions* (pp. 131–160). Washington, DC: American Psychological Association. <https://doi.org/10.1037/11265-003>
- Aldao, A. (2012). Emotion regulation strategies as transdiagnostic processes: A closer look at the invariance of their form and function. *Revista de Psicopatología y Psicología Clínica*,
- Revised draft [07-10-202025 November 2020](#)

17(3), 261–277. <https://doi.org/10.5944/rppc.vol.17.num.3.2012.11843>

Ansdell, G., & Meehan, J. (2010). “Some light at the end of the tunnel”: Exploring users’ evidence for the effectiveness of music therapy in adult mental health settings. *Music and Medicine*, 2, 29–40. <https://doi.org/10.1177/1943862109352482>

Azoulay, B., & Orkibi, H. (2015). The four-phase CBN psychodrama model: A manualized approach for practice and research. *Arts in Psychotherapy*, 42, 10–18. <https://doi.org/10.1016/j.aip.2014.12.012>

Baker, F., Jodrey, D., & Intagliata, J. (1992). Social support and quality of life of community support clients. *Community Mental Health Journal*, 28(5), 397–411. <https://doi.org/10.1007/BF00761058>

Bakker, G. M. (2019). A new conception and subsequent taxonomy of clinical psychological problems. *BMC Psychology*, 7(1), 46. <https://doi.org/10.1186/s40359-019-0318-8>

Barlow, D. H., Allen, L. B., & Choate, M. L. (2004). Toward a unified treatment for emotional disorders. *Behavior Therapy*, 35(2), 205–230. [https://doi.org/10.1016/S0005-7894\(04\)80036-4](https://doi.org/10.1016/S0005-7894(04)80036-4)

Barrington, K. A. (2005). *Music Therapy: A study in professionalisation* (Doctoral dissertation). Retrieved from etheses.dur.ac.uk/2791/

Bartholomew, L. K., Parcel, G. S., Kok, G., Gottlieb, N. H., & Fernández, M. E. (2011). *Planning health promotion programs: An intervention mapping approach* (3rd ed.). San Francisco, CA: Jossey-Bass.

Beer, L. E. (2016). From embedded to embodied: including music in arts-based music therapy research. *Music Therapy Perspectives*, 34(1), 33–40.

<https://doi.org/10.1093/mtp/miv006>

Behrends, A., Müller, S., & Dziobek, I. (2012). Moving in and out of synchrony: A concept for a new intervention fostering empathy through interactional movement and dance. *Arts in Psychotherapy, 39*(2), 107–116. <https://doi.org/10.1016/j.aip.2012.02.003>

Bein, E., Anderson, T., Strupp, H. H., Henry, W. P., Schacht, T. E., Binder, J. L., & Butler, S. F. (2000). The effects of training in time-limited dynamic psychotherapy: Changes in therapeutic outcome. *Psychotherapy Research, 10*(2), 119–132.
<https://doi.org/10.1080/713663669>

Bergström, B., Ladd, A., Jones, A., Rosso, L., & Michael, P. (2019). A national survey of clinical psychology graduate student attitudes toward psychotherapy treatment manuals: A replication study. *International Journal of Business, Human and Social Sciences, 12*(6), 923–928. <http://doi.org/10.5281/zenodo.3299863>

Beutler, L. E. (2006). It isn't the size, but the fit. *Clinical Psychology: Science and Practice, 9*(4), 434–438. <https://doi.org/10.1093/clipsy.9.4.434>

Bitsika, V., & Sharpley, C. F. (2006). Treating the client rather than the symptoms: Moving beyond manualised treatments in psychotherapy. *Australian Journal of Guidance and Counselling, 16*(2), 159–175. <https://doi.org/10.1375/ajgc.16.2.159>

Bleijenberg, N., de Man-van Ginkel, J. M., Trappenburg, J. C. A., Ettema, R. G. A., Sino, C. G., Heim, N., ... Schuurmans, M. J. (2018). Increasing value and reducing waste by optimizing the development of complex interventions: Enriching the development phase of the Medical Research Council (MRC) Framework. *International Journal of Nursing Studies, 79*, 86–93. <https://doi.org/10.1016/j.ijnurstu.2017.12.001>

- Boddy, C. (2012). The nominal group technique: An aid to brainstorming ideas in research. *Qualitative Market Research: An International Journal*, 15(1), 6–18.
<https://doi.org/10.1108/13522751211191964>
- Booth, J. M. (2005). Music, drawing and narrative: An adaptation of the Bonny method of guided imagery and music. *Journal of the Association for Music and Imagery*, 10, 55–74.
Retrieved from <https://ami-bonnymethod.org/resources/journals>
- Borgdorff, H. (2010). The production of knowledge in artistic research. In M. Biggs & H. Karlsson (Eds.), *The Routledge Companion to Research in the Arts* (pp. 44–63). New York, NY: Routledge. Retrieved from <https://doi.org/10.4324/9780203841327.ch3>
- Borrelli, B. (2011). The assessment, monitoring, and enhancement of treatment fidelity in public health clinical trials. *Journal of Public Health Dentistry*, 71(s1), S52–S63.
<https://doi.org/10.1111/j.1752-7325.2011.00233.x>
- Brabender, V., & Fallon, A. (2018). *Group psychotherapy in inpatient, partial hospital, and residential care settings*. Washington, DC: American Psychological Association.
Retrieved from <https://doi.org/10.1037/0000113-000>
- Bradt, J. (2012). Randomized controlled trials in music therapy: Guidelines for design. *Journal of Music Therapy*, 49(2), 120–149. <https://doi.org/10.1093/jmt/49.2.120>
- Brooke, S. L. (Ed.) (2006). *Creative arts therapies manual: A guide to the history, theoretical approaches, assessment, and work with special populations of art, play, dance, music, drama and poetry therapies*. Springfield, IL: Charles C Thomas.
- Brüne, M., Abdel-Hamid, M., Sonntag, C., Lehmkämpfer, C., & Langdon, R. (2009). Linking social cognition with social interaction: Non-verbal expressivity, social competence and

“mentalising” in patients with schizophrenia spectrum disorders. *Behavioral and Brain Functions*, 5(6). <https://doi.org/10.1186/1744-9081-5-6>

Bryl, K., & Goodill, S. (2019). Development, execution and acceptance of a manualized dance/movement therapy treatment protocol for the clinical trial in the treatment of negative symptoms and psychosocial functioning in schizophrenia. *American Journal of Dance Therapy*, 1–26. <https://doi.org/10.1007/s10465-019-09312-8>

Buck, E. T., & Havsteen-Franklin, D. (2013). Connecting with the image: How art psychotherapy can help to re-establish a sense of epistemic trust. *ATOL: Art Therapy Online*, 4(1). <https://doi.org/10.25602/GOLD.atol.v4i1.310>

Bullis, J. R., Sauer-Zavala, S., Bentley, K. H., Thompson-Hollands, J., Carl, J. R., & Barlow, D. H. (2015). The unified protocol for transdiagnostic treatment of emotional disorders: preliminary exploration of effectiveness for group delivery. *Behavior Modification*, 39(2), 295–321. <https://doi.org/10.1177/0145445514553094>

Burlingame, G. M., Fuhrman, A., & Johnson, J. E. (2001). Cohesion in group psychotherapy. *Psychotherapy*, 38(4), 373–379. <https://doi.org/10.1037/0033-3204.38.4.373>

Butler, S. F., Henry, W. P., & Strupp, H. H. (1995). Measuring adherence in time-limited dynamic psychotherapy. *Psychotherapy: Theory, Research, Practice, Training*, 32(4), 629–638. <https://doi.org/10.1037/0033-3204.32.4.629>

Carroll, K. M., & Nuro, K. F. (2002). One size cannot fit all: A stage model for psychotherapy manual development. *Clinical Psychology: Science and Practice*, 9(4), 396–406. <https://doi.org/10.1093/clipsy/9.4.396>

Casey, B. J., Oliveri, M. E., & Insel, T. (2014). A neurodevelopmental perspective on the

research domain criteria (RDoC) framework. *Biological Psychiatry*, 76(5), 350–353.

<https://doi.org/10.1016/j.biopsych.2014.01.006>

Chorpita, B. F. (2006). Treatment manuals for the real world: Where do we build them?

Clinical Psychology: Science and Practice, 9(4), 431–433.

<https://doi.org/10.1093/clipsy.9.4.431>

Colbert, T. & Bent, C. (Eds.) (2018). *Working across modalities in the arts therapies: Creative collaborations*. Abingdon: Routledge.

Cotterill, S., Knowles, S., Martindale, A. M., Elvey, R., Howard, S., Coupe, N., ... Spence, M.

(2018). Getting messier with TIDieR: Embracing context and complexity in intervention reporting. *BMC Medical Research Methodology*, 18(1).

<https://doi.org/10.1186/s12874-017-0461-y>

Craig, P., Dieppe, P., Macintyre, S., Michie, S., Nazareth, I., & Petticrew, M. (2006).

Developing and evaluating complex interventions. Swindon: Medical Research Council.

Retrieved from <https://mrc.ukri.org/documents/pdf/complex-interventions-guidance/>

Cramer, A., Tschuschke, V., Koemeda, M., Schulthess, P., & von Wyl, A. (2020). The

therapists' training and their attitudes towards therapy as predictors of therapeutic interventions. *Journal of Contemporary Psychotherapy*, 50(1), 67–76.

<https://doi.org/10.1007/s10879-019-09421-y>

Cuthbert, B. N. (2014). The RDoC framework: Facilitating transition from ICD/DSM to

dimensional approaches that integrate neuroscience and psychopathology. *World*

Psychiatry, 13(1), 28–35. <https://doi.org/10.1002/wps.20087>

Cuthbert, B. N., & Insel, T. R. (2013). Toward the future of psychiatric diagnosis: The seven

pillars of RDoC. *BMC Medicine*, 11(126), 1–8.

<https://doi.org/10.1186/1741-7015-11-126>

Dies, R. R. (1985). Leadership in short-term group therapy: Manipulation or facilitation?

International Journal of Group Psychotherapy, 35(3), 435–455.

<https://doi.org/10.1080/00207284.1985.11491426>

Eyre, L. (2013). Adults in a recovery model setting. In L. Eyre (Ed.), *Guidelines for music therapy practice in mental health* (pp. 115–167). Gilsum, NH: Barcelona.

Feder, E., & Feder, B. (1981). *The expressive arts therapies*. Upper Saddle River, NJ: Prentice Hall.

Fenner, P., Abdelazim, R. S., Bräuninger, I., Strehlow, G., & Seifert, K. (2017). Provision of arts therapies for people with severe mental illness. *Current Opinion in Psychiatry*, 30(4), 306–311. <https://doi.org/10.1016/j.biopsych.2014.01.006>

Folmo, E. J., Karterud, S. W., Bremer, K., Walther, K. L., Kvarstein, E. H., & Pedersen, G. A. F. (2017). The design of the MBT-G adherence and quality scale. *Scandinavian Journal of Psychology*, 58(4), 341–349. <https://doi.org/10.1111/sjop.12375>

Forbat, L., Black, L., & Dulgar, K. (2015). What clinicians think of manualized psychotherapy interventions: Findings from a systematic review. *Journal of Family Therapy*, 37(4), 409–428. <https://doi.org/10.1111/1467-6427.12036>

France, J. & Kramer, S. (Eds.) (2001). *Communication and mental illness. Theoretical and practical approaches*. London, UK: Jessica Kingsley

Geiger, A., Shpigelman, C. N., & Feniger-Schaal, R. (2020). The socio-emotional world of adolescents with intellectual disability: A drama therapy-based participatory action

research. *Arts in Psychotherapy*, 70, 101679.

<https://doi.org/10.1016/j.aip.2020.101679>

Gerber, N., Biffi, E., Biondo, J., Gemignani, M., Hannes, K., & Siegesmund, R. (2020). Arts-based research in the social and health sciences: Pushing for change with an interdisciplinary global arts-based research initiative. *Forum Qualitative Sozialforschung*, 21(2). <https://doi.org/10.17169/fqs-21.2.3496>

Gerge, A., Hawes, J., Eklöf, L., & Pedersen, I. (2019). Proposed mechanisms of change in arts-based psychotherapies. *Voices: A World Forum for Music Therapy*, 19(2), 31.

<https://doi.org/10.15845/voices.v19i2.2564>

Gold, C., Erkkilä, J., & Crawford, M. J. (2012). Shifting effects in randomised controlled trials of complex interventions: A new kind of performance bias? *Acta Psychiatrica Scandinavica*, 125(5), 307–314. <https://doi.org/10.1111/j.1600-0447.2012.01922.x>

Goldberg, R. W., Rollins, A. L., & Lehman, A. F. (2003). Social network correlates among people with psychiatric disabilities. *Psychiatric Rehabilitation Journal*, 26(4), 393–403. <https://doi.apa.org/doi/10.2975/26.2003.393.402>

Grant, S., Mayo-Wilson, E., Montgomery, P., Macdonald, G., Michie, S., Hopewell, S., & Moher, D. (2018). CONSORT-SPI 2018. Explanation and elaboration: Guidance for reporting social and psychological intervention trials. *Trials*, 19, 406. <https://doi.org/10.1186/s13063-018-2735-z>

Grocke, D., Bloch, S., Castle, D., Thompson, G., Newton, R., Stewart, S., ... Gold, C. (2014). Group music therapy for severe mental illness: A randomized embedded- experimental mixed methods study. *Acta Psychiatrica Scandinavica*, 130, 144–153.

<https://doi.org/10.1111/acps.12224>

Hackett, S. S., Ashby, L., Parker, K., Goody, S., & Power, N. (2017). UK art therapy practice-based guidelines for children and adults with learning disabilities. *International Journal of Art Therapy*, 22(2), 84–94. <https://doi.org/10.1080/17454832.2017.1319870>

Hackett, S. S., Taylor, J. L., Freeston, M., Jahoda, A., McColl, E., Pennington, L., & Kaner, E. (2017). Interpersonal art psychotherapy for the treatment of aggression in people with learning disabilities in secure care: A protocol for a randomised controlled feasibility study. *Pilot and Feasibility Studies*, 3, 42. <https://doi.org/10.1186/s40814-017-0186-z>

Haeyen, S., & Heijman, J. (2020). Compassion focused art therapy for people diagnosed with a cluster b/c personality disorder: An intervention mapping study. *Arts in Psychotherapy*, 69. <https://doi.org/10.1016/j.aip.2020.101663>

Haeyen, S., van Hooren, S., Dehue, F., & Hutschemaekers, G. (2018). Development of an art-therapy intervention for patients with personality disorders: an intervention mapping study. *International Journal of Art Therapy: Inscape*, 23(3), 125–135. <https://doi.org/10.1080/17454832.2017.1403458>

Halford, W., & Hayes, R. (1995). Social skills in schizophrenia: Assessing the relationship between social skills, psychopathology and community functioning. *Social Psychiatry and Psychiatric Epidemiology*, 30(1), 14–19. <https://doi-org.ezproxy.library.qmul.ac.uk/10.1007/BF00784429>

Harvey, N., & Holmes, C. A. (2012). Nominal group technique: An effective method for obtaining group consensus. *International Journal of Nursing Practice*, 18(2), 188–194. <https://doi.org/10.1111/j.1440-172X.2012.02017.x>

Havsteen-Franklin, D. (2014). Consensus for using an arts-based response in art therapy.

International Journal of Art Therapy: Inscape, 19(3), 107–113.

<https://doi.org/10.1080/17454832.2014.968598>

Havsteen-Franklin, D., Jovanovic, N., Reed, N., Charles, M., & Lucas, C. (2017). Developing a shared language within arts psychotherapies: A personal construct psychology approach to understanding clinical change. *Arts in Psychotherapy*, 55, 103–110.

<https://doi.org/10.1016/j.aip.2017.05.002>

Health and Care Professions Council. (2013). *Standards of proficiency- Arts Therapists*.

Retrieved from <https://www.hcpc-uk.org/resources/standards/standards-of-proficiency-arts-therapists/>

Henry, W. P., Strupp, H. H., Butler, S. F., Schacht, T. E., & Binder, J. L. (1993). Effects of training in time-limited dynamic psychotherapy: Changes in therapist behavior. *Journal of Consulting and Clinical Psychology*, 61(3), 434–440. <https://doi.org/10.1037/0022-006X.61.3.434>

Herman-Smith, R., Pearson, B., Cordiano, T. S., & Aguirre-McLaughlin, A. M. (2008).

Addressing individual client needs in manualized treatment: Case comparisons. *Clinical Case Studies*, 7(5), 377–396. <https://doi.org/10.1177/1534650108315867>

Herrman, H., Saxena, S., & Moodie, R. (2005). *Promoting mental health: Concepts, emerging evidence, practice*. Geneva: World Health Organisation. Retrieved from

http://whqlibdoc.who.int/publications/2005/9241562943_eng.pdf?ua=1

Heynen, E., Roest, J., Willemars, G., & van Hooren, S. (2017). Therapeutic alliance is a factor of change in arts therapies and psychomotor therapy with adults who have mental

health problems. *Arts in Psychotherapy*, 55, 111–115.

<https://doi.org/10.1016/j.aip.2017.05.006>

Hoffmann, T. C., Glasziou, P. P., Boutron, I., Milne, R., Perera, R., Moher, D., ... Michie, S.

(2014). Better reporting of interventions: Template for intervention description and replication (TIDieR) checklist and guide. *BMJ*, 348. <https://doi.org/10.1136/bmj.g1687>

Hogan, S. (2001). *Healing arts: The history of art therapy*. London: Jessica Kingsley.

Huet, V. (1997). Challenging professional confidence in arts therapies and psychiatric

rehabilitation. *Inscape*, 2(1), 14–19. <https://doi.org/10.1080/17454839708413040>

Jones, D. P. (2020). *The arts therapies: A revolution in healthcare* (2nd Ed.). London, UK:

Routledge. <https://doi.org/10.4324/9780203308066>

Jones, P. (2007). *Drama as therapy: Theory, practice and research* (2nd ed.). London:

Routledge.

Karkou, V., & Sanderson, P. (2006). *The arts therapies: A research based map of the field*.

Edinburgh: Elsevier. <https://doi.org/10.1016/B978-0-443-07256-7.X5001-3>

Karterud, S. (2016). *Mentalization-based group therapy (MBT-G): A theoretical, clinical, and*

research manual. Oxford, UK: Oxford University Press.

<https://doi.org/10.1093/med:psych/9780198753742.001.0001>

Kerns, J. (2007). Verbal communication impairments and cognitive control components in

people with schizophrenia. *Journal of Abnormal Psychology*, 116(2), 279–289.

<https://doi.org/10.1037/0021-843X.116.2.279>

Khakpoor, S., Bytamar, J. M., & Saed, O. (2019). Reductions in transdiagnostic factors as the

potential mechanisms of change in treatment outcomes in the Unified Protocol: A randomized clinical trial. *Research in Psychotherapy: Psychopathology, Process and Outcome*, 22(3), 402–412. <https://doi.org/10.4081/ripppo.2019.379>

Koch, S. C. (2017). Arts and health: Active factors and a theory framework of embodied aesthetics. *Arts in Psychotherapy*, 54, 85–91. <https://doi.org/10.1016/j.aip.2017.02.002>

Knill, P. J. (1994). Multiplicity as a tradition: Theories for interdisciplinary arts therapies- an overview. *Arts in Psychotherapy*, 21(5), 319–328. [https://doi.org/10.1016/0197-4556\(94\)90059-0](https://doi.org/10.1016/0197-4556(94)90059-0)

Kraemer, H. C., Wilson, T., Fairburn, C. G., & Agras, S. (2002). Mediators and moderators of treatment effects in randomized clinical trials. *Archives of General Psychiatry*, 59(10), 877–883. <https://doi.org/10.1001/archpsyc.59.10.877>

Landy, R. J. (1995). Isolation and collaboration in the creative arts therapies- the implications of crossing borders. *Arts in Psychotherapy*, 22(2), 83–86. [https://doi.org/10.1016/0197-4556\(95\)00010-3](https://doi.org/10.1016/0197-4556(95)00010-3)

Laposa, J. M., Mancuso, E., Abraham, G., & Loli-Dano, L. (2017). Unified protocol transdiagnostic treatment in group format: A preliminary investigation with anxious individuals. *Behavior Modification*, 41(2), 253–268. <https://doi.org/10.1177/0145445516667664>

Leamy, M., Bird, V., Le Boutillier, C., Williams, J., & Slade, M. (2011). Conceptual framework for personal recovery in mental health: Systematic review and narrative synthesis. In *British Journal of Psychiatry*, 199(6), 445–452. <https://doi.org/10.1192/bjp.bp.110.083733>

Ledger, A., & McCaffrey, T. (2015). Performative, arts-based, or arts-informed? Reflections on the development of arts-based research in music therapy. *Journal of Music Therapy*, 52(4), 441–456. <https://doi.org/10.1093/JMT/THV013>

Lee, Y., & Vakoch, D. A. (1996). Transfer and retention of implicit and explicit learning. *British Journal of Psychology*, 87(4), 637–651. <https://doi.org/10.1111/j.2044-8295.1996.tb02613.x>

Lyons, S. (2019). *Arts therapies for dementia: A systematic review and community-based case study on the value of music therapy and dance movement therapy* (Unpublished doctoral dissertation). Liverpool, UK: Edge Hill University.

Luborksy, L., & DeRubeis, R. (1984). The use of psychotherapy treatment manuals: A small revolution in psychotherapy research style. *Clinical Psychology Review*, 4(1), 5–14. [https://doi.org/10.1016/0272-7358\(84\)90034-5](https://doi.org/10.1016/0272-7358(84)90034-5)

Mac Nair-Semands, R. R., Ogradniczuk, J. S., & Joyce, A. S. (2010). Structure and initial validation of a short form of the therapeutic factors inventory. *International Journal of Group Psychotherapy*, 60(2), 245–281. <https://doi.org/10.1521/ijgp.2010.60.2.245>

MacKenzie, K. R. (1990). *Introduction to time-limited group psychotherapy*. Washington, DC: American Psychiatric Press.

Malchiodi, C. A. (Ed.) (2005). *Expressive therapies*. New York, NY: Guilford Press.

Malloch, S., & Trevarthen, C. (2009). Musicality: Communicating the vitality and interests of life. In S. Malloch & C. Trevarthen (Eds.), *Communicative musicality: Exploring the basis of human companionship* (pp. 1–11). Oxford, UK: Oxford University Press.

McAleer, J. T., Bowler, J. L., Bowler, M. C., & Schoemann, A. M. (2020). Implicit and explicit

creativity: Further evidence of the integrative model. *Personality and Individual Differences*, 154, 109643. <https://doi.org/10.1016/j.paid.2019.109643>

McCaffrey, T. (2016). Music therapy in mental health care for adults. In J. Edwards (Ed.) *The Oxford handbook of music therapy* (pp.249–267). Oxford, UK: Oxford University Press. <https://doi.org/10.1093/oxfordhb/9780199639755.013.29>

McCaffrey, T., Carr, C., Solli, H. P., & Hense, C. (2018). Music therapy and recovery in mental health: seeking a way forward. *Voices: A World Forum for Music Therapy*, 18(1). <https://doi.org/10.15845/voices.v18i1.918>

McCaffrey, T., & Edwards, J. (2015). Meeting art with art: Arts-based methods enhance researcher reflexivity in research with mental health service users. *Journal of Music Therapy*, 52(4), 515–532. <https://doi.org/10.1093/JMT/THV016>

McCaffrey, T., Edwards, J., & Fannon, D. (2011). Is there a role for music therapy in the recovery approach in mental health? *Arts in Psychotherapy*, 38(3), 185–189. <https://doi.org/10.1016/j.aip.2011.04.006>

McCaffrey, T., Higgins, P., Monahan, C., Moloney, S., Nelligan, S., Clancy, A., & Cheung, P. S. (2020). Exploring the role and impact of group songwriting with multiple stakeholders in recovery-oriented mental health services. *Nordic Journal of Music Therapy*, 1–20. <https://doi.org/10.1080/08098131.2020.1771755>

McNiff, S. (1981). *The Arts and Psychotherapy*. Springfield, IL: Charles C Thomas.

Meltzer, C. (2020). The reflexive practitioner. Using arts-based methods and research for professional development. In L. Mackay, G. Barton, S. Garvis, & V. Sappa (Eds.), *Arts-based research: Resilience and well-being across the lifespan* (pp. 209–231). Cham:

Palgrave Macmillan. https://doi.org/10.1007/978-3-030-26053-8_12

Metzner, S., Jaeger, U., Masuhr, O., Olschewski, U., Gräfe, E., Böske, A. C., & Dämpelmann, M. (2018). Forms of attunement during the initial stages of music therapy for patients with acute psychosis - a multicentre clinical study. *Nordic Journal of Music Therapy*, 27(5), 360–380. <https://doi.org/10.1080/08098131.2018.1478879>

Montgomery, C. (2002). Role of dynamic group therapy in psychiatry. *Advances in Psychiatric Treatment*, 8(1), 34–41. [https://doi.org/DOI: 10.1192/apt.8.1.34](https://doi.org/DOI:10.1192/apt.8.1.34)

Moula, Z., Karkou, V., & Powell, J. (2019). A pilot cross-over randomised controlled trial of child-focused process and outcome evaluation of arts therapies at primary mainstream schools: study protocol. *Arts and Health*.
<https://doi.org/10.1080/17533015.2019.1703198>

Najavits, L. M. (1997). Psychotherapists' implicit theories of therapy. *Journal of Psychotherapy Integration*, 7(1), 1–16. <https://doi.org/10.1037/h0101137>

National Institute of Clinical Excellence. (2014). *Psychosis and schizophrenia in adults: treatment and management. Guidance and guidelines*. National Institute of Clinical Excellence. Retrieved from <https://www.nice.org.uk/guidance/cg178>

Nelson, M. C., Cordray, D. S., Hulleman, C. S., Darrow, C. L., & Sommer, E. C. (2012). A procedure for assessing intervention fidelity in experiments testing educational and behavioral interventions. *Journal of Behavioral Health Services and Research*, 39(4), 374–396. <https://doi.org/10.1007/s11414-012-9295-x>

Nemetz, L. D. (2006). Moving with meaning: The historical progression of dance/movement therapy. In S. L. Brooke (Ed.), *Creative arts therapies manual: A guide to the history*,

theoretical approaches, assessment, and work with special populations of art, play, dance, music, drama and poetry therapies (pp. 95–108). Springfield, IL: Charles C Thomas.

O’Cathain, A., Croot, L., Duncan, E., Rousseau, N., Sworn, K., Turner, K. M., ... Hoddinott, P. (2019). Guidance on how to develop complex interventions to improve health and healthcare. *BMJ Open*, *9*(8). <https://doi.org/10.1136/bmjopen-2019-029954>

Odell-Miller, H. (2013). The development of clinical music therapy in adult mental health practice: Music, health and therapy. Case study. In V. Bates, A. Bleakley, & S. Goodman (Eds.), *Medicine health and the arts: Approaches to the medical humanities* (pp. 264–280). London, UK: Routledge. <https://doi.org/10.4324/9780203079614>

Orfanos, S., Banks, C., & Priebe, S. (2015). Are group psychotherapeutic treatments effective for patients with schizophrenia? A systematic review and meta-analysis. *Psychotherapy and Psychosomatics*, *84*(4), 241–249. <https://doi.org/10.1159/000377705>

Orfanos, S., & Priebe, S. (2017). Group therapies for schizophrenia: initial group climate predicts changes in negative symptoms. *Psychosis*, *9*(3), 225–234. <https://doi.org/10.1080/17522439.2017.1311360>

Parkinson, S., & Whiter, C. (2016). Exploring art therapy group practice in early intervention psychosis. *International Journal of Art Therapy: Inscape*, *21*(3), 116–127. <https://doi.org/10.1080/17454832.2016.1175492>

Parsons, A., Omylinska-Thurston, J., Karkou, V., Harlow, J., Haslam, S., Hobson, J., ... Griffin, J. (2020). Arts for the blues – a new creative psychological therapy for depression. *British*

Journal of Guidance & Counselling, 48(1), 5–20.

<https://doi.org/10.1080/03069885.2019.1633459>

Perlini, C., Marini, A., Garzitto, M., Isola, M., Cerruti, S., Marinelli, V., ... Brambilla, P. (2012).

Linguistic production and syntactic comprehension in schizophrenia and bipolar disorder. *Acta Psychiatrica Scandinavica*, 126(5), 363–376.

<https://doi.org/10.1111/j.1600-0447.2012.01864.x>

Priebe, S., Arenas Borrero, Á., Bird, V., Džubur Kulenović, A., Giacco, D., Gómez Restrepo, C.,

... Uribe Restrepo, J. M. (2019). Possibilities for the future of global mental health: A scenario planning approach. *BMC Psychiatry*, 19(1). <https://doi.org/10.1186/s12888-019-2381-3>

Priebe, S., Omer, S., Giacco, D., & Slade, M. (2014). Resource-oriented therapeutic models in psychiatry: Conceptual review. *British Journal of Psychiatry*, 204(4), 256–261.

<https://doi.org/10.1192/bjp.bp.113.135038>

Purdie, H. (1993). Music therapy and its relationship with creative arts therapies. *British Journal of Music Therapy*, 7(2), 23–25.

<https://doi.org/10.1177%2F135945759300700204>

Reber, A. S. (2001). Tacit knowledge, psychology of. In N. J. Smelser, & P. B. Baltes (Eds.), *International Encyclopedia of the Social & Behavioral Sciences* (pp.15431–15435).

Oxford: Elsevier.

<https://doi.org/10.1016/B0-08-043076-7/01492-3>

Robb, S. L., Burns, D. S., Docherty, S. L., & Haase, J. E. (2011). Ensuring treatment fidelity in a multi-site behavioral intervention study: Implementing NIH behavior change

consortium recommendations in the SMART trial. *Psycho-Oncology*, 20(11), 1193–1201. <https://doi.org/10.1002/pon.1845>

Rogers, C. R. (2011). *On Becoming a Person. A therapist's view of psychotherapy*. London: Constable & Robinson. (Original work published in 1967)

Rolvsjord, R. (2010). *Resource-oriented music therapy in mental health care*. Gilsum, NH: Barcelona. Retrieved from <https://www.barcelonapublishers.com/resource-oriented-music-therapy-mental-health-care>

Rolvsjord, R. (2015). What clients do to make music therapy work: A qualitative multiple case study in adult mental health care. *Nordic Journal of Music Therapy*, 24(4), 296–321. <https://doi.org/10.1080/08098131.2014.964753>

Rolvsjord, R., Gold, C., & Stige, B. (2005). Research rigour and therapeutic flexibility: Rationale for a therapy manual developed for a randomised controlled trial. *Nordic Journal of Music Therapy*, 14(1), 15–32. <https://doi.org/10.1080/08098130509478122>

Rousseau, N., Turner, K. M., Duncan, E., O’Cathain, A., Croot, L., Yardley, L., & Hoddinott, P. (2019). Attending to design when developing complex health interventions: A qualitative interview study with intervention developers and associated stakeholders. *PLoS ONE*, 14(10). <https://doi.org/10.1371/journal.pone.0223615>

Sajjani, N., Marxen, E. & Zarate, R. (2017). Critical perspectives in the arts therapies: Response/ability across a continuum of practice. *Arts in Psychotherapy*, 54, 28–37. <https://doi.org/10.1016/j.aip.2017.01.007>

Sajjani, N., Mayor, C., & Tillberg-Webb, H. (2020). Aesthetic presence: The role of the arts in the education of creative arts therapists in the classroom and online. *Arts in*

Psychotherapy, 69, 101668. <https://doi.org/10.1016/j.aip.2020.101668>

Samaritter, R. (2018). The aesthetic turn in mental health: reflections on an explorative study into practices in the arts therapies. *Behavioral Sciences*, 8(4), 41.

<https://doi.org/10.3390/bs8040041>

Sánchez, J., Sung, C., Phillips, B. N., Tschopp, M. K., Muller, V., Lee, H. –L., & Chan, F. (2019).

Predictors of perceived social effectiveness of individuals with serious mental illness.

Psychiatric Rehabilitation Journal, 42(1), 88–99.

<https://doi.apa.org/doi/10.1037/prj0000321>

Sanislow, C. (2016). Connecting psychopathology meta-structure and mechanisms. *Journal of Abnormal Psychology*, 125(8), 1158–1165.

<https://content.apa.org/doi/10.1037/abn0000207>

Sholomskas, D. E., Syracuse-Siewert, G., Rounsaville, B. J., Ball, S. A., Nuro, K. F., & Carroll, K.

M. (2005). We don't train in vain: A dissemination trial of three strategies of training clinicians in cognitive-behavioral therapy. *Journal of Consulting and Clinical Psychology*,

73(1), 106–115. <https://doi.org/10.1037/0022-006X.73.1.106>

Sidani, S. & Braden, C. J. (2011). *Design, evaluation and translation of nursing interventions*.

Chichester, UK: John Wiley & Sons.

Solli, H. P., Rolvsjord, R., & Borg, M. (2013). Toward understanding music therapy as a

recovery-oriented practice within mental health care: A meta-synthesis of service users' experiences. *Journal of Music Therapy*, 50(4), 244–273.

<https://doi.org/10.1093/jmt/50.4.244>

Springborg, C. (2020). Designing dance into qualitative research. In J. Ward, & H. Shortt

(Eds.), *Using arts-based research methods* (pp. 41–74). Cham: Palgrave Macmillan.

https://doi.org/10.1007/978-3-030-33069-9_3

Springham, N., Dunne, K., Noyse, S., & Swearingen, K. (2012). Art therapy for personality disorder: 2012 UK professional consensus guidelines, development process and outcome. *International Journal of Art Therapy, 17*(3), 130–134.

<https://doi.org/10.1080/17454832.2012.734834>

Springham, N., & Huet, V. (2018). Art as relational encounter: An ostensive communication theory of art therapy. *Art Therapy, 35*(1), 4–10.

<https://doi.org/10.1080/07421656.2018.1460103>

Stern, D. N. (2010). *Forms of vitality: Exploring dynamic experience in psychology, the arts, psychotherapy and development*. Oxford, UK: Oxford University Press.

Story, K. M., & Beck, B. D. (2017). Guided imagery and music with female military veterans: An intervention development study. *Arts in Psychotherapy, 55*, 93–102.

<https://doi.org/10.1016/j.aip.2017.05.003>

Taylor Buck, E., & Dent-brown, K. (2014). Creative manuals. *International Journal of Art Therapy, 19*(2), 82–87. <https://doi.org/10.1080/17454832.2014.906475>

Taylor Buck, E., Dent-Brown, K., Parry, G., & Boote, J. (2014). Dyadic art psychotherapy: Key principles, practices and competences. *Arts in Psychotherapy, 41*(2), 163–173.

<https://doi.org/10.1016/j.aip.2014.01.004>

Taylor Buck, E., & Hendry, A. (2016). Developing principles of best practice for art therapists working with children and families. *International Journal of Art Therapy, 21*(2), 56–65.

<https://doi.org/10.1080/17454832.2016.1170056>

Thornicroft, G., Mehta, N., Clement, S., Evans-Lacko, S., Doherty, M., Rose, D., ... Henderson, C. (2016). Evidence for effective interventions to reduce mental-health-related stigma and discrimination. *Lancet*, *387*(10023), 12–18.

[https://doi.org/10.1016/S0140-6736\(15\)00298-6](https://doi.org/10.1016/S0140-6736(15)00298-6)

Trompenaars, F. J., Masthoff, E. D., Van Heck, G. L., de Vries, J., & Hodiament, P. P. (2007). Relationships between social functioning and quality of life in a population of Dutch adult psychiatric outpatients. *International Journal of Social Psychiatry*, *53*(1), 36–47.

<https://doi.org/10.1177/0020764006074281>

Truijens, F., Zühlke-van Hulzen, L., & Vanheule, S. (2019). To manualize, or not to manualize: Is that still the question? A systematic review of empirical evidence for manual superiority in psychological treatment. *Journal of Clinical Psychology*, *75*(3), 329–343.

<https://doi.org/10.1002/jclp.22712>

Tschuschke, V., & Dies, R. (1994). Intensive analysis of therapeutic factors and outcome in long-term inpatient groups. *International Journal of Group Psychotherapy*, *44*(2), 185–208. <https://doi.org/10.1080/00207284.1994.11490742>

Tschuschke, V., Cramer, A., Koehler, M., Berglar, J., Muth, K., Staczan, P., ... Koemeda-Lutz, M. (2015). The role of therapists' treatment adherence, professional experience, therapeutic alliance, and clients' severity of psychological problems: Prediction of treatment outcome in eight different psychotherapy approaches. Preliminary results of a naturalistic study. *Psychotherapy Research*, *25*(4), 420–434.

<https://doi.org/10.1080/10503307.2014.896055>

Turner, K. M., Rousseau, N., Croot, L., Duncan, E., Yardley, L., O'Cathain, A., & Hodinott, P.

(2019). Understanding successful development of complex health and healthcare interventions and its drivers from the perspective of developers and wider stakeholders: An international qualitative interview study. *BMJ Open*, 9(5).
<https://doi.org/10.1136/bmjopen-2018-028756>

UKCP. (2012). *UKCP standards of education and training (2012) - the minimum core criteria psychotherapy with adults*.

Retrieved from <http://www.psychotherapy.org.uk/index.php?id=428>

Vakoch, D. A., & Strupp, H. H. (2000). The evolution of psychotherapy training: Reflections on manual-based learning and future alternatives. *Journal of Clinical Psychology*, 56(3), 309–318.

[https://doi.org/10.1002/\(SICI\)1097-4679\(200003\)56:3<309::AID-JCLP7>3.0.CO;2-7](https://doi.org/10.1002/(SICI)1097-4679(200003)56:3<309::AID-JCLP7>3.0.CO;2-7)

Waller, D. (1991). *Becoming a profession: The history of art therapy in Britain 1940-1982 (Psychology revivals)*. London: Routledge. <https://doi.org/10.4324/9780203750209>

Waltz, J., Addis, M. E., Koerner, K., & Jacobson, N. S. (1993). Testing the integrity of a psychotherapy protocol: Assessment of adherence and competence. *Journal of Consulting and Clinical Psychology*, 61(4), 620–630. <https://doi.org/10.1037/0022-006X.61.4.620>

Wampold, B. E. (2015). How important are the common factors in psychotherapy? An update. *World Psychiatry*, 14(3), 270–277. <https://doi.org/10.1002/wps.20238>

Wampold, B. E., & Imel, Z. E. (2015). *The great psychotherapy debate: The evidence for what makes psychotherapy work* (2nd ed.). New York, NY: Routledge.

<https://doi.org/10.4324/9780203582015>

- Westen, D. (2002). Manualizing manual development. *Clinical Psychology: Science and Practice*, 9(4), 416–418. <https://doi.org/10.1093/clipsy.9.4.416>
- Windle, E., Tee, H., Sabitova, A., Jovanovic, N., Priebe, S., & Carr, C. (2020). Association of patient treatment preference with dropout and clinical outcomes in adult psychosocial mental health interventions: a systematic review and meta-analysis. *JAMA Psychiatry*, 77(3), 294–302. <https://doi.org/10.1001/jamapsychiatry.2019.3750>
- World Health Organization (2001). *Mental Health: New understanding, new hope*. Geneva: World Health Organization.
Retrieved from https://www.who.int/whr/2001/en/whr01_en.pdf?ua=1
- Wright, T., & Holttum, S. (2020). *BAAT Guidelines on Art Therapy for People with a Psychosis-Related Diagnosis*. Retrieved from [https://www.baat.org/Assets/Docs/General/BAAT Guidelines AT Psychosis PART 1-2-3.pdf](https://www.baat.org/Assets/Docs/General/BAAT%20Guidelines%20AT%20Psychosis%20PART%201-2-3.pdf)
- Yalom, I. D., & Leszcz M. (2005). *The theory and practice of group psychotherapy* (5th ed.). New York, NY: Basic Books.
- Zubala, A. (2013). *Description and evaluation of arts therapies practice with adults suffering from depression in the UK* (Doctoral dissertation). Retrieved from <https://eresearch.qmu.ac.uk/handle/20.500.12289/7400>
- Zubala, A., MacIntyre, D. J., Gleeson, N., & Karkou, V. (2013). Description of arts therapies practice with adults suffering from depression in the UK: Quantitative results from the nationwide survey. *Arts in Psychotherapy*, 40(5), 458–464.
<https://doi.org/10.1016/j.aip.2013.09.003>

Zubala, A., MacIntyre, D. J., Gleeson, N., & Karkou, V. (2014). Description of arts therapies practice with adults suffering from depression in the UK: Qualitative findings from the nationwide survey. *Arts in Psychotherapy, 41*(5), 535–544.

<https://doi.org/10.1016/j.aip.2014.10.005>

Table 1

Development of Practice Principles Through Initial Meetings and Therapist Workshops

Initial meetings		Workshops	Final grouping
Practice principles	Proscribed principles		
<p>Non-judgemental, validating attitude</p> <p>Empathy, unconditional positive regard, congruence of therapist, modelling for the group</p>	<p>Over-sharing.</p> <p>Therapist as a ‘blank slate’.</p>	<p>Use of therapist self-disclosure as a means of building trust and sense of being understood</p>	<p>1. Engagement, interest and warmth</p>
<p>Modelling curiosity and interest in clients’ experience of the modality (as opposed to interpreting)</p>	<p>Direct interpretation without checking client’s understanding of events or ignoring client’s communicated understanding of events</p>	<p>Being with what is.</p> <p>Cultivating curiosity.</p>	<p>2. Exploration, curiosity, not-knowing stance</p>
<p>Use of the art form and stance to meet the person/group where they are at.</p>	<p>Not intervening when expression is uncontained</p>	<p>Letting go of tension.</p>	<p>3. Active and dynamic attunement</p>
<p>Supporting self-regulation through containment and transformation of expression</p>		<p>Safe space for feelings to be expressed.</p>	<p>4. Arousal regulation</p>
<p>The therapist adapts their stance and involvement with the group depending on whether the group is in early, middle or late stages.</p>		<p>Therapist stance is more active in early stages. Therapist becomes less directive but is ready to resume directions if the group requires.</p>	<p>5. Developmentally informed group leadership</p>

Initial meetings		Workshops	Final grouping
Practice principles	Proscribed principles		
Importance of supportive and effective co-working.			6. Cooperation between therapists
Modelling reliability and trustworthiness through actions	Rigid adherence to boundaries without collaborating or seeking mutual understanding with clients	Promoting safety. Elastic/flexible boundaries to be negotiated as a group. Fluidity. Boundaries as a group container and structure. Therapist responsibility.	7. Managing group boundaries
Early stages focus on building trust, safety and cohesion. Later stages, group autonomy is promoted, with the therapist becoming more directive to prepare for group ending.	Not intervening/ being completely non-directive in the group Therapist directs- does not give group opportunity to lead or direct	Make changes led by client.	8. Regulating group phases
Open and equal relationship to model safety and ways of relating in the group.	Once cohesion and trust are established, not challenging group to try out new things or take risks	Group cohesion. Commonality of experience. Arts promote a sense of belonging – community and friendship.	9. Active promotion of group cohesion
Encouraging sharing and helping between group members	Contributions ignored or dismissed by therapist		10. Caring for the group and each member
Encouraging active participation, exploration and use of the art form	Not assisting or intervening when the client is unable to use the art form	Creativity. Enabling participation. Potential space, possibilities.	11. Modelling use and potentials of art-form

	Initial meetings		Workshops	Final grouping
	Practice principles	Proscribed principles		
Encourage self-expression and creativity		Allowing the group to avoid or not use the art form	Sharing of different cultures Play, Enabling participation.	12. Encouraging engagement and exploration of the art-form
Encourage exploration and new ways of using the art form.		Persisting in an activity which is not appealing or helpful to the group Allowing long periods of talking or silence without any use of the art form.	Not dogmatic but coherent. Practice evolves with group life.	13. Developing and expanding use of the art-form
Highlighting group interactions-encouraging clients to make their own meaning from this and try out new ways of relating		Problem solving for the group	Noticing patterns → changing patterns. Acknowledging trauma and working through. Exploration of differences.	14. Focus on personal and group issues
Emotional involvement in the creative process		Therapist over-involvement in emotional processes	Working with what is within Experiencing self differently.	15. Focus on implicit experiences (arts-based, embodied, emotional)
Relating all interactions in the sessions back to the here and now Develop relationships that are shared, equal, collaborative, respectful, mutually trusting Use of transference to understanding evolving relationships within the group		Focus upon past events without connection to here and now of the group	Making connections Breaking isolation Social contact Peer support Practising interactions Shared experiences Communication. Acknowledging diversity and difference.	16. Focus on relationship between therapists and patients
Help service users to discover personal meaning and understanding from the arts process			Therapist is dynamic and moves fluidly between verbal and nonverbal ways of relating	17. Encouraging integration of arts, implicit and relational experiences in the group

Initial meetings		Workshops	Final grouping
Practice principles	Proscribed principles		
		<p>A place where emotions are ok.</p> <p>Integration</p> <p>Connecting with different parts of self, integrating using creative means.</p> <p>Other ways of seeing self/others.</p>	
<p>Support service users to build on useful and helpful experiences of using the art-form within the group by identifying ways this can be continued in their day-to-day life.</p>		<p>Use of arts in own life outside of therapy for resilience/enjoyment.</p>	<p>18. Integrating use of art-form inside and outside of sessions</p>
<p>Offering encouragement and facilitating/fostering positive experiences</p>	<p>Focus only on deficits or problems</p> <p>Avoiding problems</p>	<p>Building confidence and self-esteem.</p> <p>Resilience building.</p>	<p>19. Acknowledging and encouraging strengths, preferences and positive experiences</p>
<p>Collaborating with group members regarding goals and methods of working</p>			
<p>Focusing upon group members' strengths and potentials, recognising clients' competence in the creative process and unique creative identity</p>			

Table 2*Clinical Scenarios Where Therapists Make an Observable Therapeutic Action*

Item	Definition
1 Unmanageable affect	High / low affect that impacts upon ability to relate to or be with the art form/others and is difficult to self-regulate
2 Power dynamics	Interpersonal dynamics, criticism, scapegoating, blame, dominance, boundaries, them and us
3 Collaboration	Empathy, attunement, relating, working with others, co-production of the art form
4 Ambivalence about using arts media	Difficulty using art form due eg. to lack of experience, current feeling state, self-consciousness in front of others
5 Using the arts for retreat/distraction/avoidance	Avoiding contact with others by focusing solely on the arts medium, avoiding connection to feelings
6 Identity	Artistic biography, narratives, cultural associations, rites of passage, rituals
7 Linking arts process with personal relationships and community	Making connections between use of the art form and relationships within/outside the group.
8 Improvisation	Authentic, new artistic element, expression, play, creativity in use of art form in the moment
9 Incongruence between art form and feelings	Expression within the art-form does not match the presentation or stated affect of the group/group member
10 Concrete, rigid, stuck	Set ways of using the art-form, lack of flexibility, lack of connection to feelings and interpersonal process.
11 Sub-groups	Pairing, trios, quartets
12 Diverse uses of the arts forms	New ways of using the arts media eg. playing a chair with drumsticks, finding a different movement, different styles and genres of the art form introduced.
13 Over-compliance / need to 'fit in'	Taking a background role in the group, using the arts to hide from/avoid challenge by others. Working to please the therapist.
14 Pre-occupation with an 'aesthetic' or competence	Valuing the artistic product over and above the arts/group process. Focusing on ability/competence to use the arts without relating back to self and situation.

Table 3

Therapist Principles for Practice

Therapist principles for practice	
1. Engagement, interest and warmth	The therapist conveys empathy, authenticity, positive regard, offers encouragement and is emotionally involved both during and outside the creative process.
2. Exploration, curiosity, not-knowing stance	Overall stance of the therapist in terms of active listening, modelling curiosity and interest and conveying a stance of 'not-knowing'.
3. Active and dynamic attunement	The therapist actively meets and attunes to the presenting internal state of each individual and the group as a whole through use of the art-form, body language and words.
4. Arousal regulation	The therapist uses attunement and interactions within relationships, primarily with the art-form, to help clients to regulate arousal levels, paying attention to under/over-arousal to establish a calm and reflective holding presence. The therapist later supports clients to carefully expand upon and return to different arousal states. The therapist only challenges the client/group once safety and cohesion are established.
5. Developmentally informed group leadership	The therapist's style of group leadership changes depending on the developmental level of the group with greater activity at the beginning and end of the group process.
6. Cooperation between therapists	The therapist and co-facilitator work together to establish a safe space, model relational capacity and cooperation through confident, transparent and open communication.
7. Managing group boundaries	Active and dynamic management of the physical space and group boundaries. Supporting and encouraging attendance, maintaining contact, modelling reliability and trustworthiness through actions.
8. Regulating group phases	The therapist is active in helping the group establish and move between beginning, middle and end stages of the session and process as a whole.
9. Active promotion of group cohesion	The therapist works actively to promote a sense of group wholeness and belonging through arts-based activities and discussion especially in early stages of the group.
10. Caring for the group and each member	The therapist provides a calm, reflective and holding presence; promotes shared, equal, collaborative and mutually trusting relationships. Actively addresses destructive behaviour or threats towards the group.
11. Modelling use and potentials of art-form	The therapist demonstrates ways to begin, and use the art-form.
12. Encouraging engagement and exploration of the art-form	The therapist supports members to overcome anxiety in use of arts materials and develop curiosity as to what is possible.
13. Developing and expanding use of the art-form	

The therapist supports and begins to challenge members to play creatively to develop and expand their use of the art-form.

14. Focus on personal and group issues

The therapist supports group members to actively explore personal and group issues raised in the here and now of the group.

15. Focus on implicit experiences (arts-based, embodied, emotional)

The therapist draws attention to the embodied or felt qualities of using the art-form through arts-based or verbal responses.

16. Focus on relationship between therapists and patients

The therapist seeks to develop shared, equal, mutually trusting relationships, highlights group interactions and encourages clients to make their own meaning/try out new ways of relating; encourages clients to learn from one another.

17. Encouraging integration of arts, implicit and relational experiences in the group

The therapist encourages connection of bodily and arising emotional/felt states to the process of engaging with the art form and helps to make connections from the use of the arts to relationships within the group.

18. Integrating use of art-form inside and outside of sessions

The therapist promotes and explores connections between arts, group experiences and everyday life.

19. Acknowledging and encouraging strengths, preferences and positive experiences

The therapist nurtures positive experiences through collaboration and acknowledgement of the group's existing strengths and preferences.
