

CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

**The Coronavirus Anxiety Scale: Cross-National Measurement Invariance and Convergent  
Validity Evidence**

# CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

## Abstract

Coronavirus Anxiety Scale (CAS) is a widely used measure that captures somatic symptoms of coronavirus-related anxiety. In a large-scale collaboration spanning 60 countries with 21,513 respondents, we examined CAS's measurement invariance and the evidence of convergent validity of CAS scores in relation to the fear of COVID-19 (FCV-19S) and the satisfaction with life (SWLS-3) scales. We utilized both conventional exact invariance tests and alignment procedures, revealing that the single-factor model fit the data well in almost all countries. Partial scalar invariance was supported in a subset of 56 countries. To ensure the robustness of results, given the unbalanced samples, we employed resampling techniques both with and without replacement and found the results were more stable in larger samples. The alignment procedure demonstrated a high degree of measurement invariance with 9% of the parameters exhibiting non-invariance. We also conducted simulations of alignment using the parameters estimated in the current model. This demonstrated the reliability of the means but indicated challenges in estimating the latent variances. Positive and strong correlations between CAS and FCV-19S estimated with three different approaches were found in most countries. Correlations of CAS and SWLS-3 were weak and negative, but significantly differed from zero in several countries. Overall, the study provided support for the measurement invariance of the CAS and offered evidence of its convergent validity while also highlighting issues with variance estimation.

*Keywords:* coronavirus anxiety, measurement invariance, alignment, validity, culture.

**Public significance statement.** We found that the widely used Coronavirus Anxiety Scale is generally suitable for cross-national research. Given that previously its comparability was questioned, the current study provided evidence of invariance making international comparisons of coronavirus-related anxiety possible.

**The Coronavirus Anxiety Scale: Cross-National Measurement Invariance and Convergent Validity Evidence**

The SARs-CoV-2 (otherwise known as coronavirus or COVID-19) pandemic has had an unprecedented impact on the psychological functioning of people worldwide (Aknin et al., 2022). Notably, fear and anxiety have been common responses to the ongoing pandemic (Luo et al., 2021), due to its novelty, uncertainty, and unpredictability, and the scope of impact and dramatic consequences of the COVID-19 crisis. Terms “fear of COVID-19” (Ahorsu et al., 2022) and “coronaphobia” (Asmundson & Taylor, 2020) have been used to describe intense fear and discomfort experienced when a person is exposed to COVID-19 information or when they are thinking about the COVID-19 disease. The urgency of understanding mental health aspects of the COVID-19 outbreak has led to the development of a number of questionnaires aimed at assessing fear, anxiety, and stress related to a novel coronavirus (for a review, please see Voitsidis et al., 2021). In the present paper, we focus on a five-item Coronavirus Anxiety Scale (CAS; Lee, 2020a), one of the most popular mental health questionnaires to have emerged during the pandemic (e.g., based on Scopus the CAS received more than 100 citations in 2020, and close to 300 during 2021). We evaluate the cross-national measurement invariance of the CAS using samples from 60 countries and examine the convergent validity evidence of the CAS scores by investigating its relations to fear of COVID-19 and life satisfaction scores.

**The Coronavirus Anxiety Scale**

The CAS is an instrument designed at the onset of the pandemic (in March 2020) to assess coronavirus related anxiety (i.e., dysfunctional anxiety in the context of the ongoing pandemic). It was developed to serve as both a screening instrument and a survey measure, helping health practitioners and researchers alike (Lee, 2020a). Item content shows that the CAS

## CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

captures exclusively physical and physiological (i.e., somatic) components of coronavirus anxiety, namely dizziness (item 1), sleep disturbances (item 2), tonic immobility (item 3), appetite changes (item 4), and nausea and abdominal distress (item 5). None of the items include cognitive (e.g., worry), behavioral (e.g., avoidance), or emotional (e.g., fear) aspects of anxiety, although items referring to these features of anxiety were included in the initial pool of 20 candidate items. The five somatic items were included in the final version of the CAS because they had the strongest factor loadings on the first component of the principal component analysis, high pattern/structure and communality coefficients, as well as low cross-loadings. As argued by the author of the scale (Lee, 2020b), two CAS items (dizziness and tonic immobility) capture physiological arousal in response to coronavirus-related fear, two items (sleep disturbances and appetite loss) capture the somatic symptoms caused by intense, persistent worry about the coronavirus and are more closely associated with anxiety, and the fifth item (nausea and abdominal distress) captures somatic reactions resulting from either fear (e.g., thoughts of immediate danger) or anxiety (e.g., intense worry). Therefore, the choice of five items included in the final version of the CAS appears to be justified from both psychometrical and theoretical perspective. The original study in an adult US sample (Lee, 2020a) supported high internal consistency of the CAS, a one-factor structure, convergent validity evidence, and measurement invariance across gender and age. However, two items (item 2: sleep disturbances, and item 4: appetite changes) operated differently across Whites and non-Whites. More specifically, these two items had stronger loadings on the latent coronavirus anxiety factor among Whites than among the non-Whites (Lee, 2020a). A subsequent psychometric examination in a new adult US sample showed that a one-factor structure was invariant across age, gender, and race (Lee et al., 2020).

## CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

The favorable psychometric properties in the initial studies and its brevity made the CAS an increasingly popular measure of coronavirus-related anxiety during the pandemic. The scale has been translated in dozens of languages and psychometrically evaluated in numerous countries, such as Bangladesh (Ahmed et al., 2022), China (Chen et al., 2021), Cuba (Broche-Pérez et al., 2022), Poland (Skalski et al., 2021), Portugal (Magano et al., 2021), South Korea (Choi et al., 2022), and Turkey (Evren et al., 2022). Furthermore, the scale has been used in several studies examining mental health and well-being during the COVID-19 pandemic around the world, including some cross-national research (e.g., Linehan et al., 2020). In most studies, the results of confirmatory factor analyses have supported the one-factor model of the CAS (Broche-Pérez et al., 2022; Choi et al., 2022; Evren et al., 2022; Lee, 2020c). However, in some countries, modifications have been made (e.g., correlating residuals between a pair of items) to achieve an excellent model fit (see Ahmed et al., 2022; Magano et al., 2021; Vinaccia et al., 2022). Furthermore, in several studies the upper limit of the 90% confidence interval for the RMSEA was above the acceptable cut-off (i.e., .10) (Chen et al., 2021; Skalski et al., 2021). Therefore, the internal structure of the CAS warrants further research, as the simple one-factor solution does not hold in all countries.

### **The Importance of Testing Cross-National Measurement Invariance of the CAS**

Despite the widespread use of the CAS in many countries, to our knowledge, its cross-national measurement invariance has been investigated only in two studies to date (Caycho-Rodríguez et al., 2022; Lieven, 2021), both of which have some important limitations. Caycho-Rodríguez and colleagues (2022) examined the factor structure and measurement invariance of the Spanish version of the CAS across twelve Latin American countries. The authors found that the original one-factor model fitted poorly in most countries (RMSEA values were above .10 in

## CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

eleven out of twelve countries), and that residuals of item 4 (appetite loss) and item 5 (nausea and abdominal distress) were strongly associated. After removing item 5, the one-factor model of the abbreviated CAS (CAS-4) provided an excellent fit to the data in most countries. The authors tested the cross-national measurement invariance of the CAS-4 and concluded that both metric and scalar invariance were met, although the large drop in RMSEA between metric and configural model (.04) suggested that the metric invariance was not supported, and that testing for partial metric invariance should have been conducted. Another important limitation of this study is that convergent validity of the CAS scores in relation to other measures was not examined. Lieven (2021) investigated the measurement invariance of the CAS across 25 countries from six continents, and contrary to Caycho-Rodríguez et al. (2022) found that the original one-factor model provided acceptable or good fit to the data in all countries, and that this model was fully invariant across countries. However, this study also has some limitations, as the majority of study participants dwelled in high-income, Western countries. Furthermore, Lieven's study also did not include convergent measures. To sum up, the cross-cultural applicability of the CAS is still largely unknown, especially in non-Western samples.

There are several issues with the CAS's item content that raise a question whether this scale is appropriate for the assessment of coronavirus anxiety in different cultural settings. The CAS was developed in a Western context, i.e., on a sample of US adults and relying on the somatic symptoms of fear and anxiety typical for people in Western cultures. For example, the content validity of the CAS items in the original study (Lee, 2020a) has been supported by the relevance of the symptoms captured by the CAS for diagnoses of mental disorders such as panic disorder, generalized anxiety disorder, post-traumatic stress disorder, and major depressive disorder as defined within DSM-5 (APA, 2013), which has been repeatedly criticized as an

## CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

ethnocentric and culturally insensitive approach to mental disorders (e.g., Bredström, 2019; Marecek & Lafrance, 2021). Although physical and physiological symptoms are common features of anxiety worldwide (Kirmayer, 2001), the presentation, experience, and reporting of somatic symptoms of anxiety may vary substantially across cultures (Marques et al., 2011). For example, somatic symptoms appear to have a more central role in anxiety among individuals from non-Western cultures than from Western cultures (Lewis-Fernández et al., 2010). Another potential issue with the CAS is the inclusion of common somatic symptoms of depression, such as problems with sleep and appetite, which may be more frequently endorsed in non-Western cultures (e.g., Ryder et al., 2008). Cross-cultural variations in physical and physiological symptoms of anxiety warrant careful research on the cross-cultural validity of the CAS as a measure of coronavirus-related anxiety.

### **The Present Study**

The main goal of the current study is to investigate the cross-national measurement invariance of the CAS across 60 countries. Measurement invariance is necessary to ensure that a given instrument measures the construct of interest in the same way across different groups (Han et al., 2019). This becomes especially important: a) when the construct of interest appears to be ubiquitous (as it is related to the pandemic) and does not discriminate between groups; and b) when the groups comprise culturally diverse populations. In this case, the translation of the questionnaire and cultural differences may affect a proper understanding of the items and could then reveal biased estimates of anxiety. Although unable to guarantee that the instrument's adaptations across languages are parallel, measurement invariance tests provide evidence of similarity of its functioning (Leitgöb et al., 2022). In addition, we aimed to expand on previous research by examining the convergent validity evidence of the CAS scores in relation to

## CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

alternative measure of coronavirus-related mental health (i.e., Fear of COVID-19 Scale; FCV-19S), and in relation to the abbreviated version of the most frequently used measure of subjective well-being (Satisfaction with Life Scale; SWLS).

### Method

#### Participants and Procedure

The data was collected as part of a larger project led by [MASKED FOR REVIEW] at the [MASKED FOR REVIEW]. The project aimed at examining the impact and experiences of COVID-19 among young people and established adults in an international perspective across major geographical regions. Due to its exploratory nature, the current XX study was not pre-registered. The study was approved by the [MASKED FOR REVIEW] Ethical Committee. The samples in most countries were recruited using convenience and snowball sampling methods. **In each country the participant's primary language was used and online surveys were administered in all countries.** The sample characteristics, the CAS mean scores, and Cronbach Alphas are listed in Online Supplementary Materials (OSM), Section A. In the course of data cleaning, we removed all observations that contained more than two missing responses on the five CAS items which comprised 0.4% of the total sample. The final sample size was  $N = 21,513$  respondents in 60 countries ( $n_s = 48$  to 1,287 per country).

#### Measures

The Coronavirus Anxiety Scale (CAS; Lee, 2020a) consists of the following five items:

1. I felt dizzy, lightheaded, or faint, when I read or listened to news about the coronavirus;
2. I had trouble falling or staying asleep because I was thinking about the coronavirus;
3. I felt paralyzed or frozen when I thought about or was exposed to information about the coronavirus;
4. I lost interest in eating when I thought about or was exposed to information about the



## CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

coronavirus; and 5. I felt nauseous or had stomach problems when I thought about or was exposed to information about the coronavirus. Participants are asked to rate the frequency of each symptom over the last two weeks on a 5-point scale with options *Not at all (1)*, *Rare, less than a day or two (2)*, *Several days (3)*, *More than 7 days (4)*, and *Nearly every day (5)*.

The Fear of COVID-19 Scale (FCV-19S; Ahorsu et al., 2022) is a 7-item questionnaire (e.g., *It makes me uncomfortable to think about coronavirus-19*). Items are rated on a 7-point scale (from 1 = *Strongly disagree* to 7 = *Strongly agree*).

The abbreviated, three-item version of the Satisfaction with Life Scale (SWLS-3; Kjell & Diener, 2021) was used to assess global life satisfaction. The SWLS-3 is a brief version of the original SWLS (Diener et al., 1985), and it includes the first three items of the original scale (*In most ways my life is close to my ideal; The conditions of my life are excellent; I am satisfied with my life*). Items are rated on a 7-point scale (from 1 = *Strongly disagree* to 7 = *Strongly agree*).

### **Questionnaire Translation**

Several translations of the CAS (e.g., Spanish, Japanese, Polish) and most of the FCV-19S and SWLS-3 translations were already available at the beginning of the project, so where possible, the existing translations were used. In countries where the questionnaires' translations were not identified, a committee approach was used (e.g., van de Vijver, 2019). The committee approach to translation attempts to decrease the cultural bias inherent in the native language, by introducing collaborative, consensus translation efforts. In each country, at least two individuals with knowledge of the target language, society, and culture, translated the questionnaires independently. In the next step, the translated versions were compared, the differences were resolved by consensus, and the final version was accepted after the discussion on the items' appropriateness.

### **Data Analysis**

Descriptive analysis of the CAS items demonstrated highly skewed distributions in all the groups (for the distributions, see OSM Section B). The response option “*Not at all*” was chosen disproportionately more often than all the others (64%-84% across items), and the second most chosen option was “*Rare, less than a day or two*” (10%-19%), while the other options were chosen only occasionally. The distribution of responses is unsurprising as the CAS measures severe conditions in the general population. However, as a consequence, the responses cannot be treated as a statistically normal continuous distribution, but rather as categorical. Due to low frequencies of several response options combined with the small sample sizes in many groups, the corresponding contingency tables were sparse (i.e., contained multiple zero frequencies). The sparse response patterns are highly problematic for multiple group latent variable analyses (Forero & Maydeu-Olivares, 2009). Therefore, we could not model this as an ordinal categorical (polytomous) scale. Thus, we decided to dichotomize the responses into 0 for “*Not at all*” and 1 for all the other response options (see DiStefano et al., 2021). Dichotomization did not lead to a substantial loss of information because the crucial meaningful source of variance came from the difference between occurrence and non-occurrence of the five symptoms.

### **Measurement Invariance**

In order to test measurement invariance of the CAS we employed multiple approaches, because the samples were convenient and diverse, and the study included participants from very different populations. We used both conventional exact invariance tests, as well as a more recent procedure of alignment (Asparouhov & Muthén, 2022; Leitgöb et al., 2022), including weighted least squares and Bayesian versions. The test of the exact invariance with the dichotomous indicators involves comparison of the two models: configural and scalar invariance both

## CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

estimated with weighted least squares mean and variance adjusted (WLS). Intermediary metric invariance in case of binary indicators was not viable (Wu & Estabrook, 2016). The goodness of fit was indicated by  $CFI > .90$  and  $RMSEA < .08$  (Hu & Bentler, 1999). Configural model tests for an overall similarity of factor structure across groups, whereas scalar invariance models require equality of the factor loadings and item thresholds across groups. If the difference between the fit of the two models lies within the cutoff values of  $\Delta CFI < .008$  and  $\Delta RMSEA < .05$  (Rutkowski & Svetina, 2017;  $\Delta CFI$  comes from adding two cutoffs of .004 for factor loadings and threshold constraints), then full scalar invariance is indicated. This would imply comparability of the unstandardized regression coefficients as well as latent means across groups. We also applied the alignment procedure (Asparouhov & Muthén, 2014) because the diverse and large group differences were unlikely to show exact measurement invariance. Alignment is an iterative procedure that estimates a configural model without any group constraints and, similar to the factor loadings rotation in exploratory factor analysis, finds the most invariant set of parameters possible without changing the model fit to the data (Asparouhov & Muthén, 2014; 2022). The alignment procedure established configural invariance model using WLSMV and then rotated parameters to achieve the highest possible invariance. We followed a criterion of the maximum of 25% non-invariant parameters to establish a sufficient level of measurement invariance. In addition, we considered  $R^2$ , its higher values reflect the degree to which group differences in loadings and thresholds are explained by corresponding differences in factor variances and means. Higher values indicate higher degree of invariance. At first, we ran a free mode alignment, identified a group with a latent mean close to zero (which happened to be Malaysia), and then ran a fixed mode alignment using the identified group as a baseline.

### **Resampling Analysis**

## CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

The sample sizes of groups were very different ranging from 48 respondents in Uganda up to 1,287 in Bangladesh. This might have caused unequal influence of different groups on the overall result of the measurement invariance tests (Yoon & Lai, 2018). Therefore, another robustness check was implemented. Specifically, we employed a resampling technique described by Yoon and Lai (2018). Resampling included drawing random samples of the equal size from each group (1) and fitting multiple group factor analysis using this new sample (2). Points 1 and 2 were repeated a large number of times (we used 500 runs), and finally, the fit statistics and estimated parameters were summarized across all the runs of the resampled models. The resampling technique was applied both to the conventional tests and to the alignment. One important limitation of this technique is that the resampled  $N$  gets as small as the smallest group in the data. Since our dataset contains groups with as little sample sizes as 48, we first dropped all the groups with the sample sizes below 99 observations. It resulted in 52 groups for the resampling analysis. The resampling was performed in two ways. First, we precisely followed the instructions given by Yoon and Lai (2018) and sampled every group with the same number of observations, where  $N$  was equal to the smallest sample in the data. Secondly, we opted to use resampling *with replacement*, so that the group sample sizes can be larger. This enabled the models to be estimated using indefinitely large samples, so we chose to use 500 and 1000 observations per group. Finally, led by the inconclusive results of the planned tests, we ran a follow-up analysis applying several alternative methods of measurement invariance testing. The details are discussed in the corresponding section.

### **Convergent Validity**

Another task was to relate the CAS scores to FCV-19S and SWLS scores. A strong, positive association between the CAS and FCV-19S, and a negative, but weak association

## CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

between CAS and SWLS were expected in every country. Before testing the associations of the CAS with the other scales, measurement invariance on these scales was also assessed. Therefore, we tested it using the already mentioned methods of exact, approximate (alignment) invariance, as well as with a resampling technique. Finally, we computed correlations between these latent variables. Most analyses were conducted using Mplus 8.8 (Muthén & Muthén, 1998-2022), preparation of data, integration of results, and visualizations were conducted in R (R core team, 2022), in particular, using ‘MplusAutomation’ (Hallquist & Wiley, 2021), ‘MIE’ (Rudnev, 2022), ‘lavaan’ (Rosseel, 2012), and other packages (for the full list of the packages see the OSM). The data for the present study are available upon request from the corresponding author. All the reproduction codes are available at the Open Science Framework directory [https://osf.io/7gnz9/?view\\_only=7c345b583bb247ceb45194f6efb40392](https://osf.io/7gnz9/?view_only=7c345b583bb247ceb45194f6efb40392).

### Results

#### Exact Invariance

We tested a simple one-factor model of CAS without any correlated residuals. Before checking for measurement invariance, we tested the model in each country separately using confirmatory factor analyses. The results are listed in OSM Section C. The model fit the data sufficiently well in almost all the countries – the  $\chi^2$  tests  $p$ -values were higher than .05 in all but seven countries. Six of those had relatively large sample sizes (> 300) and good model fit in terms of CFI and RMSEA, which made significant  $\chi^2$  values less relevant. The samples from Uganda and Taiwan showed an unacceptable fit in terms of both RMSEA and  $\chi^2$  values. Also, despite the model showing a good fit to the data, the estimated factor’s variance was negative in the Czech Republic. These three countries were dropped from further analysis. In addition, Mozambique was dropped because the preliminary multiple group models persistently estimated

## CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

a negative variance of the CAS factor in this sample. The model using the pool of 56 remaining samples revealed a very good fit to the data, CFI/TLI = .998/.996, RMSEA = .042 (90% CI = .037, .047), SRMR = .018.

Next, we turned to testing the exact measurement invariance. Table 1 lists the results of the configural and scalar Multigroup Confirmatory Factor Analysis (MGCFA) model fit in 56 countries. The results showed that the configural invariance model had a very good fit to the data. Scalar invariance showed acceptable CFI and SRMR statistics, RMSEA was .085, which is higher than recommended upper cutoff value of .08 (Hu & Bentler, 1999). The examination of parameter estimates in the configural model demonstrated that thresholds of item 1 (*dizziness*) and item 3 (*tonic immobility*) showed the largest differences across groups. Therefore, we fitted a partial scalar invariance model (Byrne et al., 1989), withdrawing the equality constraints on these two items' thresholds. The partial invariance model included relaxed constraints on the two thresholds, at the same time the corresponding "scales" were fixed at 1. This way, the model had the same number of degrees of freedom as the scalar model, but was less constrained.<sup>1</sup> The resulting partial invariance model had a good fit to the data by all fit indices. The differences in CFI and RMSEA between configural and partial scalar invariance models were within the recommended range (i.e.,  $\Delta\text{CFI} = .002$ ,  $\Delta\text{RMSEA} = .009$ ). Other than that, scalar invariance model estimated non-significant variances of the latent factor in approximately half of the samples, which signals a problematic solution. Nevertheless, we can conclude that partial invariance could be supported and the means and regression coefficients of the COVID anxiety factor can be compared across cultural groups.

### **Resampling of the Exact Invariance Tests**

---

<sup>1</sup> In the delta parameterization for CFA with categorical indicators, scales are additional parameters replacing residuals which are normally not identified.

## CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

As already mentioned, we applied two resampling approaches – without replacement and with replacements. To make these resampling solutions comparable to the non-resampled models, we further excluded four countries with the smallest samples (Costa Rica, Bosnia and Herzegovina, Nigeria, and Qatar, all  $n < 99$ ), therefore, the number of countries was reduced to 52 (see Table 2). Several notable points arose from this analysis. The conventional invariance tests on the slightly reduced sample showed almost identical results. Every resampling supported well-fitting configural models, rejected full scalar models, and confirmed partial invariance as its fit was only slightly worse than the fit of the configural models. In resampling runs with  $n = 500$  and  $n = 1000$ , the partial invariance model had marginal RMSEA values of .083 and .084, respectively. These results imply that if we had balanced and large samples in each country partial invariance would probably not be supported. This adds some ambiguity to the results, and asks for further robustness tests. Therefore, we turned to an approximate invariance testing using the alignment procedure.<sup>2</sup>

### Measurement Invariance Testing Using Alignment

The model fit in the alignment procedure is the same as in the configural model (see Table 1). The results of the alignment procedure are listed in Table 3. All the factor loadings occurred to be approximately invariant across all 56 countries. Only 9% of the parameters were non-invariant, which is substantially lower than the recommended upper cutoff of 25%. The most non-invariant parameter was threshold of item 1 (*dizziness*). It was non-invariant in 21 countries. Item 3 (*tonic immobility*) threshold was non-invariant in 14 countries. This coincides

---

<sup>2</sup> Many resampled solutions had a high number of non-positive definite matrices (namely negative variances of observed variables), which were more common in smaller samples and less common in larger samples. Sparse contingency tables likely caused this, as some cross-tabulations had frequencies close to zero, and the resampling reproduced this issue. Given that residuals of binary observed indicator estimated with Delta parameterization are not typical parameters and do not necessarily point to a problem, we considered the results of the resampling solutions sufficiently reliable.

## CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

with the results of the partial invariance analysis. The other thresholds had negligible numbers of non-invariant countries. Even in countries with the weakest invariance, namely, Brazil and Romania, only three thresholds were non-invariant whereas two other thresholds and all the five factor loadings were invariant. Overall, the alignment demonstrated a high degree of approximate measurement invariance.

Next, we ran simulations of alignment with parameters estimated in the current model (Asparouhov & Muthén, 2014). Simulations can demonstrate whether the current model is able to correctly estimate latent means and variances. We ran simulations for different numbers of observations per group – 100, 500, and 1000. The general approach assumes a balanced design (equal sample sizes in each group), which is not the case in our data. This might have resulted in overly optimistic estimates of the reliability of alignment. Given the unbalanced nature of our data, actual sample sizes were used in the simulation as well. There were two measures of replication. For the first one, each set of estimated means is correlated with the (true) means found in the original alignment, and then these correlations are averaged across all samples. Another way to aggregate the results is to compute average estimates across all the generated samples, and only after that correlate these average estimates to the true means. Both approaches should arrive at correlation equal or greater than .98 to support reliability of the alignment results (ibid.). The results for different sample sizes are listed in OSM, Section D. The simulations with actual  $n$  and  $n = 100$  had a large proportion of NPD solutions (which is not necessarily problematic, see footnote 2). It was not the case for balanced and larger samples though. The correlations of true and estimated means were acceptable ( $r > .98$ ) in balanced sample condition with  $n = 500$  and higher. It follows that the latent means could have been reliable if the sampling was balanced, and the samples were at least 500 per group. But for the current sampling



## CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

approach, the model might have slightly misestimated the latent means of the anxiety factor. In contrast, the correlations between true and estimated variances were extremely small, the highest estimated correlation for a balanced sample and  $n = 1000$  was only .736. This result implies that the variances of the latent variables could not be reliably estimated by the current alignment model. Yet, the other measure of reliability of variances, correlation of average estimated variance with the population values, shows higher reliability, but again reaches an acceptable level only at  $n = 500$ .

Finally, we also applied resampling to the alignment. We successfully computed 491 resampled alignments ( $n = 500$  for each group, with replacement)<sup>3</sup>, which resulted in 500 sets of means for each of 52 groups. The replication rates were high. The average correlation was  $r = .991$  and its SD = .002. The correlation of average estimates with the true values was  $r = .999$ . As mentioned above, a similar method, but for simulations rather than resampling, was suggested by Asparouhov and Muthén (2014), who also provided a cutoff of  $r = .98$  as a measure of successful replication. Although there are no correlation guidelines for the resampling, the correlations seem high enough to suggest that the results of alignment might have revealed the unbiased means. The correlations between the estimated variances in resampling and true variances were also very high ( $r = .999$ ).

Overall, however, the results were inconclusive. On one hand, both exact invariance tests and alignment suggested a high degree of (approximate) invariance. On the other hand, however, the solutions contained many NPDs, means and variances differed across methods. In order to add more certainty, we also ran a more flexible Bayesian alignment (Asparouhov & Muthén,

---

<sup>3</sup> Resampling using  $n = 99$  without replacement resulted in 91% solutions with NPD, yet the correlations of true and replicated means were relatively high; correlation of averages with the true means was  $r = .999$  (and  $r = .994$  for variances), average of correlations with true means was  $r = .972$  and SD = .007.

## CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

2014), maximum likelihood alignment, and Bayesian approximate invariance tests (BSEM – see details in Van de Schoot et al., 2014). The details of these analyses are listed in OSM Section D. The correlations between latent means and variances estimated with six different methods are listed in Table 4. The means, including a simple group mean score, converged. The correlation between WLSMV alignment and partial exact invariance is  $r = .96$ . Figure D1 in OSM demonstrates similarity of means estimated by these two methods and their similarity to average summative scores. The other methods estimated means that correlated with each other  $r = .90$  or higher, except the means estimated by the full scalar model. The latter is expected as the model was rejected due to its fit to the data and estimation problems.

Unlike the means, the estimation of variances produced very different results (see Table 4 above the diagonal). The variances of the summative index demonstrated the weakest correlations with variances estimated by the other methods. Nevertheless, partial invariance model and different methods of alignment estimated factor variances that correlate around  $r = .90$  which is sufficient for most practical purposes. Overall, all the non-exact invariance methods estimated highly consensual means and variances, and arrived at similar conclusions of a high level of approximate/partial invariance of the CAS. For further analysis of convergent validity, we opted to use WLS alignment because it seems to represent the most consensual pattern of means and variances compared to all the other methods.

### **Convergent Validity Evidence**

Before examining the relations between CAS with FCV-19S and SWLS-3 scales, the latter also needed to be tested for invariance across countries. Although both scales have been shown to be invariant across various countries in other studies (e.g., Sawicki et al., 2022), it was necessary to demonstrate it with the current data to ensure that the within-country correlations

## CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

were not biased (requires metric invariance) and that country-level correlations can be interpreted (requires scalar invariance). The results listed in OSM Section E demonstrate that both scales had a sufficient level of approximate invariance.

Given the inconsistent results of the tests of invariance and also to test the relations between CAS and the two convergent measures, we used methodological triangulation. That is, we applied three different strategies. The first strategy used predicted individual factor scores and then computed correlations. The second strategy involved merging MGCFA models with established levels of invariance into a single MGCFA model and estimating covariances between latent variables simultaneously. Finally, we made use of the possibility of conducting alignment for the entire model with factors for CAS, SWLS-3, and FCV-19S. The results are summarized in Table 5 (for details see OSM Section F). The correlations between the two factors of the FCV-19S (physiological arousal and psychological distress) and CAS were positive and significant both at the individual and at the country level in all countries except Pakistan (as well as Nepal – for psychological distress factor and USA for physiological arousal factor). The exclusion of these countries, however, barely changed the results, which supported the convergent validity of CAS. Correlations of CAS with SWLS-3 were much weaker and less stable across countries. The average correlation across different methods was between  $-.07$  and  $-.15$ , with large standard deviations pointing to their marginal difference from zero. Indeed, the correlation was significantly different from zero only in a small fraction of countries. However, in almost all groups, this correlation was consistently negative (Table 5).

Finally, we checked how consistent the correlations were between different methods. Consistency (correlation of Fisher-standardized correlations estimated by different methods) of within-country correlations estimated with factor scores vs. MGCFA was  $r = .97$  (psychological

## CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

distress factor);  $r = .97$  (physiological arousal factor);  $r = .94$  (life satisfaction). Likewise, meta-correlations based on estimates from alignment factor scores and MGCFA were .93; .80; .96; and the ones based on MGCFA and the common alignment were .93, .80, .96, respectively. Mean differences between values of correlation coefficients were .10 and .08 for physiological arousal and psychological distress factors, respectively, and .04 for life satisfaction.

### **Discussion**

The present study is the largest cross-national investigation of CAS to this date, examining the factor structure, measurement invariance, and convergent validity evidence of the scale on a sample of 21,513 respondents from all five continents and 60 countries. From a methodological standpoint, this is the first study to apply both exact and approximate approaches to testing measurement invariance of the CAS, and to examine cross-national associations between the CAS and two concurrent measures of mental health and well-being. The CAS is a new measure designed to assess COVID-19 related anxiety, but its content can easily be adapted for other health crises and negative life experiences. Furthermore, the CAS's brevity and ease of administration makes it a promising tool for large surveys and cross-national research. Therefore, testing its measurement invariance across a large set of countries can offer valuable insights for future cross-national research.

Overall, the scale demonstrated the theoretically expected single-factor structure in the vast majority of countries included in this research and showed consistent strong, positive associations with fear of COVID-19 and negative (albeit less consistent) links with satisfaction with life. The one-factor model of the CAS had a poor fit to the data only in a few samples, whereas an excellent fit was observed in the vast majority of countries. This is in line with the findings by Lieven (2021) that were drawn from 10,232 respondents across 25 countries. Low

## CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

RMSEA values of the single-factor model observed in most countries in the present study were not in accordance with the results of Caycho-Rodríguez et al.'s study (2022), who found large RMSEA values in eleven out of twelve Latin American countries. Although RMSEA values can be overestimated in models with small degrees of freedom (Kenny et al., 2015), such is the case with the one-factor model of the CAS ( $df = 5$ ), these discrepancies in RMSEA values in our and Caycho-Rodríguez et al.'s study (2022) suggest that the single-factor model might operate differently across samples and languages, thus warranting further research on the CAS's structure. However, it is also important to note that our results are not directly comparable to previous research due to several methodological and statistical differences. Namely, Lieven (2021) and Caycho-Rodríguez et al. (2022) used maximum likelihood (ML) and robust ML estimation method, respectively, whereas we relied on the WLSMV. The data collection in two previous cross-cultural studies were restricted to March 2021 (Lieven, 2021) and February-March 2021 (Caycho-Rodríguez et al., 2022). On the other hand we used data that was collected from the beginning until the end of 2021. Most importantly, we dichotomized the data and treated them as binary, whereas the earlier cross-national studies relied on the original five response options in their analyses.

The results of our study showed that cross-national measurement invariance of the CAS needs to be approached with caution and that alternative methods of assessing invariance should be considered when evaluating this scale across a wide range of countries. There were several important details which we explored in depth. First, the results of various approaches to testing measurement invariance of CAS resulted in generally optimistic but unstable results. Despite good fit of the models, they often arrived at inadmissible solutions and, depending on the method, varied in the estimated levels of COVID-19-related anxiety. In order to arrive at a stable

## CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

result, we applied a methodological triangulation which involves application of several methods to analyze the same data (Heesen et al., 2019). The results suggested that the most stable solution was found in Bayesian alignment.

Virtually all methods pointed on the lower invariance of thresholds for items relating to *dizziness* (item 1) and *tonic immobility* (item 3). These results indicate that these two symptoms appear across samples to different degrees regardless of the overall anxiety. In other words, it might imply that culture moderates the expression of anxiety through *dizziness* (feeling dizzy, lightheaded, faint) and *tonic immobility* (feeling paralyzed, frozen) or at least regulates the self-reports of these conditions. Contrary to the other items (sleep disturbances, appetite loss, nausea/abdominal distress) capturing somatic symptoms caused primarily by intense worry about the coronavirus, dizziness and tonic immobility refer to physiological arousal in response to intense fear. Both dizziness (and other dissociative symptoms; Schalinski et al., 2015) and tonic immobility (e.g., Abrams et al., 2008) might occur in situations of perceived inescapability, threat of death, and in the context of panic attacks. Although both dizziness and tonic immobility are involuntary, automatically activated defense behaviors (Kozłowska et al., 2015), they were assessed via self-report in the present study, which might bring out some cross-cultural differences in the catastrophic interpretation of these symptoms (see Hinton & Polack, 2009) and provoke culture-specific meanings of these bodily symptoms (Hofmann & Hinton, 2014). Furthermore, the linguistic meaning of dizzy/lightheaded/faint (item 1) and paralyzed/frozen (item 3) probably varies more across languages compared to sleeping (item 2) and eating problems (item 4), which refer to basic physiological needs essential for human survival. Thus, items referring to eating and sleeping can be expected to have higher translatability and a more similar meaning across languages, as they capture motivational states with a high level of

## CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

universality across languages (e.g., Saucier et al., 2014). In addition, nausea/stomach problems (item 5) are closely associated with an affective state of disgust also essential for survival (Panksepp, 2007), and thus also can be expected to have less variation across languages than terms related to dizziness and tonic immobility.

The triangulated analysis of CAS association with fear of COVID-19 and satisfaction with life did not show exactly the same strength of correlation coefficients in every studied country, but these correlations' directions were consistent with our expectations. As expected, the coronavirus anxiety (as measured by the CAS) had strong positive correlations with physiological arousal and psychological distress factors of the Fear of COVID-19 scale, and weak negative correlations with satisfaction with life. The findings support the convergent validity of the CAS scores, as it has been shown that somatic symptoms of coronavirus anxiety are more closely associated with psychological and somatic aspects of COVID-19 related fear (i.e., FCV-19S) than with a measure of people's overall evaluation of their life. Life satisfaction judgments are relatively stable and strongly influenced by objective conditions and chronically accessible information (Pavot & Diener, 2008), thus weak correlations with context-specific fears and anxiety are expected. The magnitude of correlations between CAS and two FCV-19S factors found in the present research is comparable to those observed in only a few previous studies that adopted a two-factor structure of the FCV-19S (e.g., Magano et al., 2021), and suggest that the CAS and FCV-19S measure related, yet distinct constructs. The CAS focuses exclusively on somatic aspects of anxiety, whereas the FCV-19S includes items covering somatic symptoms of fear (insomnia, heart palpitations, and clammy hands), but also psychological aspects (being afraid, uncomfortable, afraid to die, and nervous/anxious), thus their relatively modest intercorrelation is as expected.

## CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

Based on our results and some previous findings on the structure of the FCV-19S (e.g., Sawicki et al., 2022) and the CAS (Lieven, 2021) across cultures, it can be concluded that the CAS is a better alternative for cross-cultural research than FCV-19S. However, it has to be noted that the CAS is limited in its scope (no psychological, cognitive, or social dimensions of anxiety are included), so it would be preferable to complement this scale with measures that capture other aspects of anxiety beyond somatic symptoms.

### **Constraints on Generality**

Several important limitations of the present study should be noted. First, although we covered a wide range of countries across the globe, the participants were recruited using convenience sampling and the samples in most countries included mostly young adults. This limits the generalizability of our findings as anxiety symptoms might differ substantially across age groups (e.g., Carlucci et al., 2018; Teachman & Gordon, 2009). Second, the sample sizes varied greatly across countries, which poses a challenge to testing cross-national measurement invariance. Previous studies have shown that unequal sample sizes across groups might jeopardize the results of invariance testing and lead to biased invariance findings (Yoon & Lai, 2018). Although we applied a resampling strategy (i.e., random samples of balanced groups), future studies should aim to recruit more balanced samples across cultures in terms of gender, age, and socioeconomic status. Third, the current study demonstrated characteristics of the CAS in the general population but the study's findings are limited to indicating the mere presence of anxiety conditions.. A future study targeted at the affected populations could focus on the viability of the scale for indicating severity of each condition. Fourth, all the CAS items were not reverse-coded which might have resulted in a confusion between measurement of the anxiety and response styles which in turn are known to boost reliability and unidimensionality. On one hand,



## CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

our conclusions on comparability of CAS might indicate the invariance of response style in addition to invariance of the COVID-19 anxiety; on the other hand, there is no reason to expect close relations between response style and anxiety, hence, the confusion is unlikely.

Additionally, dichotomization of items could partially mitigate the response style bias, removing the scale use variance. Fifth, the convergent validity evidence of the CAS scores was evaluated only in relation to life satisfaction and fear of COVID-19. Future studies should strive to examine the relationships between the CAS and other well-established measures of anxiety (for a review of self-report anxiety scales see Wall & Lee, 2022). Finally, we did not explore potential factors that might have influenced the measurement invariance results, i.e., source of non-invariance of items 1 and 3. A recently developed procedure for disentangling different sources of item bias - the culture, comprehension, and translation bias procedure (CCT; Bader et al., 2021) - might be a useful resource for future studies aimed at understanding whether the lack of invariance on certain CAS items results from translation bias or from systematic differences in social and cultural contexts between groups. Despite these limitations, the present study suggests that the CAS is a valid measure of somatic symptoms of health-related anxiety for purposes of cross-national research.

**References**

- Abrams, M. P., Carleton, R. N., & Asmundson, G. J. G. (2012). Tonic immobility does not uniquely predict posttraumatic stress symptom severity. *Psychological Trauma: Theory, Research, Practice, and Policy*, *4*(3), 278–284. <https://doi.org/10.1037/a0023272>
- Ahmed, O., Faisal, R. A., Sharker, T., Lee, S. A., & Jobe, M. C. (2022). Adaptation of the Bangla version of the COVID-19 Anxiety Scale. *International Journal of Mental Health and Addiction*, *20*(1), 284–295. <https://doi.org/10.1007/s11469-020-00357-2>
- Ahorsu, D. K., Lin, C. Y., Imani, V., Saffari, M., Griffiths, M. D., & Pakpour, A. H. (2022). The fear of COVID-19 scale: Development and initial validation. *International Journal of Mental Health and Addiction*, *20*(3), 1537–1545. <https://doi.org/10.1007/s11469-020-00270-8>
- Aknin, L., De Neve, J.-E., Dunn, E., Fancourt, D., Goldberg, E., Helliwell, J. F., Jones, S. P., Karam, E., Layard, R., Lyubomirsky, S., Rzepa, A., Saxena, S., Thornton, E. M., VanderWeele, T. J., Whillans, A. V., Zaki, J., Karadag Caman, O., & Ben Amor, Y. (2022). Mental health during the first year of the COVID-19 pandemic: a review and recommendations for moving forward. *Perspectives on Psychological Science*, *17*(4), 915–936. <https://doi.org/10.1177/17456916211029964>
- American Psychiatric Association (APA). (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Author.
- Asmundson, G. J. G., & Taylor, S. (2020). Coronaphobia: Fear and the 2019-nCoV outbreak [Editorial]. *Journal of Anxiety Disorders*, *70*, Article 102196. <https://doi.org/10.1016/j.janxdis.2020.102196>

## CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

- Asparouhov, T., & Muthén, B. (2014). Multiple-group factor analysis alignment. *Structural Equation Modeling, 21*(4), 495–508. <https://doi.org/10.1080/10705511.2014.919210>
- Asparouhov, T., & Muthén, B. (2022). *Multiple group alignment for exploratory and structural equation models.*
- Bader, M., Jobst, L. J., Zettler, I., Hilbig, B. E., & Moshagen, M. (2021). Disentangling the effects of culture and language on measurement noninvariance in cross-cultural research: The culture, comprehension, and translation bias (CCT) procedure. *Psychological Assessment, 33*(5), 375–384. <https://doi.org/10.1037/pas0000989>
- Bredström, A. (2019). Culture and context in mental health diagnosing: Scrutinizing the DSM-5 revision. *Journal of Medical Humanities, 40*, 347–363. <https://doi.org/10.1007/s10912-017-9501-1>
- Broche-Pérez, Y., Fernández-Castillo, E., Fernández-Fleites, Z., Jiménez-Puig, E., Vizcaíno-Escobar, A., Ferrer-Lozano, D., Martínez-Rodríguez, L., & Martín-González, R. (2022). Adaptation of the Cuban version of the Coronavirus Anxiety Scale. *Death Studies, 46*(3), 603–607. <https://doi.org/10.1080/07481187.2020.1855610>
- Byrne, B. M., Shavelson, R. J., & Muthén, B. (1989). Testing for the equivalence of factor covariance and mean structures: The issue of partial measurement invariance. *Psychological Bulletin, 105*(3), 456–466. <https://doi.org/10.1037/0033-2909.105.3.456>
- Carlucci, L., Watkins, M. W., Sergi, M. R., Cataldi, F., Saggino, A., & Balsamo, M. (2018). Dimensions of anxiety, age, and gender: assessing dimensionality and measurement invariance of the state-trait for cognitive and somatic anxiety (STICSA) in an Italian sample. *Frontiers in Psychology, 9*, 2345. <https://doi.org/10.3389/fpsyg.2018.02345>

## CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

- Caycho-Rodríguez, T., Valencia, P. D., Vilca, L. W., Carbajal-León, C., Vivanco-Vidal, A., Saroli-Araníbar, D., Reyes-Bossio, M., White, M., Rojas-Jara, C., Polanco-Carrasco, R., Gallegos, M., Cervigni, M., Martino, P., Palacios, D. A., Moreta-Herrera, R., Samaniego-Pinho, A., Lobos-Rivera, M. E., Figares, A. B., Puerta-Cortés, D. X., Corrales-Reyes, I. E., ... Flores-Mendoza, C. (2022). Cross-cultural validation of the new version of the Coronavirus Anxiety Scale in twelve Latin American countries. *Current Psychology*, 1–18. <https://doi.org/10.1007/s12144-021-02563-0>
- Chen, J. H., Tong, K. K., Su, X., Yu, E. W-y., Wu, A. M.S. (2021). Measuring COVID-19 related anxiety and obsession: Validation of the Coronavirus Anxiety Scale and the Obsession with COVID-19 Scale in a probability Chinese sample. *Journal of Affective Disorders*, 295, 1131–1137. <https://doi.org/10.1016/j.jad.2021.08.104>.
- Choi, E., Lee, J., & Lee, S. A. (2022). Validation of the Korean version of the obsession with COVID-19 scale and the Coronavirus anxiety scale. *Death Studies*, 46(3), 608–614. <https://doi.org/10.1080/07481187.2020.1833383>
- Diener, E., Emmons, R. A., Larsen, R. J., & Griffin, S. (1985). The Satisfaction with Life Scale. *Journal of Personality Assessment*, 49, 71–75. [https://doi.org/10.1207/s15327752jpa4901\\_13](https://doi.org/10.1207/s15327752jpa4901_13)
- DiStefano, C., Shi, D., & Morgan, G. B. (2021). Collapsing categories is often more advantageous than modeling sparse data: investigations in the CFA framework. *Structural Equation Modeling*, 28(2), 237–249. <https://doi.org/10.1080/10705511.2020.1803073>
- Evren, C., Evren, B., Dalbudak, E., Topcu, M., & Kutlu, N. (2022). Measuring anxiety related to COVID-19: A Turkish validation study of the Coronavirus Anxiety Scale. *Death Studies*, 46(5), 1052–1058. <https://doi.org/10.1080/07481187.2020.1774969>

## CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

- Forero, C. G., & Maydeu-Olivares, A. (2009). Estimation of IRT graded response models: limited versus full information methods. *Psychological Methods, 14*(3), 275–299.  
<https://doi.org/10.1037/a0015825>
- Hallquist, M. N., & Wiley, J. F. (2018). MplusAutomation: An R package for facilitating large-scale latent variable analyses in Mplus. *Structural Equation Modeling, 25*(4), 621–638.  
<https://doi.org/10.1080/10705511.2017.1402334>.
- Han, K., Colarelli, S. M., & Weed, N. C. (2019). Methodological and statistical advances in the consideration of cultural diversity in assessment: A critical review of group classification and measurement invariance testing. *Psychological Assessment, 31*(12), 1481–1496.  
<https://doi.org/10.1037/pas0000731>
- Heesen, R., Bright, L. K., & Zucker, A. (2019). Vindicating methodological triangulation. *Synthese, 196*, 3067–3081. <https://doi.org/10.1007/s11229-016-1294-7>
- Hinton, D. E., & Pollack, M. H. (2009). Introduction to the special issue: anxiety disorders in cross-cultural perspective. *CNS Neuroscience & Therapeutics, 15*(3), 207–209.  
<https://doi.org/10.1111/j.1755-5949.2009.00097.x>
- Hofmann, S. G., & Hinton, D. E. (2014). Cross-cultural aspects of anxiety disorders. *Current Psychiatry Reports, 16*(6), 450. <https://doi.org/10.1007/s11920-014-0450-3>
- Hu, L. T., & Bentler, P. M. (1999). Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Structural Equation Modeling, 6*(1), 1–55.  
<https://psycnet.apa.org/doi/10.1080/10705519909540118>
- Kenny, D. A., Kaniskan, B., & McCoach, D. B. (2015). The performance of RMSEA in models with small degrees of freedom. *Sociological Methods & Research, 44*(3), 486–507.  
<https://doi.org/10.1177/0049124114543236>

## CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

Kirmayer, L. J. (2001). Cultural variations in the clinical presentation of depression and anxiety:

Implications for diagnosis and treatment. *The Journal of Clinical Psychiatry*, *62*, 22–28.

Kjell, O. N. E., & Diener, E. (2021). Abbreviated three-item versions of the Satisfaction with Life Scale and the Harmony in Life Scale yield as strong psychometric properties as the original scales. *Journal of Personality Assessment*, *103*(2), 183–194.

<https://doi.org/10.1080/00223891.2020.1737093>

Kozłowska, K., Walker, P., McLean, L., & Carrive, P. (2015). Fear and the defense cascade: Clinical implications and management. *Harvard Review of Psychiatry*, *23*(4), 263–287.

<https://doi.org/10.1097/HRP.0000000000000065>

Lee, S. A. (2020). Replication analysis of the Coronavirus Anxiety Scale. *Dusunen Adam: The Journal of Psychiatry and Neurological Science*, *33*, 203–205.

<https://doi.org/10.14744/DAJPNS.2020.00079>

Lee, S. A. (2020a). Coronavirus Anxiety Scale: A brief mental health screener for COVID-19 related anxiety. *Death Studies*, *44*(7), 393–401.

<https://doi.org/10.1080/07481187.2020.1748481>

Lee, S. A. (2020b). Measuring coronaphobia: the psychological basis of the Coronavirus Anxiety Scale. *Dusunen Adam: The Journal of Psychiatry and Neurological Sciences*, *33*(2), 107–

108. <https://doi.org/10.14744/DAJPNS.2020.00069>

Lee, S. A., Mathis, A. A., Jobe, M. C., & Pappalardo, E. A. (2020). Clinically significant fear and anxiety of COVID-19: A psychometric examination of the Coronavirus Anxiety Scale.

*Psychiatry Research*, *290*, Article 113112. <https://doi.org/10.1016/j.psychres.2020.113112>

Leitgöb, H., Seddig, D., Asparouhov, T., Behr, D., Davidov, E., De Roover, K., Jak, S.,

Meitinger, K., Menold, N., Muthén, B., Rudnev, M., Schmidt, P. & van de Schoot, R.

## CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

(2022). Measurement invariance in the social sciences: Historical development, methodological challenges, state of the art, and future perspectives. *Social Science Research*, 102805. <https://doi.org/10.1016/j.ssresearch.2022.102805>

Lewis-Fernández, R., Hinton, D. E., Laria, A. J., Patterson, E. H., Hofmann, S. G., Craske, M. G., Stein, D. J., Asnaani, A., & Liao, B. (2010). Culture and the anxiety disorders: recommendations for DSM-V. *Depression and Anxiety*, 27(2), 212–229. <https://doi.org/10.1002/da.20647>

Lieven T. (2021). Global validation of the Coronavirus Anxiety Scale (CAS). *Current Psychology*, 1–11. <https://doi.org/10.1007/s12144-021-02583-w>

Linehan, C., Araten-Bergam, T., Baumbusch, J., Beadle-Brown, J., Bigby, C., Birkbeck, G., Bradley, V., Brown, M., Bredewold, F., Chirwa, M., Cui, J., Godoy Gimenez, M., Gomiero, T., Kanova, S., Kroll, T., MacLachlan, M., Mirfin-Veitch, B., Narayan, J., Nearchou, F., Nolan, A., ... Tossebro, J. (2020). COVID-19 IDD: A global survey exploring family members' and paid staff's perceptions of the impact of COVID-19 on individuals with intellectual and developmental disabilities and their caregivers. *HRB Open Research*, 3, 39. <https://doi.org/10.12688/hrbopenres.13077.2>

Luo, F., Ghanei Gheshlagh, R., Dalvand, S., Saedmoucheshi, S., & Li, Q. (2021). Systematic review and meta-analysis of fear of COVID-19. *Frontiers in Psychology*, 12, 661078. <https://doi.org/10.3389/fpsyg.2021.661078>

Magano, J., Vidal, D. G., Sousa, H., Dinis, M., & Leite, Â. (2021). Validation and psychometric properties of the Portuguese version of the Coronavirus Anxiety Scale (CAS) and Fear of COVID-19 Scale (FCV-19S) and associations with travel, tourism and hospitality.

## CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

*International Journal of Environmental Research and Public Health*, 18(2), 427.

<https://doi.org/10.3390/ijerph18020427>

Marecek, J., & Lafrance, M. N. (2021). Editorial introduction: The politics of psychological suffering. *Feminism & Psychology*, 31(1), 3–18. <https://doi.org/10.1177/0959353521989537>

Marques, L., Robinaugh, D. J., LeBlanc, N. J., & Hinton, D. (2011). Cross-cultural variations in the prevalence and presentation of anxiety disorders. *Expert Review of Neurotherapeutics*, 11(2), 313–322. <https://doi.org/10.1586/ern.10.122>

Muthén, L. K., & Muthén, B. O. (1998-2022). *Mplus User's Guide*. Eighth Edition. Los Angeles, CA: Muthén & Muthén.

Panksepp, J. (2007). Criteria for basic emotions: Is DISGUST a primary “emotion”? *Cognition and Emotion*, 21(8), 1819–1828. <https://doi.org/10.1080/02699930701334302>

Pavot, W., & Diener, E. (2008). The Satisfaction With Life Scale and the emerging construct of life satisfaction. *The Journal of Positive Psychology*, 3(2), 137–152. <https://doi.org/10.1080/17439760701756946>

R Core Team (2022). R: *A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.

Rosseel, Y. (2012). lavaan: An R package for structural equation modeling. *Journal of Statistical Software*, 48, 1–36. <https://doi.org/10.18637/jss.v048.i02>

Rudnev, M. (2022). *Measurement invariance explorer*. MIE R package. <https://github.com/MaksimRudnev/MIE.package>

Rutkowski, L., & Svetina, D. (2017). Measurement invariance in international surveys: Categorical indicators and fit measure performance. *Applied Measurement in Education*, 30(1), 39–51. <https://doi.org/10.1080/08957347.2016.1243540>



## CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

- Ryder, A. G., Yang, J., Zhu, X., Yao, S., Yi, J., Heine, S. J., & Bagby, R. M. (2008). The cultural shaping of depression: somatic symptoms in China, psychological symptoms in North America? *Journal of Abnormal Psychology, 117*(2), 300–313. <https://doi.org/10.1037/0021-843X.117.2.300>
- Saucier, G., Thalmayer, A. G., & Bel-Bahar, T. S. (2014). Human attribute concepts: relative ubiquity across twelve mutually isolated languages. *Journal of Personality and Social Psychology, 107*(1), 199–216. <https://doi.org/10.1037/a0036492>
- Sawicki, A. J., Żemojtel-Piotrowska, M., Balcerowska, J. M., Sawicka, M. J., Piotrowski, J., Sedikides, C., Jonason, P. K., Maltby, J., Adamovic, M., Agada, A. M. D., Ahmed, O., Al-Shawaf, L., Appiah, S. C. Y., Ardi, R., Babakr, Z. H., Bălăţescu, S., Bonato, M., Cowden, R. G., Chobthamkit, P., ... Zand, S. (2022). The fear of COVID-19 scale: Its structure and measurement invariance across 48 countries. *Psychological Assessment, 34*(3), 294–310. <https://doi.org/10.1037/pas0001102>
- Schalinski, I., Schauer, M., & Elbert, T. (2015). The shutdown dissociation scale (Shut-d). *European Journal of Psychotraumatology, 6*, 25652. <https://doi.org/10.3402/ejpt.v6.25652>
- Skalski, S., Uram, P., Dobrakowski, P., & Kwiatkowska, A. (2021). The link between ego-resiliency, social support, SARS-CoV-2 anxiety and trauma effects. Polish adaptation of the Coronavirus Anxiety Scale. *Personality and Individual Differences, 171*, 110540. <https://doi.org/10.1016/j.paid.2020.110540>
- Teachman, B. A., & Gordon, T. (2009). Age differences in anxious responding: Older and calmer, unless the trigger is physical. *Psychology and Aging, 24*(3), 703–714. <https://doi.org/10.1037/a0016813>

## CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

- Van De Schoot, R., Kluytmans, A., Tummers, L., Lugtig, P., Hox, J., & Muthén, B. (2013). Facing off with Scylla and Charybdis: a comparison of scalar, partial, and the novel possibility of approximate measurement invariance. *Frontiers in Psychology, 4*, 770.
- van de Vijver, F. J. R. (2019). Cross-cultural research. In J. Edlund & A. Nichols (Eds.), *Advanced research methods and statistics for the behavioral and social sciences* (pp. 274–286). Cambridge University Press.
- Vinaccia, S., Bahamón, M. J., Trejos-Herrera, A. M., Lee, S. A., Quiceno, J. M., Gómez, C. A., DoLugar, S. V., & Pelaez, E. C. (2022). Validating the Coronavirus Anxiety Scale in a Colombian sample. *Death Studies, 46*(10), 2366–2375.  
<https://doi.org/10.1080/07481187.2021.1944401>
- Voitsidis, P., Dialehti Kerasidou, M., Nikopoulou, A. V., Tsalikidis, P., Parlapani, E., Holeva, V., & Diakoginnis, I. (2021). A systematic review of questionnaires assessing the psychological impact of COVID-19. *Psychiatry Research, 305*, 114183.  
<https://doi.org/10.1016/j.psychres.2021.114183>
- Wall, A. D., & Lee, E. B. (2022). What do anxiety scales really measure? an item content analysis of self-report measures of anxiety. *Journal of Psychopathology and Behavioral Assessment, 44*, 1148–1157. <https://doi.org/10.1007/s10862-022-09973-9>
- Wu, H., & Estabrook, R. (2016). Identification of confirmatory factor analysis models of different levels of invariance for ordered categorical outcomes. *Psychometrika, 81*(4), 1014–1045. <https://doi.org/10.1007/s11336-016-9506-0>
- Yoon, M., & Lai, M. H. (2018). Testing factorial invariance with unbalanced samples. *Structural Equation Modeling, 25*(2), 201–213. <https://doi.org/10.1080/10705511.2017.1387859>

CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

**Table 1.**

*Fit Indices of the Exact Invariance Tests of the CAS*

	CFI	$\Delta$ CFI	TLI	$\Delta$ TLI	RMSEA	$\Delta$ RMSEA	SRMR	$\Delta$ SRMR
Configural	.998		.995		.047		.033	
Full Scalar	.988	.010	.984	.011	.085	.038	.048	.015
Partial Scalar	.996	.002	.983	.002	.056	.009	.043	.010

*Note.* In the Partial Scalar model, threshold of items item 1 (dizziness) and item 3 (tonic immobility) are free.

CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

**Table 2.**

*Exact Measurement Invariance Tests with the Full and Resampled Data (Number of Runs is 500; Number of Groups is 52)*

	$\chi^2$ (SD)	CFI (SD)	TLI (SD)	RMSEA (SD)	SRMR (SD)	N converged replications	NPD matrix (% of replications)
<i>Not resampled, reduced sample of groups, N=52</i>							
Configural	502.1	.997	.995	.049 [.042, .055]	.032	converged	0
Partial	945.4	.994	.993	.057 [.053, .062]	.042	converged	0
Scalar	1631.1	.987	.984	.087 [.082, .091]	.047	converged	0
$\Delta$ Configural and Scalar	1128.9	.010	.011	.038	.015		
$\Delta$ Configural and Partial	443.2	.003	.002	.006	.010		
<i>Resampling without replacement, N per group = 99</i>							
Configural	284.9 (20.1)	.999 (.001)	.998 (.001)	.028 (.014)	.055 (.003)	485	93
Partial	529.0 (27.7)	.996 (.001)	.995 (.001)	.053 (.006)	.070 (.003)	488	93
Scalar	653.7 (32.7)	.992 (.001)	.990 (.002)	.077 (.005)	.070 (.003)	477	93
$\Delta$ Configural and Scalar	368.7 (12.6)	.007 (.000)	.008 (.001)	.049 (.009)	.015 (.000)		
$\Delta$ Configural and Partial	244.0 (7.6)	.003 (.000)	.003 (.000)	.025 (.008)	.015 (.000)		
<i>Resampling with replacement, N rep per group = 500</i>							

CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

	$\chi^2$ (SD)	CFI (SD)	TLI (SD)	RMSEA (SD)	SRMR (SD)	N converged replications	NPD matrix (% of replications)
Configural	928.6 (52.3)	.995 (.000)	.990 (.001)	.072 (.003)	.044 (.002)	500	71
Partial	1831.5 (74.7)	.990 (.001)	.987 (.001)	.083 (.002)	.057 (.002)	500	59
Scalar	2511.2 (92.3)	.984 (.001)	.980 (.001)	.101 (.002)	.058 (.002)	500	77
$\Delta$ Configural and Scalar	1582.5 (40.0)	.011 (.001)	.010 (.000)	.029 (.001)	.014 (.000)		
$\Delta$ Configural and Partial	902.9 (22.4)	.005 (.001)	.003 (.000)	.011 (.001)	.013 (.000)		
<i>Resampling with replacement, N rep per group = 1000</i>							
Configural	1633.2 (72.7)	.995 (.000)	.989 (.001)	.073 (.002)	.041 (.001)	500	42
Partial	3316.0 (101.2)	.989 (.001)	.986 (.001)	.084 (.001)	.053 (.001)	500	56
Scalar	4692.8 (122.8)	.983 (.001)	.979 (.001)	.102 (.001)	.055 (.001)	500	77
$\Delta$ Configural and Scalar	3059.5 (50.1)	.012 (.001)	.010 (.000)	.029 (.001)	.014 (.000)		
$\Delta$ Configural and Partial	1682.7 (28.5)	.006 (.001)	.003 (.000)	.011 (.001)	.012 (.000)		

*Note.* Degrees of freedom for Configural, Partial, and Scalar models are 260, 413, and 413, respectively. NPD - non-positive definite model-implied covariance matrix (e.g., negative variances of observed variables).

CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

**Table 3.**

*The Results of the Fixed-Mode WLS Alignment of CAS*

Parameter	Aligned estimated parameter	R <sup>2</sup>	N nonvariant	List of countries with non-invariant parameters
<i>Thresholds</i>				
Dizziness	1.08	0.76	21	Afghanistan, Armenia, Bangladesh, Bulgaria, Colombia, Cyprus, Georgia, Germany, Indonesia, Iran, Israel, Lebanon, Malaysia, Nepal, Philippines, Poland, Romania, Singapore, Syria, Ukraine, Zambia
Sleep disturbances	0.55	0.82	7	Afghanistan, Brazil, Honduras, Lebanon, Philippines, Portugal, USA
Tonic immobility	1.33	0.74	14	Bosnia and Herzegovina, Brazil, Colombia, Cuba, Ecuador, Georgia, Germany, Honduras, Poland, Romania, Serbia, Singapore, Slovenia, Turkey
Appetite changes	0.88	0.93	3	Brazil, India, Japan
Nausea and abdominal distress	0.81	0.89	4	Kosovo <sup>a</sup> , Romania, Slovakia, Vietnam
<i>Loadings</i>				
Dizziness	0.87	0.59	0	
Sleep disturbances	0.90	0.29	0	
Tonic immobility	0.91	0.52	0	
Appetite changes	0.91	0.61	0	
Nausea and abdominal distress	0.93	0.42	0	

*Note.* <sup>a</sup> Kosovo declared its independence from Serbia in 2008, but there is no consensus on its status as a state. As of December 2022, out of 193 United Nations member states, 117 countries do recognize Kosovo as an independent state, whereas 76 countries, including Serbia, do not.

CROSS-NATIONAL INVARIANCE OF THE CORONAVIRUS ANXIETY SCALE

**Table 4.**

*Pearson Correlations Between the Latent Means (Below the Diagonal) and variances (Above the Diagonal) Estimated by Different Methods, N = 56.*

	1	2	3	4	5	6	7	8	9	10
1. Mean score	-	.22	.28*	.28*	0.33*	.34*	.37**	.22	.17	.35**
2. Configural	NA	-	.77***	.45***	.72***	.65***	.68***	.59***	.54***	.63***
3. Partial	.96***	NA	-	.58***	.94***	.92***	.92***	.82***	.80***	.78***
4. Scalar	.73***	NA	.85***	-	.55***	.49***	.51***	.51***	.53***	.55***
5. Alignment WLS	.98***	NA	.96***	.74***	-	.97***	.98***	.89***	.86***	.87***
6. Alignment MLR	.90***	NA	.93***	.81***	.91***	-	.98***	.91***	.87***	.86***
7. Alignment Bayes	.89***	NA	.92***	.82***	.89***	.93***	-	.90***	.87***	.86***
8. BSEM approximate (configural)	.91***	NA	.93***	.81***	.90***	.97***	.94***	-	.94***	.88***
9. BSEM approximate (scalar)	.94***	NA	.95***	.82***	.92***	.95***	.94***	.97***	-	.85***
10. Alignment BSEM	.93***	NA	.95***	.83***	.92***	.96***	.94***	.98***	.99***	-

*Note.* NA – means are not available in exact configural model because they are fixed to zero for identification purposes. WLS = Weighted Least Squares Mean and Variance adjusted. MLR = Robust Maximum Likelihood. BSEM = Bayesian Structural Equation Modeling.

Fit indices and other details of each model are provided in the OSM, Section D.

\*  $p < .05$ , \*\*  $p < .01$ , \*\*\*  $p < .001$

CORONAVIRUS ANXIETY SCALE

**Table 5.**

*Summary of relations between CAS, FCV-19S and SWLS-3 (excluding Pakistan)*

	Correlations between the CAS and		
	Psychological distress (FCV-19S)	Physiological arousal (FCV-19S)	Life satisfaction (SWLS-3)
<i>Factor scores (estimated with alignment) correlations</i>			
Average individual-level correlation (SD)	.41 (.11)	.44 (.10)	-.10 (.09)
Correlation of means	.42**	.51***	-.15
N	50	50	49
<i>Multiple group CFA (N= 52)<sup>a</sup></i>			
Average individual-level correlation (SD)	.55 (.14)	.51 (.17)	-.13 (.10)
<i>Common alignment (N = 48)</i>			
Average individual-level covariance (SD)	.53 (.14)	.47 (.15)	-.13 (.11)
Correlation of means	.63***	.40**	-.07

<sup>a</sup> Latent means were not available for FCV and SWLS-3 scales since the established models were metric, and thus the means were fixed to 0 in all the groups. Fit indices for the Multi-group CFA are:  $\chi^2 = 6868.3$ ,  $df = 4620$ , CFI = .973, TLI = .968, RMSEA = .035, SRMR = .055.

\*\*  $p < .01$ , \*\*\*  $p < .001$