

THE MINIMAL PROJECTIVE BUNDLE DIMENSION AND TORIC 2-FANO MANIFOLDS

CAROLINA ARAUJO, ROYA BEHESHTI, ANA-MARIA CASTRAVET, KELLY JABBUSCH,
SVETLANA MAKAROVA, ENRICA MAZZON, AND NIVEDITA VISWANATHAN,
WITH AN APPENDIX BY WILL REYNOLDS

ABSTRACT. Motivated by the problem of classifying toric 2-Fano manifolds, we introduce a new invariant for smooth projective toric varieties, the minimal projective bundle dimension. This invariant $m(X) \in \{1, \dots, \dim(X)\}$ captures the minimal degree of a dominating family of rational curves on X or, equivalently, the minimal length of a centered primitive relation for the fan of X . We classify smooth projective toric varieties with $m(X) \geq \dim(X) - 2$, and show that projective spaces are the only 2-Fano manifolds among smooth projective toric varieties with $m(X) \in \{1, \dim(X) - 2, \dim(X) - 1, \dim(X)\}$.

CONTENTS

1. Introduction	1
2. Primitive collections	5
2.1. Notation and background	5
2.2. The minimal \mathbb{P} -dimension	8
2.3. Some properties of primitive collections	11
2.4. Primitive collections on toric Fano manifolds	12
3. Toric Fano manifolds with $m(X) = 1$	13
4. Proof of Theorem 1.3	17
4.1. First case: $\{x, y, z\}$ is a primitive collection	18
4.2. Second case: none of $\{x, y, z\}$ form a primitive collection	24
5. Toric Fano manifolds with $m(X) = n - 2$	25
Appendix A. Code for computing primitive collections	27
References	29

1. INTRODUCTION

Fano varieties are projective varieties with positive first Chern class. Over the complex numbers, this condition is equivalent to the existence of a metric with positive Ricci curvature. Basic examples of Fano varieties include projective spaces and Grassmannians. The positivity condition has further geometric implications, e.g., Fano varieties over the complex numbers are *simply connected*. This has an analogue on the algebro-geometric side: any Fano variety is covered by rational curves [Mor79], and is in fact *rationally connected* [KMM92; Cam92], i.e., there are rational curves connecting any two of its points. In a series of papers, de Jong and Starr introduce and investigate possible candidates for the notion of *higher rational connectedness* [dHS11; dS07; dS06b; dS06c; Sta06], inspired

by the natural analogue in topology. In particular, in [dS06b] they define *2-Fano* manifolds. A smooth projective variety X is 2-Fano if it is Fano and its second Chern character $\text{ch}_2(T_X) = \frac{1}{2}c_1(T_X)^2 - c_2(T_X)$ is positive, i.e., $\text{ch}_2(T_X) \cdot S > 0$ for every surface S in X . In a similar way, one can define k -Fano varieties for any $k \geq 2$, and aim at their classification. For instance, \mathbb{P}^n is n -Fano, and it is conjectured that it is the only n -dimensional n -Fano manifold. The geometry of higher Fano manifolds has been fairly investigated, and in several special cases they are shown to enjoy the expected nice properties. For instance, 2-Fano manifolds satisfying some mild assumptions are covered by rational surfaces [dS07], and similar results hold for higher Fano manifolds [Suz21], [Nag19]. There is a classification of 2-Fano manifolds of high index [AC13] and, more recently, a classification of homogeneous 2-Fano manifolds [Ara+22]. On the other hand, very few examples of higher Fano manifolds are known. Quite strikingly, all known examples of 2-Fano manifolds have Picard rank 1 and relatively large index.

It is natural for algebraic geometers to turn to the pool of toric varieties when looking for intuition or examples. It is well known that projective spaces are the only projective toric manifolds with Picard rank 1. Thus, a classification of toric 2-Fano manifolds could either provide the first examples of 2-Fano manifolds with higher Picard rank, or it could be an evidence that every 2-Fano manifold has Picard rank 1. Geometric properties of a toric variety can often be checked in the combinatorics of the associated fan. This bridge has been exploited in search of new examples of toric 2-Fano manifolds [Nob11], [Nob12], [Sat12], [Sat16], [SS20], [SSS21], [Shr20]. Despite the efforts, a complete (computer aided) classification is only known up to dimension 8 [Nob11], [SSS21], and projective spaces remain the only known examples of toric 2-Fano manifolds. The sparsity of higher Fano manifolds leads to the following conjecture.

Conjecture 1.1. ([SSS21, Conjecture 4.3]) The only toric 2-Fano manifolds are projective spaces.

In this paper, we propose a new strategy to approach Conjecture 1.1. We follow the philosophy introduced in [AC12], namely, to investigate 2-Fano manifolds by studying their *minimal dominating families of rational curves*. By [CFH14], minimal dominating families of rational curves on a smooth projective toric variety X correspond to *primitive relations* of the form

$$(1) \quad x_0 + \cdots + x_m = 0,$$

satisfied by some of the primitive integral generators x_i of the corresponding fan. These primitive relations are called *centered of order $m + 1$* . By [CFH14], a centered primitive relation of order $m + 1$ yields a \mathbb{P}^m -bundle structure $X^\circ \rightarrow T$ on a dense open subset X° of X . If $\dim(T) \geq 1$, and the complement $X \setminus X^\circ$ has codimension at least 2 in X , then one can construct a complete surface $S \subset X^\circ$ such that $\text{ch}_2(T_X) \cdot S \leq 0$, showing that X is not 2-Fano. So our basic strategy consists of trying to describe, in a rather explicit way, a suitable birational map $\varphi : X \dashrightarrow Y$ transforming X into a projective toric variety Y admitting a \mathbb{P}^m -bundle structure on a big open subset. We then hope to be able to compare the second Chern characters $\text{ch}_2(T_X)$ and $\text{ch}_2(T_Y)$ to show that X is not 2-Fano, except if $X = \mathbb{P}^m$ and φ is the identity.

To follow this strategy, we introduce a new invariant of a smooth projective toric variety X , the *minimal projective bundle dimension* of X , *minimal \mathbb{P} -dimension* in short, which is

of independent interest (Definition 2.10):

$$m(X) = \min \{ m \in \mathbb{Z}_{>0} \mid \text{there is a relation as in (1)} \} \in \{1, \dots, \dim X\}.$$

Building on [Bat99; Sat00; Øbr07] and data provided by Øbro, the Graded Ring Database [BK] collects the combinatorial data of toric Fano manifolds of dimension up to 6. This data is also accessible via NormalToricVarieties package ([Smi]) in Macaulay2 [GS]. For such manifolds, the code in Appendix A by Will Reynolds computes their primitive collections, and thus allows to extract their minimal \mathbb{P} -dimension.

$\dim(X)$	# Fanos	$\#(m=1)$	$\#(m=2)$	$\#(m=3)$	$\#(m=4)$	$\#(m=5)$	$\#(m=6)$
4	124	107	15	1	1		
5	866	744	112	8	1	1	
6	7622	6333	1174	105	8	1	1

TABLE 1. The number of toric Fano manifolds of dimension 4, 5, 6 according to their minimal \mathbb{P} -dimension.

When $m(X) = 1$, Casagrande constructs in [Cas03b] a sequence of blowdowns and flips from X to a toric variety admitting a global \mathbb{P}^1 -bundle structure. This allows us to make the basic strategy work, yielding the following result.

Theorem 1.2. Let X be a smooth toric Fano variety with $m(X) = 1$. Then X is not 2-Fano.

Table 1 suggests that this result covers “most” toric varieties, and not just the fringe cases.

Next we turn our attention to toric Fano manifolds X with large values of $m(X)$. Projective spaces are the only smooth projective toric varieties admitting a centered primitive relation of order $\dim(X) + 1$. In [CFH14, Proposition 3.8], Chen, Fu and Hwang classify toric Fano manifolds admitting a centered primitive relation of order $\dim(X)$. There are three such varieties, and two of them also admit a centered primitive relation of order 2. As a consequence, the only n -dimensional toric Fano manifold X with $m(X) = n - 1$ is the blowup of \mathbb{P}^n along a linear \mathbb{P}^{n-2} . In [BW23], Beheshti and Wormleighton investigate smooth projective toric varieties admitting a centered primitive relation of order $\dim(X) - 1$, showing that they have Picard rank $\rho(X) \leq 5$. Most of these varieties also admit centered primitive relations of order 2 or 3, and we prove the following bound for the remaining ones. Theorem 1.4 shows that this bound is sharp.

Theorem 1.3. Let X be a smooth toric Fano variety with $\dim(X) = n \geq 6$ and $m(X) \geq 3$. If X has a centered primitive relation of order $n - 1$,

$$x_0 + x_1 + \dots + x_{n-2} = 0,$$

then $\rho(X) \leq 3$. Moreover, $m(X) = n - 2$ and the above relation is the only centered primitive relation of X .

Using Theorem 1.3 and Batyrev’s description of smooth projective toric varieties with Picard rank 3, we are able to classify n -dimensional smooth toric Fano varieties with $m(X) = n - 2$. There are eight distinct isomorphism classes when $n \geq 6$, which can be

explicitly described. The following statement summarizes the classification of toric Fano manifolds with $m(X) \geq \dim(X) - 2$.

Theorem 1.4. We have the following classification of smooth toric Fano varieties with $m(X) \geq \dim(X) - 2$.

- (1) The only n -dimensional smooth toric Fano variety X with $m(X) = n$ is \mathbb{P}^n .
- (2) For $n \geq 3$, the only n -dimensional smooth toric Fano variety X with $m(X) = n - 1$ is the blowup of \mathbb{P}^n along a linear \mathbb{P}^{n-2} .
- (3) For $n \geq 6$, there are eight distinct isomorphism classes of n -dimensional smooth toric Fano varieties X with $m(X) = n - 2$. Namely:
 - (a) $X = \mathbb{P}_S(\mathcal{E})$ is a \mathbb{P}^{n-2} -bundle over a toric surface S , where (S, \mathcal{E}) is one of the following:
 - $S = \mathbb{P}^2$ and $\mathcal{E} = \mathcal{O}_{\mathbb{P}^2}(1) \oplus \mathcal{O}_{\mathbb{P}^2}^{\oplus n-2}$,
 - $S = \mathbb{P}^2$ and $\mathcal{E} = \mathcal{O}_{\mathbb{P}^2}(1) \oplus \mathcal{O}_{\mathbb{P}^2}(1) \oplus \mathcal{O}_{\mathbb{P}^2}^{\oplus n-3}$,
 - $S = \mathbb{P}^2$ and $\mathcal{E} = \mathcal{O}_{\mathbb{P}^2}(2) \oplus \mathcal{O}_{\mathbb{P}^2}^{\oplus n-2}$,
 - $S = \mathbb{P}^1 \times \mathbb{P}^1$ and $\mathcal{E} = \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(1, 1) \oplus \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}^{\oplus n-2}$,
 - $S = \mathbb{P}^1 \times \mathbb{P}^1$ and $\mathcal{E} = \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(1, 0) \oplus \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(0, 1) \oplus \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}^{\oplus n-3}$,
 - $S = \mathbb{F}_1$ and $\mathcal{E} = \mathcal{O}_{\mathbb{F}_1}(e + f) \oplus \mathcal{O}_{\mathbb{F}_1}^{\oplus n-2}$, where $e \subset \mathbb{F}_1$ is the -1 -curve, and $f \subset \mathbb{F}_1$ is a fiber of $\mathbb{F}_1 \rightarrow \mathbb{P}^1$.

In the first three cases, $\rho(X) = 2$, while in the latter three cases, $\rho(X) = 3$.

- (b) Let $Y \simeq \mathbb{P}_{\mathbb{P}^2}(\mathcal{O}_{\mathbb{P}^2}(1) \oplus \mathcal{O}_{\mathbb{P}^2}^{\oplus n-2})$ be the blowup of \mathbb{P}^n along a linear subspace $L = \mathbb{P}^{n-3}$, and denote by $E \subset Y$ the exceptional divisor. Then X is the blowup of Y along a codimension 2 center $Z \subset Y$, where:
 - Z is the intersection of E with the strict transform of a hyperplane of \mathbb{P}^n containing the linear subspace L , or
 - Z is the intersection of the strict transforms of two hyperplanes of \mathbb{P}^n , one containing the linear subspace L , and the other one not containing it.

In both cases, $\rho(X) = 3$.

Corollary 1.5. The projective space \mathbb{P}^n is the only smooth n -dimensional toric 2-Fano variety with $m(X) \in \{1, n - 2, n - 1, n\}$.

This paper is organized as follows. In Section 2, we review some results from toric geometry and fix notation. In particular, we discuss centered primitive relations on smooth projective toric varieties, describing explicitly their open subsets admitting a projective space bundle structure (Proposition 2.14). In Section 3, we study smooth toric Fano varieties with $m(X) = 1$, and prove Theorem 1.2. In Section 4, we investigate smooth projective n -dimensional toric varieties admitting a centered primitive relation of order $n - 1$, and prove Theorem 1.3. In Section 5, we use this result, together with Batyrev's description of smooth projective toric varieties with Picard rank 3, to prove Theorem 1.4.

Acknowledgements. This collaboration started as a working group at the ICERM “Women in Algebraic Geometry Collaborative Research Workshop” in July 2020. A great part of this work was developed during the follow-up “Collaborate@ICERM” meeting in May 2022. We thank ICERM for the financial support and the great working conditions provided to us during our visit. We are very grateful to Cinzia Casagrande and Milena

Hering for many enlightening discussions and explanations about toric varieties, and to Will Reynolds for running his code and providing us with precious data about primitive relations of toric Fano manifolds. We thank the anonymous referees for their careful reading of the paper and suggestions about terminology and exposition.

Carolina Araujo was partially supported by CAPES/COFECUB, CNPq and FAPERJ Research Fellowships. Roya Beheshti was supported by NSF grant DMS-2101935. Ana-Maria Castravet was partially supported by the ANR 20-CE40-0023 grant *FanoHK* and the ANR-22-CE40-0009-01 grant *FRACASSO*. Enrica Mazzon was supported by the collaborative research center SFB 1085 *Higher Invariants - Interactions between Arithmetic Geometry and Global Analysis* funded by the Deutsche Forschungsgemeinschaft. Nivedita Viswanathan was supported by the EPSRC New Horizons Grant No.EP/V048619/1.

2. PRIMITIVE COLLECTIONS

2.1. Notation and background. A toric variety is a normal complex variety X that contains a torus $T = (\mathbb{C}^*)^n$ as a dense open subset, together with an action of T on X that extends the natural action of T on itself. There is a one-to-one correspondence between n -dimensional toric varieties and *fans* in \mathbb{Q}^n . Let N be a free abelian group of rank n , and consider the vector space $N_{\mathbb{Q}} = N \otimes_{\mathbb{Z}} \mathbb{Q}$. A *fan* in $N_{\mathbb{Q}}$ is a nonempty finite collection Σ of strongly convex polyhedral cones in $N_{\mathbb{Q}}$ such that every face of a cone in Σ is also a cone in Σ , and the intersection of two cones in Σ is a face of each. We write $\delta \prec \tau$ to express that δ is a face of τ . One-dimensional cones in Σ are called rays, and each ray is generated by a primitive vector in N . The set of primitive vectors of N generating rays of Σ is denoted by $G(\Sigma)$. We will write a cone $\tau \in \Sigma$ in terms of its primitive generators, $\tau = \langle v_1, \dots, v_l \rangle$, saying that the v_i 's generate τ , and setting $G(\tau) := \{v_1, \dots, v_l\} \subseteq G(\Sigma)$.

We denote by X_{Σ} the toric variety corresponding to a fan Σ . Conversely, given a toric variety X , we denote by Σ_X the fan associated to X . There is a one-to-one inclusion-reversing correspondence between cones in Σ and T -orbit closures in X_{Σ} . Given a cone $\tau \in \Sigma$, we write $V(\tau) \subset X_{\Sigma}$ for the corresponding T -orbit closure, or $V(v_1, \dots, v_l)$ when $G(\tau) = \{v_1, \dots, v_l\}$. Note that $\dim(\tau) = \text{codim}_{X_{\Sigma}} V(\tau)$. We refer to [Ful93] and [CLS11] for the background on toric varieties.

In this paper, we are mostly interested in *smooth* and *proper* toric varieties. The smoothness conditions translates into the fan Σ being *regular* (also called *unimodular*), i.e., for each cone $\tau \in \Sigma$, the set of generators $G(\tau)$ is part of a basis of N ([CLS11, Definition 1.2.16]). The properness condition translates into the fan Σ being *complete*, i.e., its support being the whole $N_{\mathbb{Q}}$. In what follows, smooth and proper toric varieties will be simply called *toric manifolds*. We would like to classify toric 2-Fano manifolds. Given a toric manifold X , there is an exact sequence ([CLS11, Theorem 8.1.1]):

$$0 \rightarrow \Omega_X^1 \rightarrow N^{\vee} \otimes_{\mathbb{Z}} \mathcal{O}_X \rightarrow \bigoplus_{v \in G(\Sigma_X)} \mathcal{O}_{V(v)} \rightarrow 0,$$

from which one easily computes:

$$c_1(X) = \sum_{v \in G(\Sigma_X)} V(v) \quad \text{and} \quad \text{ch}_2(X) = \frac{1}{2} \sum_{v \in G(\Sigma_X)} V(v)^2.$$

Definition 2.1. ([Bat91, Definition 2.6]) Let Σ be a regular complete fan in $N_{\mathbb{Q}}$. A *primitive collection* $P \subseteq G(\Sigma)$ is a nonempty set of primitive vectors of N that does not

generate a cone of Σ , but such that any proper subset of P does. Equivalently, $P = \{v_1, \dots, v_r\} \subseteq G(\Sigma)$ is a primitive collection if and only if

$$\langle v_1, \dots, v_r \rangle \notin \Sigma \quad \text{and} \quad \langle v_1, \dots, \check{v}_i, \dots, v_r \rangle \in \Sigma$$

for any $i = 1, \dots, r$. We denote by $\text{PC}(\Sigma)$ the set of primitive collections of Σ . For a toric manifold X , we will talk about primitive collections of X and write $\text{PC}(X)$, meaning $\text{PC}(\Sigma_X)$.

Definition 2.2. Let Σ be a regular complete fan in $N_{\mathbb{Q}}$. Given a primitive collection $P = \{v_1, \dots, v_r\} \in \text{PC}(\Sigma)$, let $\sigma(P) = \langle w_1, \dots, w_s \rangle$ be the minimal cone of Σ such that $v_1 + \dots + v_r \in \sigma(P)$. Then there is a relation

$$r(P): v_1 + \dots + v_r = \mu_1 w_1 + \dots + \mu_s w_s,$$

where $\mu_j \in \mathbb{Z}_{\geq 0}$ for $j = 1, \dots, s$. We call $r(P)$ the *primitive relation* associated to P . We define the *order* of P as $\text{ord}(P) = |P| = r$, while the *degree* of P as $\deg(P) = r - \sum_{j=1}^s \mu_j$.

By [Bat91, Proposition 3.1], for any primitive collection P , we have $P \cap \sigma(P) = \emptyset$. In particular, $\{v_1, \dots, v_r\} \cap \{w_1, \dots, w_s\} = \emptyset$.

Definition 2.3. Let Σ be a regular complete fan in $N_{\mathbb{Q}}$. A primitive collection $P = \{x_0, \dots, x_k\}$ of Σ is called *centered* if $\sigma(P) = \{0\}$, i.e.

$$r(P): x_0 + \dots + x_k = 0.$$

Lemma 2.4. Let Σ be a regular complete fan in $N_{\mathbb{Q}}$, and let P, Q be two distinct centered primitive collections. Then $P \cap Q = \emptyset$.

Proof. Write $r(P): x_0 + \dots + x_k = 0$ and $r(Q): y_0 + \dots + y_l = 0$. Assume that $P \cap Q \neq \emptyset$, then without loss of generality we may assume that $x_0 = y_0$. But then subtracting this vector from both relations, we get

$$x_1 + \dots + x_k = y_1 + \dots + y_l,$$

which shows that interiors of two distinct cones intersect. This is a contradiction. \square

Lemma 2.5. Let Σ be a regular complete fan in $N_{\mathbb{Q}}$, and let P, Q be two distinct centered primitive collections. Then $\text{Span } P \cap \text{Span } Q = \{0\}$, in particular $|P| + |Q| - 2 \leq \dim N_{\mathbb{Q}}$.

Proof. Write $r(P): x_0 + \dots + x_k = 0$ and $r(Q): y_0 + \dots + y_l = 0$. Take any vector $v \in \text{Span } P \cap \text{Span } Q$, so we can write it as

$$v = \sum a_i x_i = \sum b_j y_j.$$

By possibly adding $r(P)$ and $r(Q)$ to the sums, we can get that all $a_i, b_j \geq 0$, and up to relabelling the a_i, b_j , we can assume $a_0 = b_0 = 0$. But this shows that v is in the intersection of two cones, $\langle x_1, \dots, x_k \rangle \cap \langle y_1, \dots, y_l \rangle$, and the sets of generators are disjoint by Lemma 2.4, so $\langle x_1, \dots, x_k \rangle \cap \langle y_1, \dots, y_l \rangle = \{0\}$ and $v = 0$.

The last claim follows from considering the dimensions of $\text{Span } P$ and $\text{Span } Q$. \square

Let $A_1(X_{\Sigma})$ be the group of algebraic 1-cycles on X_{Σ} modulo numerical equivalence, and set $\mathcal{N}_1(X_{\Sigma}) = A_1(X_{\Sigma}) \otimes_{\mathbb{Z}} \mathbb{Q}$. The Mori cone $\text{NE}(X_{\Sigma}) \subset \mathcal{N}_1(X_{\Sigma})$ is the cone generated

by the classes of effective curves. A primitive integral class generating an extremal ray of $\text{NE}(X_\Sigma)$ is called an *extremal class*. There is an exact sequence:

$$\begin{aligned} 0 \longrightarrow A_1(X_\Sigma) &\longrightarrow \mathbb{Z}^{G(\Sigma)} \longrightarrow N \longrightarrow 0, \\ [C] &\longrightarrow (C \cdot V(v))_{v \in G(\Sigma)} \\ &\quad (\nu_v)_{v \in G(\Sigma)} \longrightarrow \sum_{v \in G(\Sigma)} \nu_v v. \end{aligned}$$

Thus the elements of $A_1(X_\Sigma)$ are identified with integral relations between the elements of $G(\Sigma)$. If the class $[C]$ corresponds to the relation $\sum_v \nu_v v = 0$, then we have $-K_{X_\Sigma} \cdot C = \sum_v \nu_v$.

Proposition 2.6. ([Cas03a, Lemma 1.4]) Let Σ be a regular complete fan in $N_\mathbb{Q}$. A relation

$$\alpha_1 x_1 + \cdots + \alpha_l x_l - \beta_1 y_1 - \cdots - \beta_m y_m = 0,$$

with $\alpha_i, \beta_j \in \mathbb{Z}_{>0}$, defines an effective class in $\mathcal{N}_1(X_\Sigma)$ provided that $\langle y_1, \dots, y_m \rangle$ is a cone of Σ .

We will usually write the above relation as

$$\alpha_1 x_1 + \cdots + \alpha_l x_l = \beta_1 y_1 + \cdots + \beta_m y_m.$$

It follows that primitive relations correspond to effective curve classes. By abuse of notation, we will identify a primitive relation $r(P)$ with the corresponding curve class. Note that $\deg(P) = -K_{X_\Sigma} \cdot r(P)$. In the projective case we have the following description of $\text{NE}(X_\Sigma)$.

Proposition 2.7. ([Bat91, Theorem 2.15]) Let Σ be a regular complete fan in $N_\mathbb{Q}$, and assume that X_Σ is projective. Then the Mori cone is generated by primitive relations:

$$\text{NE}(X_\Sigma) = \sum_{P \in \text{PC}(X_\Sigma)} \mathbb{Q}_{\geq 0} r(P).$$

Curve classes corresponding to primitive relations (and in particular centered primitive relations) are not necessarily extremal. When this occurs, the following result holds.

Proposition 2.8. ([Rei83, Theorem 2.4]) Let Σ be a regular complete fan in $N_\mathbb{Q}$, and assume that X_Σ is projective. Let γ be an extremal class in $\text{NE}(X_\Sigma)$ whose corresponding primitive relation is

$$r(P): v_1 + \cdots + v_r = \mu_1 w_1 + \cdots + \mu_s w_s.$$

Let $\tau = \langle z_1, \dots, z_l \rangle$ be a cone of Σ such that $G(\tau) \cap P = G(\tau) \cap G(\sigma(P)) = \emptyset$, and such that $\langle \sigma(P), \tau \rangle = \langle w_1, \dots, w_s, z_1, \dots, z_l \rangle$ is a cone of Σ . Then, for each $i = 1, \dots, r$,

$$\langle P \setminus \{v_i\}, \sigma(P), \tau \rangle = \langle v_1, \dots, \check{v}_i, \dots, v_r, w_1, \dots, w_s, z_1, \dots, z_l \rangle$$

is also a cone of Σ .

2.2. The minimal \mathbb{P} -dimension. Let X be a toric manifold with regular complete fan Σ_X in $N_{\mathbb{Q}}$. In this section, we discuss centered primitive collections, introduced in Definition 2.3.

Proposition 2.9. ([Bat91, Proposition 3.2]) If X is projective, then Σ_X has a centered primitive collection of order $k + 1$

$$(2) \quad r(P): x_0 + \cdots + x_k = 0$$

for some $k \in \{1, \dots, \dim(X)\}$.

Definition 2.10. For a projective toric manifold X , we define the *minimal \mathbb{P} -dimension* as

$$m(X) := \min \left\{ m \in \mathbb{Z}_{>0} \mid \begin{array}{c} \Sigma_X \text{ has a centered} \\ \text{primitive collection of order } m + 1 \end{array} \right\}.$$

The next remark explains the terminology of Definition 2.10 and highlights the significance of studying centered primitive collections.

Remark 2.11. In [CFH14], Chen, Fu and Hwang provide a new geometric proof of Proposition 2.9 by relating centered primitive collections to *minimal dominating families of rational curves*. We review some aspects of the theory of rational curves on varieties and refer to [Kol96] for details. Given a smooth and proper uniruled variety X , there is a scheme $\text{RatCurves}^n(X)$ parametrizing rational curves on X . A *dominating family of rational curves* on X is an irreducible component of $\text{RatCurves}^n(X)$ parametrizing rational curves that sweep out a dense open subset of X . A dominating family of rational curves H is said to be *minimal* if, for a general point $x \in X$, the subvariety of H parametrizing curves through x is proper. When X is projective, there always exists a minimal dominating family of rational curves on X . For instance, one can take H to be a dominating family of rational curves on X having minimal degree with respect to some fixed ample line bundle on X . The minimal anti-canonical degree l_X of a minimal dominating family of rational curves is a natural invariant of a Fano variety X . In [CD15], Casagrande and Druel investigate n -dimensional Fano varieties admitting a minimal dominating family of rational curves of anti-canonical degree n , and classify the cases when $l_X = n$. Currently, there are no general results when the anti-canonical degree is $n - 1$.

When $X = X_{\Sigma}$ is a toric variety, there is a one-to-one correspondence between minimal dominating families of rational curves H on X and centered primitive collections of Σ ([CFH14, Proposition 3.2]). Moreover, if the centered primitive collection has order $k + 1$ as in Equation (2) above, then there is a dense T -invariant open subset U of X and a \mathbb{P}^k -bundle $\pi: U \rightarrow W$ such that the general curve parametrized by H is a line on a general fiber π ([CFH14, Corollary 2.6]).

It follows from this discussion that the minimal \mathbb{P} -dimension $m(X) = l_X - 1$ is the smallest integer k such that X admits a generic \mathbb{P}^k -bundle structure. We have

$$m(X) \in \{1, \dots, n = \dim(X)\},$$

and $m(X) = n$ if and only if $X \simeq \mathbb{P}^n$. By [CFH14, Proposition 3.8], there are three toric Fano manifolds admitting a centered primitive relation of order $n = \dim(X)$, namely: $\mathbb{P}^{n-1} \times \mathbb{P}^1$, the blowup of $\mathbb{P}^{n-1} \times \mathbb{P}^1$ along a linear \mathbb{P}^{n-2} , and the blowup of \mathbb{P}^n along a linear \mathbb{P}^{n-2} . The first two varieties also admit a generic \mathbb{P}^1 -bundle structure. As a consequence, the only n -dimensional toric Fano manifold X with $m(X) = n - 1$ is the blowup of \mathbb{P}^n

along a linear \mathbb{P}^{n-2} . In Section 5, we shall classify n -dimensional toric Fano manifolds X with $m(X) = n - 2$.

Let $P \in \text{PC}(X)$ be a centered primitive collection of order $k + 1$. As explained in Remark 2.11, P induces a \mathbb{P}^k -bundle structure on a dense T -invariant open subset U of X . In [CFH14, Corollary 2.6], the T -invariant open subset U was taken as small as possible, namely, $U \cong \mathbb{P}^k \times (\mathbb{C}^*)^{n-k}$. For our purposes, we want to take U as big as possible. So our next goal is to describe explicitly the biggest T -invariant open subset of X on which P induces a \mathbb{P}^k -bundle structure.

Notation 2.12. Let $P = \{x_0, \dots, x_k\} \in \text{PC}(X)$ be a centered primitive collection. Denote by \mathcal{E}_P the set of cones $\sigma = \langle v_1, \dots, v_r \rangle \in \Sigma_X$ such that $P \cap G(\sigma) = \emptyset$, and $\{v_1, \dots, v_r, x_{j_1}, \dots, x_{j_s}\} \in \text{PC}(X)$ for some $s \geq 1$, i.e.,

$$\mathcal{E}_P := \{\sigma \in \Sigma_X \mid P \cap G(\sigma) = \emptyset \text{ and } \exists P' \subsetneq P \text{ such that } P' \cup G(\sigma) \in \text{PC}(X)\}.$$

We write

$$V(\mathcal{E}_P) := \bigcup_{\sigma \in \mathcal{E}_P} V(\sigma) \subset X.$$

Example 2.13. We give an example that illustrates how $V(\mathcal{E}_P)$ arises from non-extremality of a centered primitive collection P . The toric Fano 6-fold X corresponding to the Macaulay ID (6, 942) in the package *NormalToricVarieties* has a fan generated by nine vectors v_1, \dots, v_9 , and admits a unique centered primitive collection $P = \{v_4, v_6, v_7, v_8\}$. All other primitive relations are

$$\begin{aligned} r_1: \quad & v_1 + v_2 + v_3 + v_9 = v_4 + v_6 + v_8, \\ q: \quad & v_1 + v_2 + v_5 + v_9 = v_8, \\ r_2: \quad & v_3 + v_7 + v_8 = v_5, \\ r_3: \quad & v_4 + v_5 + v_6 = v_3. \end{aligned}$$

In this case, we have $\mathcal{E}_P = \{\langle v_5 \rangle, \langle v_3 \rangle\} = \{\sigma(r_2), \sigma(r_3)\}$ and $V(\mathcal{E}_P)$ is the union of the two corresponding toric divisors. Note that $r(P) = r_2 + r_3$. For completeness of the example, one can also note that $q = r_1 + r_3$, and that r_1, r_2, r_3 are the three extremal primitive relations that generate the Mori cone $\text{NE}(X)$.

Proposition 2.14. Let $P = \{x_0, \dots, x_k\} \in \text{PC}(X)$ be a centered primitive collection, and let $V(\mathcal{E}_P)$ be as in Notation 2.12. Then the open subset $U = X \setminus V(\mathcal{E}_P)$ admits a \mathbb{P}^k -bundle structure over a smooth toric variety.

In order to prove Proposition 2.14, we first prove two auxiliary lemmas.

Lemma 2.15. Let $P = \{x_0, \dots, x_k\} \in \text{PC}(X)$ be a centered primitive collection, let $V(\mathcal{E}_P)$ be as in Notation 2.12, and set $U = X \setminus V(\mathcal{E}_P)$. Then the fan Σ_U of U consists of all cones of Σ_X of the form

$$(3) \quad \tau' = \langle \tau, x_{j_1}, \dots, x_{j_m} \rangle,$$

where $0 \leq m \leq k$, and $\tau \in \Sigma_X$ is such that $\langle \tau, P \setminus \{x_i\} \rangle \in \Sigma$ for every $i \in \{0, \dots, k\}$. (When $m = 0$, Equation (3) means that $\tau' = \tau$.)

Proof. Recall that a cone $\sigma \in \Sigma_X$ corresponds to a T -orbit, which is dense and open in $V(\sigma)$. Hence, a cone $\sigma \in \Sigma_X$ is in Σ_U if and only if the corresponding orbit does not intersect $V(\mathcal{E}_P)$, which is equivalent to saying that $V(\sigma) \not\subseteq V(\mathcal{E}_P)$. It is immediate that the cones of the form (3) define a fan $\Sigma' \subset \Sigma_X$ in $N_{\mathbb{Q}}$, and $X_{\Sigma'}$ is a dense open subset of X . We now prove that the toric variety $X_{\Sigma'}$ coincides with U by showing that a cone $\sigma \in \Sigma_X$ is of the form (3) if and only if $V(\sigma) \not\subseteq V(\mathcal{E}_P)$.

Consider $\sigma \in \Sigma_X \setminus \Sigma'$, which means that $\langle G(\sigma) \cup P \setminus \{x_i\} \rangle \notin \Sigma_X$ for some i . Then the set $G(\sigma) \cup P \setminus \{x_i\}$ contains a primitive collection S , so the cone $\tau := \langle S \setminus P \rangle$ is in \mathcal{E}_P . But notice that $\tau \prec \sigma$, so $V(\sigma) \subseteq V(\tau) \subseteq V(\mathcal{E}_P)$ and hence σ is not in Σ_U . Conversely, if $\sigma \in \Sigma_X \setminus \Sigma_U$, then $V(\sigma) \subseteq V(\mathcal{E}_P)$, hence there exists $\tau \in \mathcal{E}_P$ such that $V(\sigma) \subseteq V(\tau)$ and $G(\tau) \cup P' \in \text{PC}(X)$ for some $P' \subset P$. Since $G(\tau) \subseteq G(\sigma)$, we conclude that $\langle G(\sigma) \cup P' \rangle \notin \Sigma_X$, i.e., $\sigma \notin \Sigma'$. \square

Consider the sequences

$$\begin{aligned} 0 &\longrightarrow N_P := \ker(\phi) \longrightarrow N \xrightarrow{\phi} \overline{N} = N/\mathbb{Z}\langle x_0, \dots, x_k \rangle \longrightarrow 0, \\ 0 &\longrightarrow (N_P)_{\mathbb{Q}} \longrightarrow N_{\mathbb{Q}} \xrightarrow{\phi_{\mathbb{Q}}} \overline{N}_{\mathbb{Q}} \longrightarrow 0, \\ \Sigma_0 &\longrightarrow \Sigma_U \longrightarrow \overline{\Sigma}_U, \end{aligned}$$

where ϕ is the quotient map, the fan Σ_0 of $(N_P)_{\mathbb{Q}} \simeq \mathbb{Q}^{k+1}$ is the subfan of Σ_U of cones of the form (3) with $\tau = \{0\}$ (in particular note that $X_{\Sigma_0} \simeq \mathbb{P}^k$), and $\overline{\Sigma}_U = \{\phi_{\mathbb{Q}}(\sigma) \mid \sigma \in \Sigma_U\}$.

Lemma 2.16. Let the notation be as above. Then $\overline{\Sigma}_U$ is a toric fan, and the linear map $\phi_{\mathbb{Q}}$ is compatible with the fans Σ_U and $\overline{\Sigma}_U$.

Proof. The cones of $\overline{\Sigma}_U$ are exactly $\phi_{\mathbb{Q}}(\tau)$ for $\tau \in \Sigma_U$ such that $G(\tau) \cap P = \emptyset$, so for simplicity we only consider these τ .

- It is immediate that the cones of $\overline{\Sigma}_U$ are rational polyhedral, and that the faces of $\phi_{\mathbb{Q}}(\tau)$ are $\phi_{\mathbb{Q}}(\delta)$, for all subcones $\delta \prec \tau$.
- We need to show that the cone $\phi_{\mathbb{Q}}(\tau)$ is strongly convex, i.e., if $y \in \phi_{\mathbb{Q}}(\tau)$ and $-y \in \phi_{\mathbb{Q}}(\tau)$, then $y = 0$. This follows automatically from the fact that the images of the generators $\phi_{\mathbb{Q}}(G(\tau)) = \{\bar{v}_1, \dots, \bar{v}_r\}$ are linearly independent, which we prove by contradiction. If they are linearly dependent, then there exist $a_1, \dots, a_r \in \mathbb{Q}$, not all 0, such that $\sum_{i=1}^r a_i \bar{v}_i = 0$ in $\overline{N}_{\mathbb{Q}}$. This implies that there exist $b_j \in \mathbb{Q}$ for $j = 1, \dots, k$ such that $\sum_{i=1}^r a_i v_i = \sum_{j=1}^k b_j x_j$, which is a contradiction since $\langle G(\tau) \cup P \setminus \{x_0\} \rangle \in \Sigma$ and hence its primitive generators are linearly independent in $N_{\mathbb{Q}}$.
- Σ_U and $\overline{\Sigma}_U$ are compatible with $\phi_{\mathbb{Q}}$ as we have $\phi_{\mathbb{Q}}(\tau') \in \overline{\Sigma}_U$ for any $\tau' \in \Sigma_U$.

\square

Proof of Proposition 2.14. Let the notation be as above. Let $\hat{\Sigma}_U$ be the collection of fans of the form (3) above with $m = 0$. It follows from the description of the cones of Σ_U in Lemma 2.15 that

- (1) $\phi_{\mathbb{Q}}$ maps each cone $\hat{\tau} \in \hat{\Sigma}_U$ bijectively to a cone $\bar{\tau} \in \overline{\Sigma}_U$ such that $\phi(\hat{\tau} \cap N) = \bar{\tau} \cap \overline{N}$. Furthermore, the map $\hat{\tau} \mapsto \bar{\tau}$ defines a bijection $\hat{\Sigma}_U \rightarrow \overline{\Sigma}_U$;

- (2) given cones $\hat{\tau} \in \hat{\Sigma}_U$ and $\tau_0 \in \Sigma_0$, the sum $\hat{\tau} + \tau_0$ lies in Σ_U and every cone of Σ_U arises in this way.

In the notation of [CLS11, Definition 3.3.18], we say that Σ_U is *split by $\bar{\Sigma}_U$ and Σ_0* . We conclude by [CLS11, Theorem 3.3.19] that $U = X \setminus V(\mathcal{E}_P)$ is a locally trivial fiber bundle over $X_{\bar{\Sigma}_U}$ with fiber $X_{\Sigma_0} \simeq \mathbb{P}^k$. It follows automatically that $X_{\bar{\Sigma}_U}$ is smooth, since it is the base of a locally trivial fibration with a smooth total space. \square

2.3. Some properties of primitive collections. Before focusing on toric Fano manifolds, we collect here two useful properties of primitive collections of arbitrary toric manifolds. The first one, by Sato, describes the behaviour of primitive collections under a smooth toric blowdown. The second one, by Batyrev, describes primitive collections on toric manifolds of Picard rank 3.

Proposition 2.17. ([Sat00, Corollary 4.9]) Let X be a toric manifold, and let $f: X \rightarrow Y$ be the contraction associated to an extremal class in $\text{NE}(X)$, corresponding to a primitive relation of the form

$$r(Q): t_1 + \cdots + t_s = z.$$

Then the fan Σ_Y is obtained from Σ_X by removing the ray generated by z , and X is the blowup of Y along $V(t_1, \dots, t_s)$. Furthermore, the primitive collections of Y are precisely the following $P_Y \in \text{PC}(Y)$:

- $P_Y = P_X$ for some $P_X \in \text{PC}(X)$ such that $z \notin P_X$ and $P_X \neq Q = \{t_1, \dots, t_s\}$;
- $P_Y = (P_X \setminus \{z\}) \cup \{t_1, \dots, t_s\}$ for some $P_X \in \text{PC}(X)$ such that $z \in P_X$ and $(P_X \setminus \{z\}) \cup S \notin \text{PC}(X)$ for any subset $S \subsetneq \{t_1, \dots, t_s\}$.

Proposition 2.18. ([Bat91, Theorem 5.7, Theorem 6.6]) Let X be a projective toric manifold with $\rho(X) = 3$. Then the number of primitive collections of Σ_X is either $l = 3$ or $l = 5$. Moreover, the set of generators $G(\Sigma_X)$ can be written as a disjoint union of l nonempty subsets

$$G(\Sigma_X) = X_0 \sqcup \cdots \sqcup X_{l-1}$$

that define primitive collections and relations as follows:

- Case $l = 3$. Each X_0, X_1, X_2 is a primitive collection, and the corresponding primitive relations are extremal.
- Case $l = 5$. There are five primitive collections of the form $X_i \sqcup X_{i+1}$, $0 \leq i \leq 4$, where $X_5 := X_0$. To describe the primitive relations of X , we use the following notation. We fix a labelling (v_1, \dots, v_k) for the elements of X_i . If $\bar{c} = (c_1, \dots, c_k) \in \mathbb{Z}^k$, then $\bar{c} \cdot X_i$ stands for $c_1 v_1 + \cdots + c_k v_k$. Moreover, we set $\bar{1} = (1, \dots, 1)$. Then there are vectors \bar{c} and \bar{b} of nonnegative integers such that at least one entry in \bar{c} is zero (up to relabelling, we may assume that $c_1 = 0$), and the primitive relations of X are the following:

$$\begin{aligned} r_0: \quad & \bar{1} \cdot X_0 + \bar{1} \cdot X_1 = \bar{c} \cdot X_2 + (\bar{b} + \bar{1}) \cdot X_3 \\ r_1: \quad & \bar{1} \cdot X_1 + \bar{1} \cdot X_2 = \bar{1} \cdot X_4, \\ r_2: \quad & \bar{1} \cdot X_2 + \bar{1} \cdot X_3 = 0, \\ r_3: \quad & \bar{1} \cdot X_3 + \bar{1} \cdot X_4 = \bar{1} \cdot X_1, \\ r_4: \quad & \bar{1} \cdot X_4 + \bar{1} \cdot X_0 = \bar{c} \cdot X_2 + \bar{b} \cdot X_3. \end{aligned}$$

The relations r_0 , r_1 and r_3 are extremal, while $r_2 = r_1 + r_3$ and $r_4 = r_0 + r_3$.

Remark 2.19. In [SS20, Corollary 1.2], Sato and Suyama use Proposition 2.18 to show that projective spaces are the only toric 2-Fano manifolds with Picard rank $\rho \leq 3$.

2.4. Primitive collections on toric Fano manifolds. Let X be a projective toric manifold with regular complete fan Σ_X in $N_{\mathbb{Q}}$.

Proposition 2.20. ([Bat99, Proposition 2.3.6]) The toric variety X is Fano if and only if all primitive collections of Σ_X have strictly positive degree.

Proposition 2.21. ([Cas03a, Corollary 4.4]) Assume that X is Fano, and let $P \in \text{PC}(X)$. If $\deg(P) = 1$, then the corresponding curve class is extremal.

Proposition 2.22. Assume that X is Fano, and let $x \in G(\Sigma_X)$.

- (1) There is at most one primitive collection of order 2 and degree 2 containing x . If it exists, then it is of the form $x + (-x) = 0$, and $m(X) = 1$.
- (2) ([Cas03b, Lemma 3.3]) There are at most two primitive collections of order 2 and degree 1 containing x . If there are exactly two of them, then they are of the form $x + y = (-w)$ and $x + w = (-y)$, and $m(X) = 1$.

Corollary 2.23. Assume that X is Fano and $m(X) > 1$. Then any $x \in G(\Sigma_X)$ is contained in at most one primitive collection of order 2. If there is such a primitive collection, then it is of the form $x + y = z$.

Definition 2.24. Let $x \in G(\Sigma_X)$. We say that $y \in G(\Sigma_X)$ is an *opponent* of x if $\langle x, y \rangle \notin \Sigma_X$.

Notation 2.25. Assume that X is Fano and $m(X) > 1$. By Corollary 2.23, each vector $x \in G(\Sigma_X)$ has at most one opponent. If such an opponent exists, we denote it by x' .

Lemma 2.26. Assume that X is Fano and $m(X) > 1$. Consider a pair of opponents $x, x' \in G(\Sigma_X)$. If there exist $y, z \in G(\Sigma_X)$ such that $x + x' = y + z$, then $z = y'$.

Proof. If $y = x$ or $y = x'$, the claim follows automatically. So we assume otherwise. Note that $\{x, x'\} \in \text{PC}(X)$, and y and z do not form a cone, as otherwise $x + x' = y + z$ would give us a primitive relation of degree 0, which is impossible for a toric Fano manifold. \square

Lemma 2.27. Assume that X is Fano and $m(X) > 1$. Assume that there exist x, y, z, u, v in $G(\Sigma_X)$ such that $(*)$ $x + y + z = u + v$ and such that $\langle u, v \rangle \in \Sigma_X$. Then exactly one of the following must happen:

- a. The vectors x, y, z are pairwise distinct, and $\{x, y, z\}$ is a primitive collection with primitive relation $(*)$. In particular, the corresponding curve class is extremal.
- b. Up to relabeling, $v = z$, $y = x'$ and $x + x' = u$.

Proof. Assume two of $\{x, y, z\}$ do not form a cone. For example, assume x, y do not form a cone. Then $y = x'$, the opponent of x . Let $x + x' = \alpha$, for some $\alpha \in G(\Sigma_X)$. We have that $\alpha + z = u + v$. As $\langle u, v \rangle \in \Sigma_X$, Proposition 2.6 implies that $\alpha + z = u + v$ corresponds to an effective class of degree 0, which therefore implies that $\{\alpha, z\} = \{u, v\}$. Up to relabeling, we may assume $v = z$, and hence, $x + x' = u$ and we are in the situation b.

Assume now that any two vectors in $\{x, y, z\}$ form a cone. Then x, y, z are mutually disjoint and $\langle x, y, z \rangle \notin \Sigma_X$, as otherwise by Proposition 2.6 we obtain an effective curve

class of degree -1 , contradicting the fact that X is Fano. It follows that $\{x, y, z\}$ is a primitive collection. Since $\langle u, v \rangle \in \Sigma_X$, it follows that $(*)$ is the associated primitive relation. Proposition 2.21 now implies that the corresponding curve class is extremal. \square

3. TORIC FANO MANIFOLDS WITH $m(X) = 1$

In this section, we study toric Fano manifolds with $m(X) = 1$, and follow the strategy outlined in the introduction to show that they cannot be 2-Fano.

For any $x \in G(\Sigma_X)$, the set of primitive collections containing x is denoted by

$$\text{PC}_x(X) = \{P \in \text{PC}(X) \mid x \in P\}.$$

Proposition 3.1. ([Cas03b, Lemma 3.1]) Assume that X is a toric Fano manifold and that $P = \{x, -x\} \in \text{PC}(X)$.

- (1) Any $Q \in \text{PC}_x(X) \setminus \{P\}$ has degree 1 (hence is extremal by Proposition 2.21), and $r(Q)$ is of the form

$$r(Q): x + \underbrace{y_1 + \cdots + y_h}_{\in \langle Q \setminus \{x\} \rangle} = \underbrace{z_1 + \cdots + z_h}_{\in \sigma(Q)},$$

where we denote $\langle Q \setminus \{x\} \rangle := \langle y_1, \dots, y_h \rangle$ and $\sigma(Q) := \langle z_1, \dots, z_h \rangle$.

- (2) For any $R \in \text{PC}_x(X) \setminus \{P, Q\}$, we have

$$V(R \setminus \{x\}) \cap V(Q \setminus \{x\}) = \emptyset \quad \text{and} \quad V(R \setminus \{x\}) \cap V(\sigma(Q)) = \emptyset.$$

- (3) For any $Q \in \text{PC}_x(X) \setminus \{P\}$ with $r(Q): x + y_1 + \cdots + y_h = z_1 + \cdots + z_h$ we have $Q' = \{-x, z_1, \dots, z_h\} \in \text{PC}_{-x}(X)$, Q' has degree 1 (hence is extremal) and

$$r(Q'): -x + \underbrace{z_1 + \cdots + z_h}_{\in \langle Q' \setminus \{-x\} \rangle = \sigma(Q)} = \underbrace{y_1 + \cdots + y_h}_{\in \sigma(Q') = \langle Q \setminus \{x\} \rangle}.$$

Corollary 3.2. Let X be a toric manifold, and $P = \{x, -x\} \in \text{PC}(X)$. With Notation 2.12,

$$(4) \quad \mathcal{E}_P = \{\langle Q \setminus \{v\} \rangle \mid Q \in \text{PC}_v(X) \setminus \{P\}, \quad v = \pm x\}.$$

If moreover X is Fano, then $V(\mathcal{E}_P)$ has 0, 2 or 4 components of codimension 1 in X .

Proof. Let X be a toric manifold, and $P = \{x, -x\} \in \text{PC}(X)$. The description of $V(\mathcal{E}_P)$ in Equation (4) follows from Notation 2.12. In the Fano case, the number of components of codimension 1 of $V(\mathcal{E}_P)$ equals the number of primitive collections of order 2 and degree 1 containing x or $-x$, which is 0, 2 or 4 by Proposition 2.22 and Proposition 3.1. \square

Proposition 3.3. Let X be a toric Fano manifold, and $P = \{x, -x\} \in \text{PC}(X)$. Then there exists a birational morphism $f: X \rightarrow Y$ such that

- $P_Y := \{x, -x\} \in \text{PC}(Y)$,
- $V(\mathcal{E}_{P_Y})$ has codimension ≥ 2 in Y ,

- f is a composition of at most two blow-downs with disjoint centers and smooth target:

$$(5) \quad \begin{aligned} \text{Exc}(f) &= \bigcup_{\substack{Q \in \text{PC}_x(X) \setminus \{P\}: \\ \text{ord}(Q)=2}} V(\sigma(Q)) \subset X, \\ f(\text{Exc}(f)) &= \bigcup_{\substack{Q \in \text{PC}_x(X) \setminus \{P\}: \\ \text{ord}(Q)=2}} V(Q) \subset Y. \end{aligned}$$

Proof. If $V(\mathcal{E}_P)$ has codimension ≥ 2 in X , i.e., if P is the unique primitive collection of order 2 containing x , then the statement holds with $f = \text{Id}$.

Assume now that $V(\mathcal{E}_P)$ has 2 components of codimension 1, i.e. we have primitive relations

$$r(P): x + (-x) = 0, \quad r(Q_1): x + y = z, \quad r(Q'_1): -x + z = y,$$

and, for any other $R \in \text{PC}_x(X) \cup \text{PC}_{-x}(X)$, one has $\text{ord}(R \setminus \{\pm x\}) \geq 2$. Let $f_1: X \rightarrow Y$ be the smooth blow-down induced by the extremal ray of $\text{NE}(X)$ corresponding to $r(Q_1)$. By Proposition 2.17 and Proposition 3.1, $\text{PC}_x(Y) \cup \text{PC}_{-x}(Y)$ consists of

- $P_Y = \{x, -x\}$;
- $R_Y = R_X$ for some $R_X \in \text{PC}_x(X) \cup \text{PC}_{-x}(X) \setminus \{Q_1\}$ such that $z \notin R_X$. In particular we have $\text{ord}(R_Y \setminus \{\pm x\}) \geq 2$;
- $R_Y = (R_X \setminus \{z\}) \cup \{x, y\}$ for some $R_X \in \text{PC}_z(X)$ such that $(R_X \setminus \{z\}) \cup \{x\} \notin \text{PC}(X)$ and $(R_X \setminus \{z\}) \cup \{y\} \notin \text{PC}(X)$. In particular, we have $\langle R_Y \setminus \{x\} \rangle = \langle R_X \setminus \{z\}, y \rangle$, so $\text{ord}(R_Y \setminus \{x\}) \geq 2$.

It follows that $V(\mathcal{E}_{P_Y})$ has codimension ≥ 2 in Y , so the proposition holds with $f = f_1$.

Assume now that $V(\mathcal{E}_P)$ has 4 components of codimension 1, i.e., we have


$$\begin{array}{lll} r(P): x + (-x) = 0 & r(Q_1): x + y = -w & r(Q'_1): -x + (-w) = y \\ y + (-y) = 0 & r(Q_2): x + w = -y & r(Q'_2): -x + (-y) = w \\ w + (-w) = 0 & w + y = -x & -y + (-w) = x \end{array}$$

by Proposition 2.22. By [Cas03b, p.1487, Case (3)], any other primitive collection $R \in \text{PC}(X)$ is disjoint from $\{x, -x, y, -y, w, -w\}$. Let $f_1: X \rightarrow X_1$ be the smooth blow-down induced by the extremal ray of $\text{NE}(X)$ corresponding to $r(Q_1)$. By Proposition 2.17 the primitive collections of X_1 containing x or $-x$ are only

$$P_{X_1} = P \quad (Q_2)_{X_1} = Q_2 \quad (Q'_2)_{X_1} = Q'_2.$$

It follows that $r(Q_2)$ corresponds to an extremal curve class in $\text{NE}(X_1)$. Let $f_2: X_1 \rightarrow X_2$ be the smooth blow-down induced by the extremal ray of $\text{NE}(X)$ corresponding to $r(Q_2)$. By Proposition 2.17 $P_{X_2} = \{x, -x\}$ is the only primitive collection in $\text{PC}_x(X_2) \cup \text{PC}_{-x}(X_2)$. This implies that $V(\mathcal{E}_{P_{X_2}}) = \emptyset$, so the proposition holds with $Y = X_2$ and $f = f_2 \circ f_1$. \square

Construction 3.4. Let Y be a projective toric manifold of dimension ≥ 3 , and $P = \{x, -x\} \in \text{PC}(Y)$ a centered primitive collection of order 2. Let $V(\mathcal{E}_P) \subset Y$ be the closed subset defined in Notation 2.12, set $U := Y \setminus V(\mathcal{E}_P)$, and let $\pi: U \rightarrow W$ be the \mathbb{P}^1 -bundle given by Proposition 2.14.



Assume that $V(\mathcal{E}_P)$ has codimension ≥ 2 in Y , and let $Z \subset Y$ be any given closed subset of codimension ≥ 2 in Y . Note that a toric manifold is rational, hence rationally connected, so by [Kol96, Proposition II.3.7], there is a smooth (very free) rational curve $C \subset U \setminus Z$. Consider the surface $S := \pi^{-1}(\pi(C)) \subset U$, and let $n: \tilde{S} \rightarrow S$ be its normalization. Then \tilde{S} is a Hirzebruch surface with \mathbb{P}^1 -bundle structure $\tilde{\pi}: \tilde{S} \rightarrow \mathbb{P}^1$ induced by π . The curve class of the image of a fiber of $\tilde{\pi}$ on Y corresponds to the centered relation $x + (-x) = 0$. By taking C general, we may assume that $\pi(C)$ and $\pi(Z)$ meet transversely in at most finitely many general points. Hence S and Z meet transversely in at most finitely many points.

Proposition 3.5. Let Y be a projective toric manifold of dimension ≥ 3 , and $P_Y = \{x, -x\} \in \text{PC}(Y)$ a centered primitive collection of order 2. Assume that $V(\mathcal{E}_P)$ has codimension ≥ 2 in Y , and let $S \subset Y$ be as in Construction 3.4. Then $S \cdot \text{ch}_2(Y) = 0$.

Proof. Let $S = \pi^{-1}(\pi(C))$ be the surface from Construction 3.4, $n: \tilde{S} \rightarrow S$ its normalization, $\tilde{\pi}: \tilde{S} \rightarrow \mathbb{P}^1$ the \mathbb{P}^1 -bundle structure induced by π , and F a fiber of $\tilde{\pi}$. Our goal is to compute

$$S \cdot \text{ch}_2(Y) = \frac{1}{2} \sum_{v \in G(\Sigma_Y)} S \cdot V(v)^2 = \frac{1}{2} \sum_{v \in G(\Sigma_Y)} (n^*V(v))^2.$$

Recall that the curve class of the image of F in Y is associated to the relation $x + (-x) = 0$. By restricting the divisors $n^*V(v)$ to F , we have:

$$\begin{cases} n^*V(v) \cdot F = 0 & \text{if } v \neq x, -x, \\ n^*V(x) \cdot F = 1, \\ n^*V(-x) \cdot F = 1. \end{cases}$$

Hence there are sections σ, σ' of $\tilde{\pi}: \tilde{S} \rightarrow \mathbb{P}^1$, and $\alpha, \beta, \gamma \in \mathbb{Z}$ such that, on \tilde{S} ,

$$\begin{cases} n^*V(v) = \alpha F & \text{if } v \neq x, -x, \\ n^*V(x) = \sigma + \beta F, \\ n^*V(-x) = \sigma' + \gamma F. \end{cases}$$

Therefore

$$\begin{aligned}
\sum_{v \in G(\Sigma_Y)} (n^*V(v))^2 &= \sum_{v \neq x, -x} \overline{(n^*V(v))^2} + (n^*V(x))^2 + (n^*V(-x))^2 \\
&= \sigma^2 + 2\beta + \sigma'^2 + 2\gamma \\
&= \sigma^2 - 2(\sigma \cdot \sigma') + \sigma'^2 \quad \text{as } \sigma \cdot \sigma' + \beta + \gamma = n^*V(x) \cdot n^*V(-x) = 0 \\
&= (\sigma - \sigma')^2 = 0 \quad \text{as } \sigma - \sigma' \text{ is a multiple of } F,
\end{aligned}$$

and this concludes the proof. \square

Lemma 3.6. ([dS06a, Lemma 5.1]) Consider the blowup diagram

$$\begin{array}{ccc}
E & \xhookrightarrow{j} & X := \text{Bl}_Z Y \\
\pi := f|_E \downarrow & & \downarrow f \\
Z & \hookrightarrow & Y
\end{array}$$

where both Y and Z are smooth projective varieties and $\text{codim}_Y Z = c \geq 2$. Then we have the following relation between the 2nd Chern characters of X and Y :

$$\text{ch}_2(X) = f^* \text{ch}_2(Y) + \frac{c+1}{2} E^2 - j_* \pi^* c_1(\mathcal{N}_{Z/Y}).$$

Corollary 3.7. In the setting of Lemma 3.6, assume that $c = 2$. Let $S \subset Y$ be a surface that intersects Z at most transversely at $k \geq 0$ points, and let $S_X \subset X$ be its strict transform. Then

$$\text{ch}_2(X) \cdot S_X = \text{ch}_2(Y) \cdot S - \frac{3}{2} \cdot k.$$

Proof. Write $S \cap Z = \{p_1, \dots, p_k\}$. Then S_X is isomorphic to the blowup of S at p_1, \dots, p_k , $S_X \cap E = \cup_{i=1}^k e_i$, where $e_i \simeq \mathbb{P}^1$ is the exceptional curve over p_i , and $(e_i^2)_{S_X} = -1$. By Lemma 3.6,

$$\begin{aligned}
\text{ch}_2(X) \cdot S_X &= f^* \text{ch}_2(Y) \cdot S_X + \frac{c+1}{2} E^2 \cdot S_X - j_* \pi^* c_1(\mathcal{N}_{Z/Y}) \cdot S_X \\
&= \text{ch}_2(Y) \cdot S + \frac{3}{2} (E|_{S_X})^2 - \pi^* c_1(\mathcal{N}_{Z/Y}) \cdot S_{X|E} \\
&= \text{ch}_2(Y) \cdot S + \frac{3}{2} \sum_{i=1}^k (e_i)^2 - \pi^* c_1(\mathcal{N}_{Z/Y}) \cdot \sum_{i=1}^k e_i \\
&= \text{ch}_2(Y) \cdot S - \frac{3}{2} k - \sum_{i=1}^k \overline{c_1(\mathcal{N}_{Z/Y}) \cdot p_i}.
\end{aligned}$$

\square

We are ready to prove Theorem 1.2: a toric Fano manifold X with $m(X) = 1$ is not 2-Fano.

Proof of Theorem 1.2. Let X be a toric Fano manifold with $m(X) = 1$, and fix a centered primitive relation $r(P): x + (-x) = 0$. Let $f: X \rightarrow Y$ be as in Proposition 3.3, and $\pi: U = Y \setminus V(\mathcal{E}_{P_Y}) \rightarrow W$ the \mathbb{P}^1 -bundle structure induced by $r(P_Y): x + (-x) = 0$ (see Proposition 2.14). By Equation (5), $Z := f(\text{Exc}(f))$ is empty or has codimension 2 in Y .

Let $S \subset Y$ be the surface given by Construction 3.4. Then S and Z meet transversely in at most finitely many points. It follows from Proposition 3.5 that $\text{ch}_2(Y) \cdot S = 0$. Let S_X be the strict transform of S in X . By Corollary 3.7, $\text{ch}_2(X) \cdot S_X \leq \text{ch}_2(Y) \cdot S = 0$, and so X is not 2-Fano. \square

4. PROOF OF THEOREM 1.3

In this section, we work in the setting of Theorem 1.3:

- (I) X is a Fano manifold of dimension $n \geq 6$ with fan $\Sigma = \Sigma_X$ in $N_{\mathbb{Q}}$;
- (II) $m(X) \geq 3$;
- (III) $r(P): x_0 + \cdots + x_{n-2} = 0$ is a centered primitive relation in Σ .

The equality $m(X) = n - 2$, as well as uniqueness of the centered primitive collection, follows immediately from Lemma 2.5. Now the goal is to prove that $\rho(X) \leq 3$, and this will follow from Proposition 4.11, Proposition 4.16 and Proposition 4.17. By Remark 2.19, we conclude that the projective space \mathbb{P}^n is the only smooth n -dimensional toric 2-Fano variety with $m(X) \geq n - 2$ (Corollary 1.5).

Let $P := \{x_0, \dots, x_{n-2}\}$ and set $\Gamma = \text{Span } P \subset N_{\mathbb{Q}}$. Consider the quotient

$$\pi: N_{\mathbb{Q}} \simeq \mathbb{Q}^n \rightarrow \mathbb{Q}^n / \Gamma \simeq \mathbb{Q}^2.$$

Since X is complete, the support of Σ is equal to $N_{\mathbb{Q}}$, and hence we can find generators

- (IV) $x, y, z \in G(\Sigma) \setminus P$, for which
- (V) $0 \in \text{Conv}(\pi(x), \pi(y), \pi(z))$.

Lemma 4.1. The nonnegative span $\langle x, y, z \rangle$ is not a cone in Σ .

Proof. We will argue by contradiction and assume that $\langle x, y, z \rangle \in \Sigma$. By (V), we can find a nonnegative triple of constants (c_1, c_2, c_3) such that $c_1\pi(x) + c_2\pi(y) + c_3\pi(z) = 0$, or in other words

$$v := c_1x + c_2y + c_3z = a_0x_0 + \cdots + a_{n-2}x_{n-2}$$

for some constants a_i . Note that by adding some multiple of $r(P)$ (see (III)) to the right hand side, we can assume the a_i 's are nonnegative and such that at least one of them, say a_j , is 0. It follows that v lies in two cones of Σ , namely

$$v \in \langle x, y, z \rangle \cap \langle x_0, \dots, \check{x}_j, \dots, x_{n-2} \rangle,$$

which is impossible since $\{x, y, z\} \cap S = \emptyset$. \square

Since $\{x, y, z\}$ does not span a cone, we conclude that

- (1) either it is a primitive collection,
- (2) or two of these vectors do not form a cone.

The former case is the more technical one, and we start with it in Section 4.1. After we are done analyzing it, we can assume that none of the triples $\{x, y, z\}$ as in (IV) and (V) form a primitive collection, and this case will be treated in Section 4.2.

4.1. First case: $\{x, y, z\}$ is a primitive collection. We recall that if X is a projective toric Fano manifold and $m(X) > 1$, then any $x \in G(\Sigma)$ has at most one opponent by Corollary 2.23, where the opponent of x is an element $x' \in G(\Sigma)$ such that $\{x, x'\} \in \text{PC}(X)$. When we write $\{x, x'\}$, we mean either the set of two elements if x' exists, or the singleton $\{x\}$ if x' does not exist.

Remark 4.2. In the setting (I)—(V), pick a generator $u \in G(\Sigma)$. Then (V) and plane geometry imply that the convex hull of $\pi(u)$ together with two of the vectors $\pi(x)$, $\pi(y)$, $\pi(z)$ contains 0.

Lemma 4.3. In the setting (I)—(V), assume in addition that $Q := \{x, y, z\}$ is a primitive collection. Then the corresponding primitive relation is

- either $r(Q): x + y + z = x_i + x_j$ for possibly equal $x_i, x_j \in P$,
- or $r(Q): x + y + z = v$ for some $v \in G(\Sigma)$.

Proof. Since X is Fano by (I), the degree of the primitive relation $x + y + z = A$ is positive, so we can have only three possibilities for A . The first one with $A = 0$ is actually not possible by our assumption (II). The second is $A = v$, and we cannot say much about v at the moment. The last possibility is

$$r(Q): x + y + z = u + v$$

for some, possibly equal, $u, v \in G(\Sigma)$. By Proposition 2.21, this is an extremal primitive relation, so by Proposition 2.8 applied to $r(Q)$ and $\tau = \{0\}$, we get that $\langle u, x, y \rangle$, $\langle u, y, z \rangle$, $\langle u, x, z \rangle \in \Sigma$. By Remark 4.2, we may assume without loss of generality that $0 \in \text{Conv}(\pi(u), \pi(x), \pi(y))$, hence by Lemma 4.1, we get $u \in P$. The same argument applies to conclude $v \in P$. \square

Hence we have two cases to consider: when $\deg(Q)$ is 1 and when it is 2.

4.1.1. Degree one. In this case, by Lemma 4.3, we have

(VI) a primitive relation $r(Q): x + y + z = x_i + x_j$ for possibly equal $x_i, x_j \in P$.

Lemma 4.4. In the setting (I)—(VI), assume in addition that $G(\Sigma)$ is contained in $P \cup \{x, y, z, x', y', z'\}$. Then x', y', z' do not exist. Consequently, $\rho(X) \leq 2$.

Proof. By contradiction, assume that $x' \in G(\Sigma)$ exists, so $\{x, x'\}$ is a primitive collection, and let $x + x' = \alpha$ be the corresponding primitive relation. Clearly $\alpha \neq x, x'$. The relation $r(Q)$ (see (VI)) gives

$$\alpha + y + z = x_i + x_j + x',$$

which shows $\alpha \notin P \cup \{y, z\}$ since otherwise the left hand side (LHS) forms a cone and we get an effective class of degree zero. Indeed, if $\alpha \in \{y, z\}$ it is clear that the LHS would form a cone; if $\alpha \in P$ applying Proposition 2.8 to $r(Q)$ and $\tau = \langle \alpha \rangle$ would follow that the LHS is a cone.

So without loss of generality, we can assume $x + x' = y'$. Applying the same argument to y' , we get that $y + y' = x'$ or $y + y' = z'$. The former would imply $x + y = 0$, which is not possible by (II), so $y + y' = z'$. Again, applying the same argument to z' , we get $z + z' = x'$. Summing the three primitive relations, we obtain $x + y + z = 0$, which contradicts (VI). \square

This leaves us with the case when

(VII) there exists $u \in G(\Sigma) \setminus (P \cup \{x, y, z, x', y', z'\})$.

By Remark 4.2, we can assume without loss of generality that

$$(VIII) \quad 0 \in \text{Conv}(\pi(x), \pi(y), \pi(u)).$$

Lemma 4.5. Assume (I)—(VIII), then $R := \{x, y, u\}$ is a primitive collection with primitive relation $r(R): x + y + u = v$ for some $v \in G(\Sigma)$.

Proof. By Lemma 4.1, we have $\langle x, y, u \rangle \notin \Sigma$, and since $u \neq x', y'$, we conclude that $\{x, y, u\}$ is a primitive collection. By Lemma 4.3, we can have either $r(R): x + y + u = x_k + x_l$ or $r(R): x + y + u = v$. In the former case, combining $r(R)$ with $r(Q)$ (see (VI)) provides us with a relation

$$u + x_i + x_j = z + x_k + x_l.$$

By applying Proposition 2.8 to $r(Q)$ (see (VI)) and $\tau = \langle x_k, x_l \rangle$, we get $\langle z, x_k, x_l \rangle \in \Sigma$ (here we are using the assumption $\dim(X) = n \geq 6$ in (I)). Hence, by Proposition 2.6, we get an effective curve class of degree 0, contradicting the Fano assumption (I). \square

We will write down the result of Lemma 4.5 as an additional assumption, remembering that it is implied by the previous assumptions:

$$(IX) \quad \text{We have a primitive relation } r(R): x + y + u = v \text{ for some } v \in G(\Sigma).$$

Lemma 4.6. Assume (I)—(IX), then

$$G(\Sigma) \subset P \cup \{x, y, z, u, x', y', u', v'\}.$$

In particular, $z' \in P \cup \{x, y, z, u, x', y', u', v'\}$, and since $v \neq x, y, u, v'$, we have that $v \in P$ or $v \in \{z, x', y', u'\}$.

Proof. Take any $w \in G(\Sigma) \setminus (P \cup \{x, y, z, u, x', y', u'\})$. By Remark 4.2, the convex hull of $\pi(w)$ together with two of $\pi(x), \pi(y), \pi(u)$ contains 0, yielding an analog of (VIII). By Lemma 4.1 and from $w \neq x', y', u'$, it follows that one of $\{w, x, u\}, \{w, y, u\}, \{w, x, y\}$ is a primitive collection.

We will prove that the corresponding primitive relation has the form $w + x + u = b$ or $w + y + u = b$ or $w + x + y = b$ for some $b \in G(\Sigma)$. Assume for a contradiction that this is not the case. By Lemma 4.3, we have that one of $w + x + u, w + y + u$, or $w + x + y$ equals $x_k + x_l$, for possibly equal $x_k, x_l \in P$. Hence there exist $a, b \in G(\Sigma)$ (one of which is $w \neq z$) such that either $x + a + b$ or $y + a + b$ equals $x_k + x_l$. Assume $x + a + b = x_k + x_l$. Combining this with $r(Q)$ (see (VI)), it follows that $y + z + x_k + x_l = a + b + x_i + x_j$. By applying Proposition 2.8 to $r(Q)$ (see (VI)) and $\tau = \langle x_k, x_l \rangle$, we get $\langle y, z, x_k, x_l \rangle \in \Sigma$ (here we are using the assumption $\dim(X) = n \geq 6$ in (I)). Hence, by Proposition 2.6, we get an effective curve class of degree 0. The class is non-trivial since $w \neq y, z, x_k, x_l$. This contradicts the assumption that X is Fano. The case $y + a + b = x_k + x_l$ is similar.

So we have a primitive relation of the form $w + x + u = b$ or $w + y + u = b$ or $w + x + y = b$ for some $b \in G(\Sigma)$. Combining this with $r(R)$ (see (IX)), we get $b + y = w + v$ or $b + x = w + v$ or $b + u = w + v$. All possibilities imply that $w = v'$, as otherwise, by Proposition 2.6, we obtain an effective class of degree 0 (non-trivial since $w, v \neq y, u$), which contradicts the Fano assumption (I). \square

Lemma 4.7. Assume (I)—(IX). Then x, y don't have opponents,

$$G(\Sigma) \subset P \cup \{x, y, z, u, u', v'\},$$

and we have a trichotomy: $v \in P$ or $v = z$ or $v = u'$.

Proof. We will only show that x' does not exist, and the argument for y' is symmetric. Suppose to the contrary that $x' \in G(\Sigma)$, and let $x + x' = \alpha$ be the corresponding primitive relation, so $\alpha \neq x, x'$. Then we can substitute $x = \alpha - x'$ into $r(Q)$ (see (VI)) to get

$$\alpha + y + z = x' + x_i + x_j,$$

which shows $\alpha \notin P \cup \{y, z\}$, as otherwise we would get an effective curve class of degree 0. Substituting $x = \alpha - x'$ into $r(R)$ (see (IX)) gives the relation

$$(6) \quad \alpha + y + u = v + x'.$$

Note that $\langle v, x' \rangle \in \Sigma$ by Corollary 2.23. We claim that Equation (6) is an extremal primitive relation by Lemma 2.27. Assume not, then Lemma 2.27 implies that one of $\{\alpha, y, u\}$ is in $\{v, x'\}$ and the remaining two vectors are opponents. Since $u \neq y'$, then either α, y are opponents and $u \in \{v, x'\}$, or α, u are opponents and $y \in \{v, x'\}$. From (IX), we have $u \neq v$ and $y \neq v$; (VII) means $u \neq x'$; and (VI) implies $y \neq x'$. So both cases are impossible.

So Equation (6) is an extremal primitive relation. In particular, $\alpha \neq y', u, u'$. By Proposition 2.8, v forms a cone with α , hence $\alpha \neq v'$, which contradicts Lemma 4.6. \square

Lemma 4.8. Assume (I)—(IX). Then we have

$$G(\Sigma) \subset P \cup \{x, y, z, u, v'\}$$

and a dichotomy: $v \in P$ or $v = z$. If moreover u' exists, we have $u + u' = z$, $v = x_i$ and $u = x'_j$.

Proof. Assume that u' exists and let $u + u' = \alpha$ be the corresponding primitive relation. Then substituting it into $r(R)$ (see (IX)) gives a relation

$$(7) \quad x + y + \alpha = u' + v.$$

By $r(R)$ (see (IX)), $v \neq u$, so $\langle u', v \rangle \in \Sigma$. Since there are no x' and y' , it follows from Lemma 2.27 that Equation (7) is an extremal primitive relation. It follows that $\alpha \notin \{u, u', x, y, v'\}$. Moreover, $\alpha \notin P$, as otherwise applying Proposition 2.8 to $r(Q)$ and $\tau = \langle \alpha \rangle$ would imply $\langle x, y, \alpha \rangle \in \Sigma$. So by Lemma 4.7, the only possibility is that $\alpha = z$. Therefore $u' + v = x_i + x_j$ by $r(Q)$ (see (VI)), so we have, after possibly relabeling, that $v = x_i$, $u' = x_j$, which by Lemma 4.7 proves the statement.

If instead u' does not exist, the statement follows directly from Lemma 4.7. \square

Lemma 4.9. Assume (I)—(IX). Then z doesn't have an opponent.

Proof. We argue by contradiction and assume that z' exists. Clearly $z' \neq x, y, z$ by $r(Q)$ (see (VI)) and $z' \neq u$ by (VII). Furthermore, $z' \neq u'$, otherwise we have $z = u$ by Corollary 2.23, which contradicts (VII). Finally, $z' \notin P$, otherwise $z' = x_k$ implies $z = x'_k$, and $r(Q)$ (see (VI)) becomes

$$x + y + x'_k = x_i + x_j,$$

but applying Proposition 2.8 to $r(Q)$ and $\tau = \langle x_k \rangle$, we get $\langle x_k, x'_k \rangle \in \Sigma$, a contradiction. Thus by Lemma 4.8, the only possibility is $z' = v'$, so $z = v$ by Corollary 2.23. By Lemma 4.8, this implies $G(\Sigma) \subset P \cup \{x, y, z, u, z'\}$.

Consider the primitive relation $z + z' = \beta \in G(\Sigma)$. We will show that $\beta = u$. Indeed, it is clear that $\beta \neq z, z'$. Combining $z + z' = \beta$ with $r(Q)$ (see (VI)), we have

$$x + y + \beta = x_i + x_j + z'.$$

If $\beta \in \{x, y\}$, the left hand side is a cone, and we get a non-trivial effective relation of degree 0, which is impossible by (I). If $\beta \in P$, applying Proposition 2.8 to $r(Q)$ and $\tau = \langle \beta \rangle$ implies that $\langle x, y, \beta \rangle \in \Sigma$, and we obtain a contradiction as before.

But now, substituting $z + z' = u$ into $r(R)$ (see (IX)) yields $x + y + z' = 0$, hence contradicting $m(X) \geq 3$ in (II). \square

Lemma 4.10. Assume (I)—(VII). Then $\rho(X) \leq 3$.

Proof. Let us summarize the consequences of (I)—(VII):

- (VIII) $0 \in \text{Conv}(\pi(x), \pi(y), \pi(u))$ after possibly relabeling x, y, z ;
- (IX) we have a primitive relation $r(R): x + y + u = v$ (Lemma 4.5);
 - x, y, z don't have opponents (Lemma 4.7, Lemma 4.9);
 - $G(\Sigma) \subset P \cup \{x, y, z, u, v'\}$ (Lemma 4.8), so $\rho(X) \leq 4$;
 - we have $v \in P$ or $v = z$ (Lemma 4.8), so we can consider two cases.

Case $v = z$. By Lemma 4.9, $v' = z'$ does not exist, hence it follows from Lemma 4.8 that $G(\Sigma) \subset P \cup \{x, y, z, u\}$, so $\rho(X) \leq 3$.

Case $v \in P$, say $v = x_l$. If $v = x_l$ doesn't have an opponent, then $G(\Sigma) \subset P \cup \{x, y, z, u\}$ and $\rho(X) \leq 3$. So the tricky case is when $v = x_l$ has an opponent $x'_l \notin P$.

Let $v + v' = \beta \in G(\Sigma)$. By $r(R)$, we have that $x + y + u + v' = v + v' = \beta$. Since $m(X) \geq 3$, $\beta \neq x, y, u, v'$. Hence $\beta \in P \cup \{z\}$. If $\beta \in P$, let $\beta = x_k$. Then $v' = x_k - x_l$ and $k \neq l$. Using $r(P)$, we obtain that $v' = 2x_k + \sum_{t \neq k, l} x_t$. As the vectors on the right hand side form a cone, we get an effective curve class of degree $2 - n$, which is impossible by (I).

So $\beta = z$. By $r(Q)$, we obtain that $x + y + v' = x_i + x_j - x_l$. If $l \neq i, j$, then again, by using $r(P)$ (see (III)), we may write $x_i + x_j - x_l$ as $x_i + x_j + \sum_{t \neq l} x_t$. As the vectors on the right hand side form a cone, we obtain an effective curve class of degree $3 - n$, which is impossible by (I).

So $l = i$ or $l = j$. Up to symmetry, we may assume $l = i$, so $v = x_i$, $x_i + x'_i = z$. By $r(Q)$, we have that $x + y + x'_i = x_j$. Combining this with $r(R)$, we obtain that $x'_i + x_i = u + x_j$. Furthermore, $u \neq v' = x'_i$ and $u \neq x_i$. If u, x_j form a cone, we obtain an effective non-trivial curve class of degree 0, which is impossible by (I). So $u = x'_j$, $v = x_i$ and we have two primitive relations

$$x_i + x'_i = z \quad \text{and} \quad x_j + x'_j = z.$$

Then either $i = j$, in which case $\rho(X) \leq 3$, or $i \neq j$, and we notice that they are both degree 1 and hence extremal, so we can perform the contraction associated to one of them, say we contract the curve class $x_i + x'_i = z$:

$$X \longrightarrow Y,$$

$$V(z) \longrightarrow V(x_i, x'_i).$$

By Proposition 2.17,

$$\begin{aligned} r(P_Y): x_0 + \cdots + x_{n-2} &= 0, \\ r(R_Y): x + y + x'_j &= x_i, \\ r(Q'): x + y + x'_i &= x_j, \\ r(Q''): x_j + x'_j &= x_i + x'_i \end{aligned}$$

are primitive relations in Y . Since $\rho(Y) = 3$, we apply Proposition 2.18. We adopt the same notation as in Proposition 2.18 and observe that we are in the case $l = 5$, hence

$$G(\Sigma) = \sqcup_{h=0}^4 X_h = \{x_0, \dots, x_j, \dots, x_{n-2}\} \sqcup \{x_j\} \sqcup \{x, y\} \sqcup \{x'_j\} \sqcup \{x'_i\},$$

and either $X_2 \sqcup X_3 = \{x_0, \dots, x_{n-2}\}$, or $\bar{c} = \bar{b} = \bar{0}$ and $X_4 \sqcup X_0 = \{x_0, \dots, x_{n-2}\}$. However, all possibilities for $\{x_j\}$ lead to a contradiction. Indeed:

- if $X_3 = \{x_j\}$, then we have $r_3: \bar{1} \cdot X_3 + \bar{1} \cdot X_4 = x_j + x'_j = x_i + x'_i \neq \bar{1} \cdot X_h$ for all $h = 0, \dots, 4$;
- if $X_2 = \{x_j\}$, then we have $r_1: \bar{1} \cdot X_1 + \bar{1} \cdot X_2 = x'_j + x_j = x_i + x'_i \neq \bar{1} \cdot X_h$ for all $h = 0, \dots, 4$;
- if $X_4 = \{x_j\}$, then we have $r_3: \bar{1} \cdot X_3 + \bar{1} \cdot X_4 = x'_j + x_j = x_i + x'_i \neq \bar{1} \cdot X_h$ for all $h = 0, \dots, 4$;
- if $X_0 = \{x_j\}$, then we have $r_0: \bar{1} \cdot X_0 + \bar{1} \cdot X_1 = x_j + x'_j = x_i + x'_i \neq \bar{c} \cdot X_2 + (\bar{b} + \bar{1})X_3 = \bar{1} \cdot X_3$.

This concludes the last case, and we get that $\rho(X) \leq 3$. \square

Proposition 4.11. Let X be a projective toric Fano manifold of $\dim(X) = n \geq 5$, $m(X) \geq 3$, which admits a primitive relation $r(P): x_0 + x_1 + \cdots + x_{n-2} = 0$ (assumptions (I)—(III)). Let $x, y, z \in G(\Sigma) \setminus P$ be such that $0 \in \text{Conv}(\pi(x), \pi(y), \pi(z))$ (assumptions (IV)—(V)). Assume in addition that $\{x, y, z\}$ is a primitive collection of degree 1 (assumption (VI)). Then $\rho(X) \leq 3$.

Proof. We recall that assumptions (I)—(III) imply (IV)—(V). Then either $G(\Sigma) \subset P \cup \{x, y, z, x', y', z'\}$, in which case $\rho(X) \leq 2$ by Lemma 4.4, or there exists a vector $u \in G(\Sigma) \setminus (P \cup \{x, y, z, x', y', z'\})$ as in (VII), in which case Lemma 4.10 implies $\rho(X) \leq 3$, and we are done. \square

4.1.2. *Degree two.* Still working in the setting (I)—(V), we assume that $\{x, y, z\}$ is a primitive collection whose primitive relation has degree 2, and by Proposition 4.11, we can exclude the case when degree one primitive collections as in (IV)—(VI) exist. In other words, these are the additional assumptions for this part of the proof:

- (X) we have a primitive relation $r(Q): x + y + z = v$ for some $v \in G(\Sigma)$;
- (XI) there is no primitive collection $\{a, b, c\} \subset G(\Sigma) \setminus P$ with $0 \in \text{Conv}(\pi(a), \pi(b), \pi(c))$ whose primitive relation has degree 1.

Lemma 4.12. In the setting (I)—(V) and (X)—(XI), we have

$$G(\Sigma) \subset P \cup \{x, y, z, x', y', z', v'\} \quad \text{and} \quad v \in P \cup \{x', y', z'\}.$$

Proof. If there is a generator $u \in G(\Sigma) \setminus (P \cup \{x, y, z, x', y', z'\})$, then by Remark 4.2 and (XI) we can assume that we have a primitive relation of the form

$$x + y + u = w.$$

Combining it with (X), we get $u + v = w + z$, so $u = v'$, otherwise $\langle u, v \rangle \in \Sigma$ and we get an effective non-trivial curve class of degree zero.

In particular, since $v \neq v'$, we have $v \in P \cup \{x', y', z'\}$. \square

Lemma 4.13. In the setting (I)—(V) and (X)—(XI), assume that $v = x_m \in P$. Then x'_m does not exist.

Proof. The primitive relation $r(Q)$ (see (X)) becomes $x + y + z = x_m$, and by Lemma 4.12, we have $G(\Sigma) \subset P \cup \{x, y, z, x', y', z', x'_m\}$.

Assume to the contrary that we have $x'_m \in G(\Sigma)$. Clearly, $x'_m \notin P$ because x_m forms a 2-dimensional cone in Σ with any other generator $x_i \in P$. By Remark 4.2, x'_m makes a primitive collection with two of x, y, z . Without loss of generality, assume we have a primitive relation of the form

$$x + y + x'_m = w.$$

Combining it with $r(Q)$ (see (X)), we get $x'_m + x_m = z + w$, so $w = z'$ by Lemma 2.26. Let $\alpha = x_m + x'_m = z + z'$. Then $\alpha \notin P$ because $x'_m = \alpha - x_m$ is not in $\text{Span } P$. Now substituting $z = \alpha - z'$ into $r(Q)$ (see (X)) yields

$$(8) \quad x + y + \alpha = x_m + z'.$$

Since $x_m \neq z$ by $r(Q)$ (see (X)), we have $\langle x_m, z' \rangle \in \Sigma$, so Equation (8) is an extremal primitive relation. This implies that $\alpha \neq x, y, z, x', y', z', x'_m$, a contradiction with Lemma 4.12. \square

Lemma 4.14. In the setting (I)—(V) and (X)—(XI), assume that $v = x_m \in P$. Then $\rho(X) \leq 3$.

Proof. By Lemma 4.12 and Lemma 4.13, we have $G(\Sigma) \subset P \cup \{x, y, z, x', y', z'\}$, and, as before, the primitive relation in (X) is $r(Q)$: $x + y + z = x_m$.

We show that at most one of x', y', z' exists. Suppose to the contrary that for example $x', y' \in G(\Sigma)$, and let $\alpha = x + x'$, $\beta = y + y' \in G(\Sigma)$. Since

$$\alpha + y + z = x_m + x' \quad \text{and} \quad \langle x_m, x' \rangle \in \Sigma,$$

applying Lemma 2.27 shows that this is an extremal primitive relation: indeed, either α and y are opponents and $z \in \{x_m, x'\}$, or α and z are opponents and $y \in \{x_m, x'\}$. But $y, z \notin P \cup \{x'\}$, so this is a contradiction. It follows that $\alpha \notin \{x, x', y, y', z, z'\}$, so $\alpha = x_l \in P$; similarly, $\beta = x_k \in P$. But we have

$$\alpha + \beta + z = x' + y' + x_m,$$

which is not possible since we show that the right hand side forms a cone. Indeed, applying Proposition 2.8 to $x + x' = x_l$ and $\tau = \langle x_m, x_k \rangle$ (we use here that $n \geq 5$), we obtain $\langle x_m, x_k, x' \rangle \in \Sigma$, and applying then Proposition 2.8 to $y + y' = x_k$ and $\tau = \langle x_m, x' \rangle$ we have that $\langle x_m, x', y' \rangle \in \Sigma$. So, after possibly relabelling x, y, z , we have $G(\Sigma) \subset P \cup \{x, y, z, x'\}$, hence $\rho(X) \leq 3$. \square

Lemma 4.15. In the setting (I)—(V) and (X)—(XI), assume that $v = x'$. Then $\rho(X) \leq 3$.

Proof. We have $r(Q): x+y+z = x'$. We know $G(\Sigma) \subset P \cup \{x, y, z, x', y', z'\}$ by Lemma 4.12. So it is enough to show y' and z' do not exist. Suppose to the contrary that, for example, $y' \in G(\Sigma)$, and let $\beta = y + y'$. Then

$$\beta + x + z = x' + y' \quad \text{and} \quad \langle x', y' \rangle \in \Sigma,$$

so again, by Lemma 2.27, this is an extremal primitive relation. Indeed, assume the relation is not primitive. By Lemma 2.27, either β and x are opponents and $z \in \{x', y'\}$, or β and z are opponents and $x \in \{x', y'\}$. But $z, x \notin \{x', y'\}$, since $\{x, y, z\}$ is a primitive collection. Hence $\beta + x + z = y' + x'$ is a primitive relation. It follows from Proposition 2.8 that $\langle x, x' \rangle \in \Sigma$, which is a contradiction. \square

Proposition 4.16. Let X be a projective toric Fano manifold of $\dim(X) = n \geq 5$, $m(X) \geq 3$, which admits a primitive relation $r(P): x_0 + x_1 + \cdots + x_{n-2} = 0$ (assumptions (I)–(III)). Assume that any primitive collection $\{x, y, z\}$ such that $x, y, z \in G(\Sigma) \setminus P$ and $0 \in \text{Conv}(\pi(x), \pi(y), \pi(z))$ (assumptions (IV)–(V)) has degree 2 as in (XI), and that there exists such a triple x, y, z as in (X). Then $\rho(X) \leq 3$.

Proof. The statement follows from Lemma 4.12, Lemma 4.14 and Lemma 4.15. \square

4.2. Second case: none of $\{x, y, z\}$ form a primitive collection.

Proposition 4.17. Let X be a toric Fano manifold of $\dim(X) = n \geq 5$, $m(X) \geq 3$, which admits a primitive relation $r(P): x_0 + x_1 + \cdots + x_{n-2} = 0$ (assumptions (I)–(III)). Assume in addition that none of the triples $\{x, y, z\} \subseteq G(\Sigma) \setminus P$ such that $0 \in \text{Conv}(\pi(x), \pi(y), \pi(z))$ (assumptions (IV)–(V)) form a primitive collection. Then $\rho(X) \leq 3$.

Before proving Proposition 4.17, we will formulate the following useful lemma.

Lemma 4.18. In the setting of Proposition 4.17, take any triple $\{x, y, z\} \subseteq G(\Sigma) \setminus P$ with $\langle x, y \rangle \in \Sigma$. Assume that $0 \in \text{Conv}(\pi(x), \pi(y), \pi(z))$, or equivalently, $\pi(z) \in \langle -\pi(x), -\pi(y) \rangle$. Then $z = x'$ or $z = y'$.

Proof. By Lemma 4.1, x, y, z do not span a cone, and by assumption, they do not form a primitive collection. So two of the vectors must not form a cone. Since we assumed that $\langle x, y \rangle \in \Sigma$, we must have $z = x'$ or $z = y'$. \square

Proof of Proposition 4.17. We select $x, y \in G(\Sigma)$ with $\langle x, y \rangle \in \Sigma$ and such that the cone generated by $\pi(x)$ and $\pi(y)$ is maximal among cones in $\mathbb{Q}^2 \simeq \mathbb{Q}^n / \Gamma$ coming from such pairs. If there is $z \in G(\Sigma)$ such that $\pi(z)$ is outside the cone $\langle \pi(x), \pi(y) \rangle$, then we show that $z = x'$ or $z = y'$. Indeed, the case $\pi(z) \in \langle -\pi(x), -\pi(y) \rangle$ is covered by Lemma 4.18; $\pi(z) \in \langle \pi(x) \rangle$ or $\pi(z) \in \langle \pi(y) \rangle$ cannot happen by an argument similar to Lemma 4.1; and in the remaining case, $\pi(z)$ is in $\langle \pi(x), -\pi(y) \rangle$ or $\langle -\pi(x), \pi(y) \rangle$, we use maximality of the cone generated by $\pi(x)$ and $\pi(y)$.

Let v be such that $\pi(v)$ is in the open half plane determined by $\text{Span } \pi(x)$ not containing $\pi(y)$. Up to relabelling of x, y , we can assume that such a v exists. If $v = x'$, then $x + x' = y'$, since $\pi(x + x')$ is non-zero and is outside of $\langle \pi(y), \pi(x) \rangle$. So

$$\langle \pi(x), \pi(y) \rangle \subset \langle -\pi(y), -\pi(x') \rangle \cup \langle -\pi(x'), -\pi(y') \rangle \cup \langle -\pi(y'), -\pi(x) \rangle.$$

It follows from Lemma 4.18 that $G(\Sigma) = P \cup \{x, y, x', y'\}$, which concludes this case.

If now $v = y'$, in case $\pi(v) \in \langle -\pi(x), -\pi(y) \rangle$, we notice that $y + y' = x'$ as above, and in case $\pi(v) \in \langle \pi(x), -\pi(y) \rangle$, by completeness, we find w such that $\pi(w) \in \langle -\pi(x), \pi(y) \rangle \cup \langle -\pi(x), -\pi(y) \rangle$ and notice $w = x'$. In either case, we get

$$\mathbb{Q}^2 = \langle -\pi(x), -\pi(y) \rangle \cup \langle -\pi(y), -\pi(x') \rangle \cup \langle -\pi(x'), -\pi(y') \rangle \cup \langle -\pi(y'), -\pi(x) \rangle,$$

and now Lemma 4.18 gives $G(\Sigma) = P \cup \{x, y, x', y'\}$. \square

5. TORIC FANO MANIFOLDS WITH $m(X) = n - 2$

In this section, we classify all n -dimensional toric Fano manifolds X with $m(X) = n - 2$ and $n \geq 6$ (Theorem 1.4). By Theorem 1.3, we know that $\rho(X) \leq 3$. Theorem 5.1 and Theorem 5.2 classify n -dimensional toric Fano manifolds X with $m(X) = n - 2$, $n \geq 5$, and Picard rank $\rho(X) = 2$ and 3, respectively. Together, these results yield a classification of toric Fano manifolds with $m(X) = n - 2$ and $n \geq 6$.

Theorem 5.1. Let X be a toric Fano manifold of dimension $n \geq 5$, $m(X) = n - 2$ and $\rho(X) = 2$. Then $X \simeq \mathbb{P}_{\mathbb{P}^2}(\mathcal{E})$, where \mathcal{E} is one of the following vector bundles on \mathbb{P}^2 :

- $\mathcal{E} = \mathcal{O}_{\mathbb{P}^2}(1) \oplus \mathcal{O}_{\mathbb{P}^2}^{\oplus n-2}$,
- $\mathcal{E} = \mathcal{O}_{\mathbb{P}^2}(1) \oplus \mathcal{O}_{\mathbb{P}^2}(1) \oplus \mathcal{O}_{\mathbb{P}^2}^{\oplus n-3}$,
- $\mathcal{E} = \mathcal{O}_{\mathbb{P}^2}(2) \oplus \mathcal{O}_{\mathbb{P}^2}^{\oplus n-2}$.

Proof. Let X be an n -dimensional toric Fano manifold with $m(X) = n - 2$ and $\rho(X) = 2$. We recall that toric manifolds with Picard rank 2 are classified by [Kle88]: they are projective space bundles over projective spaces. The assumption $m(X) = n - 2$ implies that X is a \mathbb{P}^{n-2} -bundle over \mathbb{P}^2 .

We can write $X = \mathbb{P}(\mathcal{E})$, where $\mathcal{E} = \mathcal{O}_{\mathbb{P}^2}(a_0) \oplus \mathcal{O}_{\mathbb{P}^2}(a_1) \oplus \cdots \oplus \mathcal{O}_{\mathbb{P}^2}(a_{n-2})$ and $a_0 \geq a_1 \geq \cdots \geq a_{n-2} = 0$. The Fano assumption on X is equivalent to saying that $\sum_{i=0}^{n-2} a_i \leq 2$. Thus we have the following cases:

- (1) $\mathcal{E} = \mathcal{O}_{\mathbb{P}^2}^{\oplus n-1}$ and $X \simeq \mathbb{P}^{n-2} \times \mathbb{P}^2$,
- (2) $\mathcal{E} = \mathcal{O}_{\mathbb{P}^2}(1) \oplus \mathcal{O}_{\mathbb{P}^2}^{\oplus n-2}$,
- (3) $\mathcal{E} = \mathcal{O}_{\mathbb{P}^2}(1) \oplus \mathcal{O}_{\mathbb{P}^2}(1) \oplus \mathcal{O}_{\mathbb{P}^2}^{\oplus n-3}$,
- (4) $\mathcal{E} = \mathcal{O}_{\mathbb{P}^2}(2) \oplus \mathcal{O}_{\mathbb{P}^2}^{\oplus n-2}$.

In Case (1), we have $m(X) = 2$. Cases (2)–(4) provide the complete list of toric Fano manifolds of Picard rank 2 and $m(X) = n - 2$. \square

Theorem 5.2. Let X be a toric Fano manifold of dimension $n \geq 5$, $m(X) = n - 2$ and $\rho(X) = 3$. Then one of the following holds:

- (1) $X = \mathbb{P}_S(\mathcal{E})$ is a \mathbb{P}^{n-2} -bundle over a toric surface S , where (S, \mathcal{E}) is one of the following:
 - $S = \mathbb{P}^1 \times \mathbb{P}^1$ and $\mathcal{E} = \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(1, 1) \oplus \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}^{\oplus n-2}$,
 - $S = \mathbb{P}^1 \times \mathbb{P}^1$ and $\mathcal{E} = \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(1, 0) \oplus \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(0, 1) \oplus \mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}^{\oplus n-3}$,
 - $S = \mathbb{F}_1$ and $\mathcal{E} = \mathcal{O}_{\mathbb{F}_1}(e + f) \oplus \mathcal{O}_{\mathbb{F}_1}^{\oplus n-2}$, where $e \subset \mathbb{F}_1$ is the -1 -curve, and $f \subset \mathbb{F}_1$ is a fiber of $\mathbb{F}_1 \rightarrow \mathbb{P}^1$.
- (2) Let $Y \simeq \mathbb{P}_{\mathbb{P}^2}(\mathcal{O}_{\mathbb{P}^2}(1) \oplus \mathcal{O}_{\mathbb{P}^2}^{\oplus n-2})$ be the blowup of \mathbb{P}^n along a linear subspace $L = \mathbb{P}^{n-3}$, and denote by $E \subset Y$ the exceptional divisor. Then X is the blowup of Y along a codimension 2 center $Z \subset Y$, where:

- Z is the intersection of E with the strict transform of a hyperplane of \mathbb{P}^n containing the linear subspace L , or
- Z is the intersection of the strict transforms of two hyperplanes of \mathbb{P}^n , one containing the linear subspace L , and the other one not containing it.

Proof. We apply Batyrev's classification of toric manifolds with $\rho(X) = 3$, stated in Proposition 2.18. We adopt the same notation as in Proposition 2.18, and treat separately the cases when the number of primitive collections is $l = 3$ and $l = 5$.

Case $l = 3$. As $G(\Sigma) = X_0 \sqcup X_1 \sqcup X_2$ and $m(X) = n - 2$, we have $X_0 = \{x_0, \dots, x_{n-2}\}$, $X_1 = \{v_1, v_2\}$ and $X_2 = \{z_1, z_2\}$. The corresponding primitive relations are all extremal by Proposition 2.18. By Proposition 2.14, X is a \mathbb{P}^{n-2} -bundle over a surface S . Up to relabelling in X_0 , X_1 and X_2 , three possible choices for the remaining two primitive relations are (noting that $m(X) = n - 2 > 1$):

$$\begin{cases} v_1 + v_2 = x_0, \\ z_1 + z_2 = v_1; \end{cases} \quad \begin{cases} v_1 + v_2 = x_0, \\ z_1 + z_2 = x_0; \end{cases} \quad \begin{cases} v_1 + v_2 = x_0, \\ z_1 + z_2 = x_1. \end{cases}$$

It follows that S is isomorphic to \mathbb{F}_1 when $r(X_1)$ and $r(X_2)$ are as in the first column above, or to $\mathbb{P}^1 \times \mathbb{P}^1$ in two other cases.

Case $l = 5$. We denote by $l_i = |X_i| \geq 1$ the cardinality of X_i .

If $(\bar{c}, \bar{b}) = (\bar{0}, \bar{0})$, then both $r_2: \bar{1} \cdot X_2 + \bar{1} \cdot X_3 = 0$ and $r_4: \bar{1} \cdot X_4 + \bar{1} \cdot X_0 = 0$ are centered primitive relations. By the assumption $m(X) = n - 2$, we have $2n - 2 \leq l_2 + l_3 + l_4 + l_0 \leq |G(\Sigma)| - 1 = n + 2$, which implies $n \leq 4$, a contradiction.

So we have $(\bar{c}, \bar{b}) \neq (\bar{0}, \bar{0})$, and $P := X_2 \sqcup X_3$ is the only centered primitive collection, so $l_2 + l_3 = n - 1$ and $l_0 + l_1 + l_4 = 4$, with $l_0, l_1, l_4 \in \{1, 2\}$. As $\deg(r_0) > 0$, we get the inequality

$$3 \geq l_0 + l_1 > \sum c_i + \sum b_j + l_3,$$

which is only satisfied when $l_3 = 1$, $l_0 + l_1 = 3$ and exactly one entry in

$$(c_1, c_2, \dots, c_{l_2}, b_1)$$

equals one, while the others are all zero. Up to relabelling, there are two cases: $c_1 = 1$ or $b_1 = 1$.

We then have $l_4 = 4 - (l_0 + l_1) = 1$. From $\deg(r_3) > 0$, we get

$$l_3 + l_4 > l_1,$$

which means $2 > l_1$, and this ensures $l_1 = 1$, hence $(l_0, l_1, l_4) = (2, 1, 1)$.

To sum up, we denote $X_0 = \{v_1, v_2\}$, $X_1 = \{y\}$, $X_2 = \{x_1, \dots, x_{n-2}\}$, $X_3 = \{x_0\}$ and $X_4 = \{z\}$, and get the following two possibilities for X :

- $b_1 = 1$ and $c_j = 0$ for every j :

$$\begin{aligned} r_0: & \quad v_1 + v_2 + y = 2x_0, \\ r_1: & \quad y + x_1 + \dots + x_{n-2} = z, \\ r_2: & \quad x_0 + \dots + x_{n-2} = 0, \\ r_3: & \quad x_0 + z = y, \\ r_4: & \quad z + v_1 + v_2 = x_0. \end{aligned}$$

- $c_1 = 1$, $b_1 = 0$ and $c_j = 0$ for every $j > 1$:

$$\begin{aligned} r_0: \quad & v_1 + v_2 + y = x_1 + x_0, \\ r_1: \quad & y + x_1 + \cdots + x_{n-2} = z, \\ r_2: \quad & x_0 + \cdots + x_{n-2} = 0, \\ r_3: \quad & x_0 + z = y, \\ r_4: \quad & z + v_1 + v_2 = x_1. \end{aligned}$$

By Proposition 2.14, the open $U = X \setminus (V(y) \cup V(z))$ has a \mathbb{P}^{n-2} -bundle structure. The relation r_3 corresponds to an extremal curve class by Proposition 2.21, which induces a smooth blow-down $h: X \rightarrow Y$. By Proposition 2.17, the primitive relations in Y in cases $b_1 = 1$ and $c_1 = 1$ are, respectively:

$$\begin{cases} x_0 + \cdots + x_{n-2} = 0, \\ z + v_1 + v_2 = x_0; \end{cases} \quad \begin{cases} x_0 + \cdots + x_{n-2} = 0, \\ z + v_1 + v_2 = x_1. \end{cases}$$

It follows that $Y \simeq \mathbb{P}(\mathcal{O}_{\mathbb{P}^2}(1) \oplus \mathcal{O}_{\mathbb{P}^2}^{\oplus n-2})$, i.e., Y is the blowup of \mathbb{P}^n along a linear subspace $L = V(z, v_1, v_2) \simeq \mathbb{P}^{n-3} \subset \mathbb{P}^n$, with exceptional divisor $E = V(x_0) \subset Y$ in the first case, and $E = V(x_1) \subset Y$ in the second case. The center of the blowup $h: X \rightarrow Y$ is $V(x_0, z) \subset Y$, yielding the two varieties described in (2). \square

APPENDIX A. CODE FOR COMPUTING PRIMITIVE COLLECTIONS

by WILL REYNOLDS

This appendix explains the algorithm used by its author to compute the primitive collections of the smooth toric Fano varieties up to dimension 6. This algorithm was implemented in Macaulay2 ([GS]), making use of the NormalToricVarieties package ([Smi]), which includes a database of all of the smooth toric Fano varieties up to dimension 6. It is also possible to compute primitive collections using the software package Polymake ([GJ00]) and its command `MINIMAL_NON_FACES`.

Let Σ be the fan of a toric manifold of dimension n . For each integer $k \in \{1, \dots, n\}$, we denote by $\Sigma(k)$ the subset of Σ consisting of k -dimensional cones. To determine whether a given subset $P \subseteq G(\Sigma)$ is a primitive collection, we proceed in two steps. First make sure there does not exist $\sigma \in \Sigma(n)$ with $P \subseteq G(\sigma)$, and if there does, stop; otherwise, for each $v \in P$, make sure there exists $\sigma \in \Sigma(n)$ with $P \setminus \{v\} \subseteq G(\sigma)$. In the special case $|P| = 2$ it suffices to check that there does not exist $\sigma \in \Sigma(n)$ with $P \subseteq G(\sigma)$.

Note that, by definition, if P is a primitive collection and $P \subsetneq Q$, then Q is not a primitive collection. Therefore, when looking for primitive collections, we go through subsets of $G(\Sigma)$ in increasing cardinality. With the above in place, a reasonably efficient way to list all of the primitive collections of a fan is to arrive at such a list by eliminating $P \subseteq G(\Sigma)$ which are not primitive collections. The first step is to remove any P with $|P| = 1$. Then for each P we check whether it is a primitive collection: if it is, keep it and remove all sets properly containing it; if it is not, remove it.

This (straightforward) algorithm is impractical if $G(\Sigma)$ is too large, and this is the main reason we ran it only up to dimension 6. In practice, at the time these computations were

carried out, runtime remained fairly reasonable up to about $|G(\Sigma)| = 17$. Part of the reason such a direct approach remained feasible to this extent is perhaps of wider interest: the relative abundance of primitive collections of size 2, at least among toric Fano varieties, means that eliminating all of the supersets of any P with $|P| = 2$ often cuts down the search space quite significantly. For example, the 124 toric Fano 4-folds altogether have 785 primitive collections, of which 566 have cardinality 2. This curious abundance makes the computation of $m(X)$ for a given toric Fano variety X easier as well. Of the toric Fano varieties of a given dimension n (for $n \leq 6$), those with $m(X) = 1$ make up an overwhelming majority. Hence, even the most direct way of computing $m(X)$ is usually extremely fast. Table 1 in the Introduction summarizes the number of toric Fano manifolds according to the value $m(X)$ for $\dim(X) \in \{4, 5, 6\}$.

For convenience, we also provide Macaulay2 code implementing the algorithm for computing primitive collections.

```

coneExistenceCheck = (S, fan) -> (
  for cone in fan do (
    if isSubset(S, cone) then (
      return true;
    );
  );
  return false;
);

properSubsetCheck = (S, fan) -> (
  for ray in S do (
    if coneExistenceCheck(S-set{ray}, fan) == false then (
      return false;
    );
  );
  return true;
);

isPrimitiveCollection = (P, Var) -> (
  if coneExistenceCheck(P, orbits(Var, 0)) then (
    return false
  )
  else (
    return properSubsetCheck(P, orbits(Var, 0))
  );
);

supsetsOfPrimColl = (E, B) -> (
  return set{for P in E-set{B} when isSubset(B, P) list P};
);

primitiveCollections = (Var) -> (
  n := length rays Var;

```

```

primColls := select(subsets(toList(0..n-1)), x -> length x > 1);
for P in subsets(toList(0..n-1), 2) do (
  if coneExistenceCheck(P, orbits(Var, 0)) == false then (
    primColls = primColls - supsetsOfPrimColl(primColls, P);)
  else (
    primColls = primColls - set{P};
  );
);
for i in toList(3..n) do (
  for P in subsets(toList(0..n-1), i) do (
    if member(P, primColls) == false then continue;
    if isPrimitiveCollection(P, Var) then (
      primColls = primColls - supsetsOfPrimColl(primColls, P);
    ) else (
      primColls = primColls - set{P};
    );
  );
);
return sort primColls;
);

```

REFERENCES

- [AC12] C. Araujo and A.-M. Castravet. “Polarized minimal families of rational curves and higher Fano manifolds”. In: *Amer. J. Math.* 134.1 (2012), pp. 87–107. ISSN: 0002-9327.
- [AC13] C. Araujo and A.-M. Castravet. “Classification of 2-Fano manifolds with high index”. In: *A celebration of algebraic geometry*. Vol. 18. Clay Math. Proc. Amer. Math. Soc., Providence, RI, 2013, pp. 1–36.
- [Ara+22] C. Araujo, R. Beheshti, A.-M. Castravet, K. Jabbusch, S. Makarova, E. Mazzon, L. Taylor, and N. Viswanathan. “Higher Fano manifolds”. In: *Rev. Un. Mat. Argentina* 64.1 (2022), pp. 103–125. ISSN: 0041-6932.
- [Bat91] V. V. Batyrev. “On the classification of smooth projective toric varieties”. In: *Tohoku Math. J. (2)* 43.4 (1991), pp. 569–585. ISSN: 0040-8735.
- [Bat99] V. V. Batyrev. “On the classification of toric Fano 4-folds”. In: vol. 94. 1. *Algebraic geometry*, 9. 1999, pp. 1021–1050.
- [BK] G. Brown and A. Kasprzyk. *Graded Ring Database*. Available at <http://www.grdb.co.uk/forms/toricsmooth>.
- [BW23] R. Beheshti and B. Wormleighton. “Bounds on the Picard rank of toric Fano varieties with minimal curve constraints”. In: *Proc. Amer. Math. Soc.* 151.4 (2023), pp. 1503–1518. ISSN: 0002-9939,1088-6826.
- [Cam92] F. Campana. “Connexité rationnelle des variétés de Fano”. In: *Ann. Sci. École Norm. Sup. (4)* 25.5 (1992), pp. 539–545. ISSN: 0012-9593.

- [Cas03a] C. Casagrande. “Contractible classes in toric varieties”. In: *Math. Z.* 243.1 (2003), pp. 99–126. ISSN: 0025-5874.
- [Cas03b] C. Casagrande. “Toric Fano varieties and birational morphisms”. In: *Int. Math. Res. Not.* 27 (2003), pp. 1473–1505. ISSN: 1073-7928.
- [CD15] C. Casagrande and S. Druel. “Locally unsplit families of rational curves of large anticanonical degree on Fano manifolds”. In: *Int. Math. Res. Not. IMRN* 21 (2015), pp. 10756–10800. ISSN: 1073-7928.
- [CFH14] Y. Chen, B. Fu, and J.-M. Hwang. “Minimal rational curves on complete toric manifolds and applications”. In: *Proc. Edinb. Math. Soc. (2)* 57.1 (2014), pp. 111–123. ISSN: 0013-0915.
- [CLS11] D. A. Cox, J. B. Little, and H. K. Schenck. *Toric varieties*. Vol. 124. Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 2011, pp. xxiv+841. ISBN: 978-0-8218-4819-7.
- [dHS11] A. J. de Jong, X. He, and J. M. Starr. “Families of rationally simply connected varieties over surfaces and torsors for semisimple groups”. In: *Publ. Math. Inst. Hautes Études Sci.* 114 (2011), pp. 1–85. ISSN: 0073-8301.
- [dS06a] A. J. de Jong and J. M. Starr. “A note on Fano manifolds whose second Chern character is positive”. In: *arXiv Mathematics e-prints* (Feb. 2006). arXiv: [math/0602644](https://arxiv.org/abs/math/0602644) [math.AG].
- [dS06b] A. J. de Jong and J. M. Starr. *A note on Fano manifolds whose second Chern character is positive*. arXiv:math/0602644. 2006. arXiv:math: [0602644](https://arxiv.org/abs/math/0602644) (math.AG).
- [dS06c] A. J. de Jong and J. M. Starr. *Low degree complete intersections are rationally simply connected*. 2006.
- [dS07] A. J. de Jong and J. Starr. “Higher Fano manifolds and rational surfaces”. In: *Duke Math. J.* 139.1 (2007), pp. 173–183. ISSN: 0012-7094.
- [Ful93] W. Fulton. *Introduction to toric varieties*. Vol. 131. Annals of Mathematics Studies. The William H. Roever Lectures in Geometry. Princeton University Press, Princeton, NJ, 1993, pp. xii+157. ISBN: 0-691-00049-2.
- [GJ00] E. Gawrilow and M. Joswig. “Polymake: a framework for analyzing convex polytopes”. In: *Polytopes—combinatorics and computation*. Springer. 2000, pp. 43–73.
- [GS] D. R. Grayson and M. E. Stillman. *Macaulay2, a software system for research in algebraic geometry*. Available at <http://www.math.uiuc.edu/Macaulay2/>.
- [Kle88] P. Kleinschmidt. “A classification of toric varieties with few generators”. In: *aequationes mathematicae* 35.2 (1988), pp. 254–266. ISSN: 1420-8903.
- [KMM92] J. Kollár, Y. Miyaoka, and S. Mori. “Rational connectedness and boundedness of Fano manifolds”. In: *J. Differential Geom.* 36.3 (1992), pp. 765–779. ISSN: 0022-040X.
- [Kol96] J. Kollár. *Rational curves on algebraic varieties*. Vol. 32. Ergebnisse der Mathematik und ihrer Grenzgebiete. Berlin: Springer-Verlag, 1996.
- [Mor79] S. Mori. “Projective manifolds with ample tangent bundles”. In: *Ann. of Math. (2)* 110.3 (1979), pp. 593–606. ISSN: 0003-486X.
- [Nag19] T. Nagaoka. “On a sufficient condition for a Fano manifold to be covered by rational N -folds”. In: *J. Pure Appl. Algebra* 223.11 (2019), pp. 4677–4688. ISSN: 0022-4049.

- [Nob11] E. E. Nobili. “Classification of Toric 2-Fano 4-folds”. In: *Bulletin of the Brazilian Mathematical Society, New Series* 42.3 (2011), p. 399. ISSN: 1678-7714.
- [Nob12] E. E. Nobili. *Birational Geometry of Toric Varieties*. Thesis (Ph.D.) 2012.
- [Øbr07] M. Øbro. “Classification of terminal simplicial reflexive d -polytopes with $3d - 1$ vertices”. In: *manuscripta mathematica* 125.1 (Aug. 2007), pp. 69–79. ISSN: 1432-1785.
- [Rei83] M. Reid. “Decomposition of Toric Morphisms”. In: *Arithmetic and Geometry: Papers Dedicated to I.R. Shafarevich on the Occasion of His Sixtieth Birthday. Volume II: Geometry*. Ed. by M. Artin and J. Tate. Boston, MA: Birkhäuser Boston, 1983, pp. 395–418. ISBN: 978-1-4757-9286-7.
- [Sat00] H. Sato. “Toward the classification of higher-dimensional toric Fano varieties”. In: *Tohoku Math. J. (2)* 52.3 (2000), pp. 383–413. ISSN: 0040-8735.
- [Sat12] H. Sato. “The numerical class of a surface on a toric manifold”. In: *Int. J. Math. Math. Sci.* (2012), Art. ID 536475, 9. ISSN: 0161-1712.
- [Sat16] H. Sato. “Toric 2-Fano manifolds and extremal contractions”. In: *Proc. Japan Acad. Ser. A Math. Sci.* 92.10 (2016), pp. 121–124. ISSN: 0386-2194.
- [Shr20] M. Shrieve. *On the γ_2 -positivity of Smooth Toric Threefolds*. Thesis (Ph.D.)—University of Washington. ProQuest LLC, Ann Arbor, MI, 2020, p. 54. ISBN: 979-8684-67090-9.
- [Smi] G. G. Smith. *NormalToricVarieties: A Macaulay2 package. Version 1.8*. A *Macaulay2* package available at <https://github.com/Macaulay2/M2/tree/master/M2/Macaulay2/packages>.
- [SS20] H. Sato and Y. Suyama. “Remarks on toric manifolds whose Chern characters are positive”. In: *Comm. Algebra* 48.6 (2020), pp. 2528–2538. ISSN: 0092-7872.
- [SSS21] Y. Sano, H. Sato, and Y. Suyama. “Toric Fano manifolds of dimension at most eight with positive second Chern characters”. In: *Kumamoto J. Math.* 34 (2021), pp. 1–13.
- [Sta06] J. M. Starr. *Hypersurfaces of low degree are rationally simply-connected*. 2006. arXiv:math: [0602641](https://arxiv.org/abs/0602641) (math.AG).
- [Suz21] T. Suzuki. “Higher order minimal families of rational curves and Fano manifolds with nef Chern characters”. In: *J. Math. Soc. Japan* 73.3 (2021), pp. 949–964. ISSN: 0025-5645.

CAROLINA ARAUJO, IMPA, ESTRADA DONA CASTORINA 110, 22460-320 RIO DE JANEIRO, BRAZIL
Email address: caraujo@impa.br

ROYA BEHESHTI, DEPARTMENT OF MATHEMATICS & STATISTICS, WASHINGTON UNIVERSITY IN ST. LOUIS, ST. LOUIS, MO, 63130
Email address: beheshti@wustl.edu

ANA-MARIA CASTRAVET, UNIVERSITÉ PARIS-SACLAY, UVSQ, LABORATOIRE DE MATHÉMATIQUES DE VERSAILLES, 45 AVENUE DES ÉTATS UNIS, 78035 VERSAILLES, FRANCE
Email address: ana-maria.castravet@uvsq.fr

KELLY JABBUSCH, DEPARTMENT OF MATHEMATICS & STATISTICS, UNIVERSITY OF MICHIGAN-DEARBORN, 4901 EVERGREEN RD, DEARBORN, MICHIGAN, 48128, USA
Email address: jabbusch@umich.edu

SVETLANA MAKAROVA, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF PENNSYLVANIA, 209 S 33RD ST, PHILADELPHIA, PA 19104, USA
Email address: murmuno@gmail.com

ENRICA MAZZON, FAKULTÄT FÜR MATHEMATIK, UNIVERSITÄT REGENSBURG, UNIVERSITÄTSSTRASSE 31, 93040 REGENSBURG, GERMANY, AND DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MICHIGAN, 48109, USA
Email address: e.mazzon15@alumni.imperial.ac.uk

NIVEDITA VISWANATHAN, SCHOOL OF MATHEMATICS, CENTER FOR ENGINEERING, DESIGN AND PHYSICAL SCIENCES, BRUNEL UNIVERSITY LONDON, KINGSTON LANE, UXBRIDGE, MIDDLESEX UB8 3PH, UNITED KINGDOM
Email address: Nivedita.Viswanathan@brunel.ac.uk

WILL REYNOLDS, SCHOOL OF MATHEMATICS, THE UNIVERSITY OF EDINBURGH, EDINBURGH, EH9 3FD, UK
Email address: W.R.N.Reynolds@sms.ed.ac.uk