

**Running Title:** The effects of elastic kinesiology tape on shoulder proprioception

# Effects of Elastic Kinesiology Taping on Shoulder Proprioception: a Systematic Review

Amanda L. Ager,<sup>1,2</sup> Fabio Carlos Lucas de Oliveira,<sup>3</sup> Jean-Sébastien Roy,<sup>1,4</sup> Dorien Borms,<sup>2</sup>  
Michiel Deraedt,<sup>2</sup> Morgane Huyge,<sup>2</sup> Arne Deschepper,<sup>2</sup> Ann M. Cools<sup>2</sup>

<sup>1</sup>*Center for Interdisciplinary Research in Rehabilitation and Social Integration, Rehabilitation Institute, Quebec City, QC, Canada*

<sup>2</sup>*Department of Rehabilitation Sciences and Physiotherapy, Faculty of Medicine and Health Sciences, Ghent University, Ghent, Belgium*

<sup>3</sup>*Research Unit in Sport and Physical Activity (CIDAF), Faculty of Sport Sciences and Physical Education, University of Coimbra, Coimbra, Portugal.*

<sup>4</sup>*Department of Rehabilitation, Faculty of Medicine, Université Laval, Quebec City, QC, Canada*

## Correspondence to:

**Amanda L. Ager,**

Address: Campus UZ Ghent, Corneel Heymanslaan 10, B3, ingang 46, 9000 Ghent

Email: [Amanda.Ager@UGent.be](mailto:Amanda.Ager@UGent.be)

Received: 23 January 2022; Revised: 03 August 2022; Accepted: 13 April 2023

Systematic Review

26 **Supplementary Material**

- 27 • Key terms and MeSH strategy employed during the literature review, based on the PICOS question
- 28 • Tools used for quality assessments of the included studies, classification of the risk of bias and quality of
- 29 the studies
- 30 • Grades of evidence used to assign the certainty of evidence per shoulder proprioception outcome
- 31 • Summary of key findings from the included studies
- 32 • Visual depiction of employed elastic KT protocols of included studies
- 33 • List of excluded studies after full-text reading

34

35

36

37

38

39

40

41

42 **ABSTRACT**

43 **Background:** Shoulder injuries are associated with proprioceptive deficits. Elastic kinesiology tape  
44 (KT) is used for treating musculoskeletal disorders, including shoulder injuries, as it arguably  
45 improves proprioception.

46 **Objective:** To synthesize the evidence on the effects of elastic KT on proprioception in healthy and  
47 pathological shoulders.

48 **Methods:** Four databases (PubMed, WoS, CINAHL, SPORTDiscus) were searched for studies  
49 investigated the effects of elastic KT on shoulder proprioception. Outcome measures were active joint  
50 position sense (AJPS), passive joint position sense (PJPS), kinesthesia, sense of force (SoF), and  
51 velocity (SoV). Risk of bias (RoB) was assessed using the Cochrane Collaboration RoB tool for  
52 randomized controlled trials (RCTs), and the ROBINS-1 for non-RCTs, while the certainty of  
53 evidence was determined using GRADE.

54 **Results:** Eight studies (5 RCTs, 3 non-RCTs) were included, yielding 174 participants (102 healthy  
55 and 85 pathological shoulders). RoB ranged from low (2 studies), moderate (5 studies), to high (1  
56 study). Elastic KT has a mixed effect on AJPS of healthy shoulders (n=79) (low certainty). Elastic  
57 KT improves AJPS (subacromial pain syndrome and rotator cuff tendinopathy, n=52) and PJPS  
58 (chronic hemiparetic shoulders, n=13) among pathological shoulders (very low certainty). Elastic KT  
59 has no effect on kinesthesia among individuals with subacromial pain syndrome (n=30) (very low  
60 certainty).

61 **Conclusion:** There is very low to low certainty of evidence that elastic KT enhances shoulder AJPS  
62 and PJPS. The aggregate of evidence is currently so low that any recommendation on the  
63 effectiveness of elastic KT on shoulder proprioception remains speculative.

64  
65 **Keywords:** elastic taping; joint position sense; kinesthesia; proprioception; shoulder; upper limbs.

66  
67 **Highlights**

- 68
- 69 • Elastic kinesiology tape (KT) has a mixed effect on active joint position sense (AJPS) of  
70 healthy shoulders (low certainty).
  - 71 • Elastic KT improves active or passive JPS among pathological shoulders (very low certainty).
  - 72 • Elastic KT has no effect on kinesthesia with subacromial pain syndrome (very low certainty).
  - 73 • Overall, evidence remains speculative as to the effects of elastic KT on shoulder  
74 proprioception.
- 75

76 **INTRODUCTION**

77 The importance of shoulder pain within orthopaedic medicine and rehabilitation has been well  
78 established, with an estimated 30-50% of adults experiencing at least one episode of shoulder pain  
79 annually.<sup>1</sup> While being a meaningful reason to seek medical care, shoulder pain continues to affect a  
80 person's ability to work and their capacity to participate in activities of daily living;<sup>2</sup> while being a  
81 costly problem to the individual and society.<sup>3,4</sup> In search of more effective treatments, elastic  
82 kinesiology taping (KT) has often been used in clinical practice<sup>5</sup> as an additional therapeutic resource  
83 for treating shoulder pain. Kinesiology or elastic taping, also referred to as neuro-proprioceptive  
84 taping,<sup>6</sup> is a popular clinical tool theorized to improve proprioception.<sup>5,7-11</sup> It is described as being  
85 therapeutic<sup>7</sup> with a wide range of theoretical benefits, including (i) mimicking the elasticity of skeletal  
86 muscle<sup>12,13</sup> while allowing unrestricted range of motion (ROM);<sup>12,13</sup> (ii) improving sensory  
87 mechanisms,<sup>12</sup> correcting muscle function,<sup>5</sup> and facilitating motor activity;<sup>8</sup> (iii) the promotion of a  
88 neutral postural alignment and joint stability<sup>8</sup> and; (iv) decreased pain through neurological  
89 suppression.<sup>5,7,8</sup> Despite the widespread application of elastic KT in clinical practice,<sup>14,15</sup> its scientific  
90 effectiveness remains unclear,<sup>16,17</sup> particularly as it applies to shoulder proprioception.

91 Proprioception, or our limb's sensory awareness,<sup>18</sup> provides essential guidance to the shoulder  
92 through feedback regarding positioning in space (joint position sense, JPS), movement (kinesthesia),  
93 sense of force (SOF) (or sense of effort),<sup>19</sup> and sense of joint velocity (SOV).<sup>20</sup> Collectively,  
94 proprioception is essential to shoulder neuromuscular control throughout movements of the  
95 inherently unstable glenohumeral (GH) joint,<sup>21</sup> while also playing a crucial role in our daily lives by  
96 guiding our interactions with the world around us.<sup>22</sup> It is also well established that proprioception  
97 contributes to sports performance and complex tasks of daily living.<sup>23</sup> Therefore, improving shoulder  
98 proprioception is an important clinical goal following an injury.

99 The use of elastic KT in rehabilitation is thought to improve neuromuscular control,<sup>5,24</sup> enhance  
100 postural alignment by aiding in repositioning the humeral head within the glenoid fossa, increasing  
101 the subacromial space,<sup>5</sup> and also correcting scapular positioning.<sup>25</sup> The application of elastic KT to  
102 the skin is suggested to improve proprioception via the stimulation of local cutaneous  
103 mechanoreceptors and proprioceptors within surrounding tissues,<sup>7,10</sup> collecting mechanical  
104 information on tissue deformation (stretch, tension, vibration, movement, and positioning).<sup>18</sup>

105 Previous systematic reviews have been published regarding the use of elastic KT to manage  
106 musculoskeletal injuries;<sup>26</sup> more specifically low back pain,<sup>27</sup> patellofemoral pain syndrome,<sup>28</sup> ankle  
107 instabilities,<sup>29</sup> rotator cuff tendinopathies,<sup>30</sup> as well as among overhead athletes.<sup>31</sup> However, most  
108 reviews did not address the effects of elastic KT specifically on proprioception. Recently, Turgut et  
109 al.<sup>31</sup> evaluated the effects of all types of taping (rigid tape, elastic taping, or a combination thereof)  
110 on shoulder proprioception of overhead athletes. The authors reported minor improvements to

111 shoulder proprioception and suggested mixed results and insufficient evidence for the effects of  
112 elastic KT on shoulder proprioception. Their results pinpoint a strong need for a review of the  
113 literature addressing the specific effects of elastic KT on shoulder proprioception. Indeed, despite the  
114 wide application of elastic KT, there is insufficient evidence to suggest that it directly affects  
115 proprioception,<sup>27,28,30-32</sup> except among individuals affected by ankle instabilities.<sup>29</sup> To our knowledge,  
116 a critical literature review has yet to address the effects of elastic KT on shoulder proprioception.  
117 Therefore, this study aims to review and synthesize the evidence of the impact of elastic KT on  
118 shoulder proprioception in healthy and pathological shoulders.

119 **METHODS**

120 **Identification and selection of trials**

121 Four databases (PubMed, Web of Science, CINAHL, and SPORTDiscus) were systematically  
122 searched from their inception until December 1<sup>st</sup> 2021, to identify articles that investigated the effects  
123 of elastic KT, primarily or secondarily, in both healthy and pathological populations on shoulder  
124 proprioception; including JPS, kinesthesia, SOF, and SOV. A search strategy using PICOS  
125 (Population, Intervention, Comparison, Outcome, Study design) approach, was performed without  
126 date, geographical location, gender, sex, or language restrictions. The search was tailored for each  
127 database using their specific building block, truncation, Boolean operators, and nesting features for  
128 combining medical subject heading (MeSH) and free-text words. Details from the search strategy are  
129 available in the Supplementary Material – Table S1.

130 The selection of the articles followed the Preferred Reporting Items for Systematic Reviews and  
131 Meta-Analyses (PRISMA) guidelines.<sup>33</sup> The search yield was exported to EndNote, and, after  
132 removing duplicates, titles and abstracts of the pre-selected studies were screened by three  
133 independent reviewers (M.D., M.H., A.D.). For a double-blinded process, potentially eligible studies  
134 were randomly assigned to a pair of reviewers of a three-member blinded team working in three pairs  
135 (M.D./A.D., M.D./M.H., A.D./M.H.).

136

137 ***Eligibility criteria***

138 To be selected for full-text screening, the article had to be a (i) RCT or a non-RCT studies of  
139 intervention investigating the effectiveness of elastic KT at the shoulder amongst healthy adult (18  
140 and 65 years old) or symptomatic individuals with any painful shoulder condition; (ii) report at least  
141 one shoulder proprioception outcome measure (JPS/kinesthesia/SoF/SoV); and (iii) be published in  
142 English, French, or Dutch. The same three pairs of reviewers scrutinized the full-text to determine  
143 their inclusion in this review, resulting in two independent reviewers per citation. Screening results  
144 were openly discussed until a unanimous consensus was reached. Manual searching on the reference  
145 lists was conducted to find additional articles not found in the previous bibliographical searches.

146

147 **Assessment of characteristics of trials**

148 ***Risk of bias***

149 The risk of bias (RoB) of the included studies was assessed using two assessment tools: the Cochrane  
150 Collaboration Risk of Bias tool (ROB 2) for RCTs (Table 1),<sup>34, 35</sup> and the Risk of Bias in Non-  
151 randomized Studies - of Interventions (ROBINS-I) for non-RCTs (Table 2). Details for both tools,  
152 are available in the Supplementary Material - Table S2.

153 The quality of the non-RCTs assessed with ROBINS-I tools was quantified based on the overall  
154 scores. As the summary scores for quantifying the quality of the studies assessed with the ROBINS-  
155 I checklist are not yet associated with qualitative categories, the following index, suggested by de  
156 Oliveira et al.,<sup>36</sup> was used. A study was deemed “high quality” (HQ) for scores greater than 80.0%,  
157 “good quality” (GQ) for scores between 70% and 80.0%, “moderate quality” (MQ) for scores  
158 between 50.0% and 69.9%, and “low quality” (LQ) for scores less than 50.0%.<sup>36</sup> This quality  
159 assessment index allowed us to evaluate the quality of the included studies categorically, based on  
160 proprioception outcome measures.

161

### 162 *Certainty of evidence for proprioception outcomes*

163 Two independent reviewers (A.L.A., F.O.) evaluated the included studies according to the grading of  
164 recommendations assessment, development and evaluation (GRADE) framework,<sup>37-39</sup> to establish the  
165 certainty of evidence regarding the effectiveness of elastic KT on shoulder proprioception outcomes  
166 among healthy and pathological shoulders. The evaluation of the evidence took into consideration  
167 five key domains: i) study design limitations; ii) results inconsistency; iii) indirectness of evidence;  
168 iv) imprecision, and v) publication bias.

169

170 The body of evidence for an outcome may be determined to have serious (downgraded one point),  
171 very serious (downgraded two points), or critically serious (downgraded three points, for RoB when  
172 ROBINS-I was used) issues for each domain ([Supplementary material - Table S3](#)). The GRADE  
173 quality of evidence was based on the following:

174

- 175 • **High quality:** Further research is very unlikely to change our confidence in the estimate of  
176 effect. Consistent findings among 75% of pooled participants in RCTs and non-RCTs of  
177 intervention with low RoB are generalizable to the population in question. Sufficient data,  
178 with narrow confidence intervals, are available. No reporting biases are known or suspected  
179 (all domains are met).
- 180 • **Moderate quality:** Further research is likely to have an important impact on our confidence  
181 in the estimate of effect and may change the estimate (one domain is not met).
- 182 • **Low quality:** Further research is very likely to have an important impact on our confidence  
183 in the estimate of effect and is likely to change the estimate (two domains are not met).
- 184 • **Very low quality:** We are very uncertain about the estimate (three domains are not met).
- 185 • **No evidence:** We identified no RCT or non-RCT of intervention that measured the outcome.

186

187 ***Data extraction***

188 The following data were systematically extracted by a three-member blinded team working in three  
189 pairs (M.D./A.D., M.D./M.H., A.D./M.H.). from the included studies: author and year of publishing,  
190 study design, sample/population, intervention/taping (type, application, technique used) and the  
191 control group, proprioception outcome measures, and overall results (see the Supplementary Material  
192 – Table S4).

193

194 ***Data analysis***

195 Because of the heterogeneity of the studies included in this review (e.g., differing populations,  
196 shoulder taping protocols, and proprioception outcome measures), the data could not be pooled into  
197 a meta-analysis. Therefore, only qualitative analyses were performed.

198

199



200 **RESULTS**

201 **Flow of trials through the review**

202 The literature search yielded 261 citations. After removing 69 duplicates, 192 remaining citations  
203 were screened, and eight studies were included in this review (Figure 1). Of the eight studies,<sup>5,8,11,12,</sup>  
204 <sup>40-42</sup> five were RCTs,<sup>11, 12, 40-42</sup> and three were non-RCTs studies of intervention.<sup>5,8,9</sup>

205

206 **Participants**

207 A total of 174 participants (88 men, 86 women), including 187 shoulders (102 healthy<sup>9,11,40,41</sup> and 85  
208 pathological shoulders<sup>5,8,41,42</sup>) were included. The mean (SD) age of the participants was 27.9 (3.9)  
209 years, and 49.4% were male. The diagnosis of the pathological shoulders included subacromial pain  
210 syndrome (SAPS, n=50)<sup>8, 42</sup> (including overhead athletes with SAPS [n=30])<sup>42</sup> and rotator cuff  
211 tendinopathy (n=22),<sup>5</sup> and chronic hemiparesis following a stroke (n=13).<sup>41</sup> All included studies  
212 evaluated the dominant shoulder only,<sup>8,11,12,40,42</sup> except de Oliveira et al.<sup>5</sup> and dos Santos et al.,<sup>41</sup> who  
213 evaluated both the healthy and pathological shoulders. It is unclear which shoulder (dominant or non-  
214 dominant) was evaluated by Lin et al.<sup>9</sup>

215

216 **Proprioception subcategories**

217 The most studied proprioception outcome measure included active (AJPS) (6 studies,  
218 n=131)<sup>5,9,11,12,40,42</sup> while one study evaluated passive joint position sense (PJPS) (1 study, n=13).<sup>41</sup>  
219 The proprioception error (PE) was understood to be the reproduction error in degrees between the  
220 target angle and the performed angle.<sup>5,9,11,12,40-42</sup> One study<sup>8</sup> (n=30) investigated the sense of  
221 movement (kinesthesia) through a time to detection of passive motion (TTDPM) protocol. The PE  
222 was recorded as the difference between the start and stop angles and was captured as the mean  
223 absolute average error in degrees. No studies evaluating the SoF or SoV were identified.

224

225 **Equipment**

226 Isokinetic dynamometer (Biodex Systems) (2 studies, n=43)<sup>8, 41</sup> and FASTRAK 3-Space magnetic  
227 tracking system (2 studies, n=39),<sup>9,12</sup> were the most used equipment to quantify the angle differential  
228 (PE) during active or passive movements. The bubble inclinometer (n=16),<sup>40</sup> the Apple iPod touch  
229 with an internal accelerometer and gyroscope (n=24),<sup>11</sup> wireless inertial measurement unit (IMU)  
230 system (n=22),<sup>5</sup> and a custom-built scale ruler with a pole mounted on a 4-wheeled arm support device  
231 (n=30) were used in a single study.<sup>42</sup>

232

233 **Direction of movement**

234 Both glenohumeral (GH) joint (n=144)<sup>5,8,9,11,40,41</sup> and scapular movements (n=30)<sup>42</sup> were evaluated.  
235 GH joint movements included flexion (3 studies, n=51),<sup>5,40,41</sup> extension (1 study, n=16),<sup>40</sup> internal  
236 rotation at 90° of abduction (ABD) (2 studies, n=46),<sup>8,40</sup> external rotation at 90° of ABD (2 studies,  
237 n=46),<sup>8,40</sup> ABD in the frontal plane (2 studies, n=35),<sup>5,41</sup> and scapular abduction (scapular plane  
238 elevation) (3 studies, n=63).<sup>9,11,12</sup> Scapular movements included scapular elevation (n=30),  
239 protraction (n=30), anterior/posterior tilting (n=30), and upward/downward rotation (n=30).<sup>42</sup>

241 **Taping protocols**

242 Three studies (n=38)<sup>8,41,42</sup> used placebo tape (non-elastic tape), including Cover-Roll™ (n=10),<sup>8</sup>  
243 Cramer tape™ (n=13),<sup>41</sup> and 3M Micropore tape™ (n=15).<sup>42</sup> A single study<sup>11</sup> (n=24) used elastic KT  
244 as their sham taping without applying any tension. In five studies (n=98),<sup>5,9,11,12,41</sup> participants acted  
245 as their own controls, having both the control and intervention conditions applied to the ipsilateral  
246 shoulder. Zanca et al.<sup>11</sup> (n=24) explored three conditions: (i) no tape, (ii) elastic KT with tension, and  
247 (iii) elastic KT without tension applied a week apart. Lastly, three studies (n=76)<sup>8,40,42</sup> compared an  
248 intervention group to a control group to test the effects of elastic KT on shoulder proprioception.

250 **Risk of bias**

251 The RoB of the included studies ranged from low to high, with a high level of agreement between  
252 raters for the scoring of RCTs (ICC=0.81 [0.74, 0.92]) and non-RCTs (ICC=0.94 [0.90, 0.98]). Of  
253 the included RCTs, one was deemed to have a high risk of bias,<sup>12</sup> three were assessed to have some  
254 concerns,<sup>11,40,42</sup> and a single study<sup>41</sup> supported a low risk of bias ([Table 1](#)). Regarding the non-RCTs  
255 ([Table 2](#)), two studies were found to have moderate risk of bias,<sup>8,9</sup> while the other was deemed to  
256 have low risk.<sup>5</sup>

258 **GRADE framework evidence profile and synthesis of results**

259 [Table 3](#) presents the analysed certainty of evidence by regrouping the studies according to the  
260 shoulder health conditions (healthy or pathological) and the proprioception outcomes (AJPS, PJPS,  
261 or kinesthesia). Currently, there is low certainty of evidence suggesting that elastic KT has mixed  
262 results on AJPS among healthy shoulders (4 studies, n=79).<sup>9,11,12,40</sup> Two studies<sup>11,12</sup> evaluating AJPS  
263 with elastic KT suggested no change to proprioception, whereas two studies<sup>9,40</sup> suggested a decrease  
264 in proprioception error with elastic KT; resulting in overall conflicting and low evidence with the  
265 AJPS outcomes.

266

267 In addition, there is very low certainty regarding the effects of elastic KT on AJPS in pathological  
268 shoulders (2 studies, n=52).<sup>5,42</sup> Very low certainty suggests that active scapular repositioning  
269 improved among pathological shoulders (1 study, n=30)<sup>42</sup> and no change was found with AJPS with  
270 GH joint movements (1 study, n=22).<sup>5</sup>

271

272 There is very low certainty for the improvement of PJPS among chronic hemiparetic (post-stroke)  
273 shoulders (1 study, n=13)<sup>41</sup> and also very low certainty that elastic KT has no effect on shoulder  
274 kinesthesia among individuals with SAPS (1 study, n=30)<sup>8</sup> (Table 4). No studies that examined PJPS  
275 or kinesthesia with elastic KT on healthy shoulders were identified for this review. As the certainty  
276 of evidence is very low or low concerning the effects of elastic KT in both healthy and pathological  
277 shoulders, regardless of the proprioception outcome evaluated, no concrete recommendations can be  
278 made at this time.

279 **DISCUSSION**

280 This systematic review is the first to our knowledge to evaluate the effectiveness of elastic KT on  
281 shoulder proprioception; more specifically, AJPS, PJPS, and kinesthesia among healthy and  
282 pathological shoulders. From our review, we present conflicting and inconsistent effectiveness of  
283 elastic KT on AJPS (low certainty) and PJPS with both healthy and pathological shoulders (very low  
284 certainty) as well as very low certainty of evidence to suggest that elastic KT influences kinesthesia  
285 among individuals with subacromial pain syndrome. Accordingly, we cannot encourage using elastic  
286 KT in clinical practice to improve shoulder proprioception. Our results echo those of past reviews  
287 involving the lower extremities and spine,<sup>26-28</sup> which report little to no effect of elastic KT on  
288 proprioception, except for a review addressing individuals with ankle instabilities that suggested  
289 improvements in balance, muscle strength, and proprioception.<sup>29</sup>

290 The interest in this topic arose from the common claim and belief that elastic KT can enhance  
291 proprioception; hence the clinical term “proprioceptive tape”.<sup>6,10</sup> Elastic KT is a popular therapeutic  
292 resource used by clinicians as the material is portable, economical, requires relatively little technical  
293 training, and it is suggested to be a supportive home therapy.<sup>13</sup> In addition, arguments exist for a  
294 positive placebo effect with the application of elastic KT<sup>43,44</sup> through the positive expectancy theory,<sup>45</sup>  
295 suggesting that placebo-prone personalities benefit from such outcomes in the presence of positive  
296 beliefs.<sup>47</sup> Despite substantial claims from the manufacturers and promoters<sup>13,46</sup> on the effectiveness  
297 of elastic KT tape as a therapeutic modality, there is little to no evidence to corroborate the immediate  
298 or mid-term effect of elastic KT on proprioception.

299  
300 Our very limited results can be partially explained by considering the hypothesized  
301 neurophysiological effects of elastic KT.<sup>13,46</sup> It has been argued that the main benefits of elastic KT  
302 are derived from the direct lifting of the skin,<sup>24</sup> which increases the space between the skin and  
303 subcutaneous tissues, promoting localized lymphatic drainage and increased blood flow.<sup>13</sup>  
304 Subsequently, pressure on pain receptors is relieved, reinforcing the body’s self-healing capacities.<sup>13</sup>  
305 It is further hypothesized that the “pump action” from the lymphatic and circulatory system stimulates  
306 the localized cutaneous mechanoreceptors,<sup>13</sup> generating tactile and sensorimotor changes,<sup>47</sup> including  
307 a heightened sensation of proprioception. This theory remains questionable until further examination  
308 of the specialized mechanoreceptors within the dermis and the soft tissue surrounding a joint.  
309 However, we acknowledge that evaluating the effects of elastic KT underneath the skin is illogical if  
310 elastic KT does not show any positive clinical responses.

311 To understand the results of our review, it is important to consider the current understanding of  
312 proprioception feedback, arising from both joint mechanoreceptors (providing information regarding  
313 internal mechanical forces, muscle length, joint velocity, stiffness, deep pressure,

314 acceleration/deceleration, tensile strain, joint motion, and joint position sense)<sup>48,49</sup> and cutaneous  
315 mechanoreceptors (providing information derived from external stimuli [discriminatory touch,  
316 pressure, skin movement – slip or flutter, skin stretching, vibration, and textures]).<sup>47</sup> We believe that  
317 if proprioceptive input came solely from cutaneous mechanoreceptors, our review could have found  
318 positive proprioceptive gains with PJPS and kinesthesia outcomes, as it can be theorized that a passive  
319 task does not primarily involve active mechanical tissue deformation surrounding a joint. This was  
320 not the case because the only study that investigated the effects of elastic KT on kinesthesia reported  
321 no change.<sup>8</sup> On the other hand, if our proprioception came solely from articular mechanoreceptors,  
322 we could anticipate no change in shoulder proprioception during AJPS tasks, as it can be argued that  
323 no direct stimulation to the deep joint mechanoreceptors occurs with the topical application of elastic  
324 KT. Our review presents inconsistent results, as some studies suggest positive effects,<sup>9,40-42</sup> while  
325 others have reported no effect<sup>5,8,12,41</sup> or a worsening<sup>11,12</sup> during an active joint matching task,  
326 regardless of shoulder health conditions (healthy or pathological), body segment, or joint taped. Our  
327 results raise questions whether cutaneous mechanoreceptors can be topically and superficially  
328 stimulated, as questioned by previous neurophysiological studies.<sup>50,51</sup> How sensory information is  
329 weighed and consolidated from cutaneous and articular mechanoreceptors within the nervous system  
330 is also of interest, as it would help researchers and clinicians further understand proprioceptive inputs  
331 as they pertain to injuries and athletic performance.

332

### 333 **Lack of standardisation**

334 A significant part of our inconsistent and conflicting results, resulting in very low to low certainty of  
335 evidence, can also be explained by the lack of standardisation between studies, including the various  
336 proprioception outcome measures, the populations, taping protocols, and what part of the shoulder  
337 complex is taped. Indeed, three distinct sub-modalities of proprioception were considered (AJPS,  
338 PJPS, kinesthesia), and each elastic KT protocol used was unique (see the Supplementary Material –  
339 Table S5). Taping protocols have differed regarding anatomical location, type of elastic KT, tension  
340 applied throughout the tape, and whether the effects were intended to facilitate or inhibit the  
341 underlying musculature. Only two studies<sup>5,41</sup> provided enough detail about their protocols, which  
342 allow comprehension of the purpose of the taping and encourage the replication of their studies, which  
343 would permit further testing of their results. More clearly defined taping protocols that can be  
344 accurately replicated by different researchers, in addition to psychometrically tested shoulder  
345 proprioception outcome measures, are needed to move forward.

346

347 **Strengths and limitations**

348 Strengths of this review include the meticulous search of the literature through four scientific  
349 databases, using three languages, and the application of validated risk of bias tools for critical  
350 appraisal and the development of an evidence profile using the GRADE framework. We also searched  
351 for studies evaluating all sub-modalities of shoulder proprioception, although only protocols  
352 evaluating JPS and kinesthesia were identified. Moreover, our results are systematically reported to  
353 encourage using the presented protocols and outcomes for future research on this topic.

354 Despite the methodological rigour, we recognise several limitations of this systematic review. First,  
355 weak reporting of psychometric properties, effect sizes, and small sample sizes limits the robustness  
356 of our conclusions. The certainty of evidence profile seems to have been impacted by the few  
357 identified studies and small samples for each proprioception outcome evaluated. Consequently, no  
358 concrete recommendations can be made at this time as the evidence remains conflicting and  
359 speculative (very low to low certainty of evidence).

360 Second, limited shoulder pathologies evaluated within the included studies may also hinder the  
361 broader clinical applicability of our findings. Future studies with a variety of shoulder pathologies  
362 are encouraged. Third, none of the included studies evaluated the effects of elastic KT beyond a single  
363 laboratory session, which hampers establishing the mid or long-term effects of elastic KT on shoulder  
364 proprioception. Therefore, our results can only be considered in the short-term. The aggregate of these  
365 factors limits the pooling of data for a meta-analysis and, ultimately, narrows the application of our  
366 findings for clinical practice. Standardized taping protocols and proprioception outcome measures  
367 are needed to address whether elastic KT influences shoulder proprioception in the short-, mid- or  
368 long-term.

369

370 **Clinical recommendations**

371 From our results, we have insufficient scientific evidence to recommend or discard the clinical  
372 application of elastic KT for the improvement of shoulder proprioception (very low to low certainty  
373 of evidence). Further studies investigating different shoulder elastic KT protocols and functional  
374 proprioception outcome measures are encouraged to establish the clinical effectiveness of elastic KT  
375 on known shoulder proprioception deficits across a wider variety of shoulder pathologies.<sup>52, 53</sup>

376

377

378 **CONCLUSIONS**

379

380 The application of elastic KT on healthy shoulders demonstrated mixed results with AJPS, where two  
381 studies indicated an improvement to proprioception and two indicated no change (low certainty of  
382 evidence). There is very low certainty that elastic KT improves AJPS among pathological shoulders  
383 (individuals with subacromial pain syndrome or rotator cuff tendinopathy) or PJPS (individuals with  
384 chronic hemiparetic shoulder). Furthermore, the use of elastic KT has no effect on kinesthesia  
385 (individuals with subacromial pain syndrome) (very low certainty). As the evidence suggests very  
386 low to low certainty regarding the effectiveness of elastic KT on the evaluated sub-modalities of  
387 shoulder proprioception, further research is necessary before elastic KT can be supported as an  
388 effective clinical rehabilitative approach.

389

390 **Conflict of Interest**

391 The authors declare no conflict of interest.

392

393 **REFERENCES**

- 394 1. Lewis JS. Rotator cuff tendinopathy/subacromial impingement syndrome: is it time for a new  
395 method of assessment? *Br J Sports Med.* 2009; 43: 259-64.
- 396 2. Greenberg DL. Evaluation and treatment of shoulder pain. *Med Clin North Am.* 2014; 98: 487-  
397 504.
- 398 3. Meislin RJ, Sperling JW and Stitik TP. Persistent shoulder pain: epidemiology,  
399 pathophysiology, and diagnosis. *Am J Orthop.* 2005; 34: 5-9.
- 400 4. Pribicevic M. Pain in Perspective. In: Ghosh S, (ed.). *Chapter 7: The epidemiology of shoulder*  
401 *pain: A narrative review of the literature.* Rijeka, Croatia 2012.
- 402 5. de Oliveira FCL, Pairot de Fontenay B, Bouyer LJ and Roy JS. Immediate effects of  
403 kinesiotaping on acromiohumeral distance and shoulder proprioception in individuals with  
404 symptomatic rotator cuff tendinopathy. *Clin Biomech.* 2019; 61: 16-21.
- 405 6. Morrissey D. Proprioceptive shoulder taping. *J Bodyw Mov Ther.* 2000; 4: 189-94.
- 406 7. Halseth T, McChesney JW, Debeliso M, Vaughn R and Lien J. The effects of kinesio taping  
407 on proprioception at the ankle. *J Sports Sci Med.* 2004; 3: 1-7.
- 408 8. Keenan KA, Akins JS, Varnell M, et al. Kinesiology taping does not alter shoulder strength,  
409 shoulder proprioception, or scapular kinematics in healthy, physically active subjects and subjects  
410 with Subacromial Impingement Syndrome. *Phys Ther Sport.* 2017; 24: 60-6.
- 411 9. Lin JJ, Hung CJ and Yang PL. The effects of scapular taping on electromyographic muscle  
412 activity and proprioception feedback in healthy shoulders. *J Orthop Res.* 2011; 29: 53-7.
- 413 10. Park YH and Lee JH. Effects of proprioceptive sense-based Kinesio taping on walking  
414 imbalance. *J Phys Ther Sci.* 2016; 28: 3060-2.
- 415 11. Zanca GG, Mattiello SM and Karduna AR. Kinesio taping of the deltoid does not reduce  
416 fatigue induced deficits in shoulder joint position sense. *Clin Biomech.* 2015; 30: 903-7.
- 417 12. Aarseth LM, Suprak DN, Chalmers GR, Lyon L and Dahlquist DT. Kinesio Tape and  
418 Shoulder-Joint Position Sense. *J Athl Train.* 2015; 50: 785-91.
- 419 13. Kumbrink B. *K-taping.* Springer, 2014.
- 420 14. Lim EC and Tay MG. Kinesio taping in musculoskeletal pain and disability that lasts for more  
421 than 4 weeks: is it time to peel off the tape and throw it out with the sweat? A systematic review with  
422 meta-analysis focused on pain and also methods of tape application. *Br J Sports Med.* 2015; 49: 1558-  
423 66.
- 424 15. Thelen MD, Dauber JA and Stoneman PD. The clinical efficacy of kinesio tape for shoulder  
425 pain: a randomized, double-blinded, clinical trial. *J Orthop Sports Phys Ther.* 2008; 38: 389-95.
- 426 16. Morris D, Jones D, Ryan H and Ryan CG. The clinical effects of Kinesio(R) Tex taping: A  
427 systematic review. *Physiother Theory Pract.* 2013; 29: 259-70.



- 428 17. Williams S, Whatman C, Hume PA and Sheerin K. Kinesio taping in treatment and prevention  
429 of sports injuries: a meta-analysis of the evidence for its effectiveness. *Sports medicine*. 2012; 42:  
430 153-64.
- 431 18. Roijezon U, Clark NC and Treleaven J. Proprioception in musculoskeletal rehabilitation. Part  
432 1: Basic science and principles of assessment and clinical interventions. *Man Ther*. 2015; 20: 368-77.
- 433 19. Ager AL, Roy JS, Roos M, Belley AF, Cools A and Hebert LJ. Shoulder proprioception: How  
434 is it measured and is it reliable? A systematic review. *J Hand Ther*. 2017; 30: 221-31.
- 435 20. Ager AL, Borms D, Deschepper L, et al. Proprioception: How is it affected by shoulder pain?  
436 A systematic review. *J Hand Ther*. 2020; 33: 507-16.
- 437 21. Myers JB, Wassinger CA and Lephart SM. Sensorimotor contribution to shoulder stability:  
438 effect of injury and rehabilitation. *Man Ther*. 2006; 11: 197-201.
- 439 22. Iandolo R, Squeri V, De Santis D, Giannoni P, Morasso P and Casadio M. Proprioceptive  
440 bimanual test in intrinsic and extrinsic coordinates. *Front Hum Neurosci*. 2015; 9: 72.
- 441 23. Han J, Waddington G, Adams R, Anson J and Liu Y. Assessing proprioception: A critical  
442 review of methods. *J Sport Health Sci*. 2016; 5: 80-90.
- 443 24. Kase K, Wallis J and Kase T. *Clinical therapeutic applications of the Kinesio taping method*.  
444 1st ed. Tokyo 2003.
- 445 25. Han JT, Lee JH and Yoon CH. The mechanical effect of kinesiology tape on rounded shoulder  
446 posture in seated male workers: a single-blinded randomized controlled pilot study. *Physiother*  
447 *Theory Pract*. 2015; 31: 120-5.
- 448 26. Mostafavifar M, Wertz J and Borchers J. A systematic review of the effectiveness of kinesio  
449 taping for musculoskeletal injury. *The Physician and sportsmedicine*. 2012; 40: 33-40.
- 450 27. Nelson NL. Kinesio taping for chronic low back pain: A systematic review. *J Bodyw Mov*  
451 *Ther*. 2016; 20: 672-81.
- 452 28. Logan CA, Bhashyam AR, Tisosky AJ, et al. Systematic Review of the Effect of Taping  
453 Techniques on Patellofemoral Pain Syndrome. *Sports Health*. 2017; 9: 456-61.
- 454 29. Munoz-Barrenechea IA, Garrido-Beroiza MA, Achiardi O, Seron P and Marzuca-Nassr GN.  
455 [A systematic review of the functional effectiveness of kinesiotaping in individuals with ankle  
456 instability]. *Medwave*. 2019; 19: e7635.
- 457 30. Desjardins-Charbonneau A, Roy JS, Dionne CE and Desmeules F. The Efficacy of Taping for  
458 Rotator Cuff Tendinopathy: A Systematic Review and Meta-Analysis. *Int J Sports Phys Ther*. 2015;  
459 10: 420-33.
- 460 31. Turgut E, Can EN, Demir C and Maenhout A. Evidence for taping in overhead athlete  
461 shoulders: a systematic review. *Res Sports Med*. 2021: 1-30.

- 462 32. Parreira Pdo C, Costa Lda C, Hespanhol LC, Jr., Lopes AD and Costa LO. Current evidence  
463 does not support the use of Kinesio Taping in clinical practice: a systematic review. *J Physiother.*  
464 2014; 60: 31-9.
- 465 33. Moher D, Liberati A, Tetzlaff J, Altman DG and Group P. Preferred reporting items for  
466 systematic reviews and meta-analyses: the PRISMA statement. *Int J Surg.* 2010; 8: 336-41.
- 467 34. Higgins JP, Altman DG, Gotzsche PC, et al. The Cochrane Collaboration's tool for assessing  
468 risk of bias in randomised trials. *BMJ.* 2011; 343: d5928.
- 469 35. Schünemann HJ, Higgins JP, Vist GE, et al. Completing 'Summary of findings' tables and  
470 grading the certainty of the evidence. *Cochrane Handbook for systematic reviews of interventions.*  
471 2019: 375-402.
- 472 36. de Oliveira FCL, Bouyer LJ, Ager AL and Roy JS. Electromyographic analysis of rotator cuff  
473 muscles in patients with rotator cuff tendinopathy: A systematic review. *J Electromyogr Kinesiol.*  
474 2017; 35: 100-14.
- 475 37. Granholm A, Alhazzani W and Moller MH. Use of the GRADE approach in systematic  
476 reviews and guidelines. *Br J Anaesth.* 2019; 123: 554-9.
- 477 38. Guyatt GH, Oxman AD, Vist GE, et al. GRADE: an emerging consensus on rating quality of  
478 evidence and strength of recommendations. *BMJ.* 2008; 336: 924-6.
- 479 39. Schünemann H, Brożek J, Guyatt G, , and Oxman A. Introduction to GRADE Handbook.  
480 Handbook for grading the quality of evidence and the strength of recommendations using the GRADE  
481 approach. Updated October 2013. Amsterdam: Cochrane Netherlands, 2013.
- 482 40. Burfeind SM and Chimera N. Randomized Control Trial Investigating the Effects of  
483 Kinesiology Tape on Shoulder Proprioception. *J Sport Rehabil.* 2015; 24: 405-12.
- 484 41. dos Santos GL, Souza MB, Desloovere K and Russo TL. Elastic Tape Improved Shoulder  
485 Joint Position Sense in Chronic Hemiparetic Subjects: A Randomized Sham-Controlled Crossover  
486 Study. *PLoS One.* 2017; 12: e0170368.
- 487 42. Shih YF, Lee YF and Chen WY. Effects of Kinesiology Taping on Scapular Reposition  
488 Accuracy, Kinematics, and Muscle Activity in Athletes With Shoulder Impingement Syndrome: A  
489 Randomized Controlled Study. *J Sport Rehabil.* 2018; 27: 560-9.
- 490 43. Lumbroso D, Ziv E, Vered E and Kalichman L. The effect of kinesio tape application on  
491 hamstring and gastrocnemius muscles in healthy young adults. *J Bodyw Mov Ther.* 2014; 18: 130-8.
- 492 44. Mak DN, Au IP, Chan M, et al. Placebo effect of facilitatory Kinesio tape on muscle activity  
493 and muscle strength. *Physiother Theory Pract.* 2019; 35: 157-62.
- 494 45. Price DD, Finniss DG and Benedetti F. A comprehensive review of the placebo effect: recent  
495 advances and current thought. *Annu Rev Psychol.* 2008; 59: 565-90.
- 496 46. Manufacturer KT. KT Tape: How does KT tape work? 2020.


- 497 47. Iheanacho F and Vellipuram AR. Physiology, Mechanoreceptors. *StatPearls*. Treasure Island  
498 (FL)2021.
- 499 48. Kaya D, Yertutanol FDK and Calik M. Neurophysiology and assessment of the  
500 proprioception. *Proprioception in orthopaedics, sports medicine and rehabilitation*. Springer, 2018,  
501 p. 3-11.
- 502 49. Purves D and Augustine G. *Mechanoreceptors Specialized for Proprioception*. Sunderland  
503 (MA): Sinauer Associates, 2001.
- 504 50. Haeberle H and Lumpkin EA. Merkel Cells in Somatosensation. *Chemosens Percept*. 2008;  
505 1: 110-8.
- 506 51. Nakatani M, Maksimovic S, Baba Y and Lumpkin EA. Mechanotransduction in epidermal  
507 Merkel cells. *Pflugers Arch*. 2015; 467: 101-8.
- 508 52. Ferreira GE, McLachlan AJ, Lin CC, et al. Efficacy and safety of antidepressants for the  
509 treatment of back pain and osteoarthritis: systematic review and meta-analysis. *BMJ*. 2021; 372:  
510 m4825.
- 511 53. Hultcrantz M, Rind D, Akl EA, et al. The GRADE Working Group clarifies the construct of  
512 certainty of evidence. *J Clin Epidemiol*. 2017; 87: 4-13.

513

514

515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545


Figure 1. Flowchart of the literature selection process performed according to the PRISMA statement.



546 **Table 1.** Risk of bias of randomized studies according to the Cochrane Risk of Bias Assessment (Version 2).<sup>34,35</sup>

Study ID	Experimental	Comparator	Outcome	D1	D2	D3	D4	D5	Overall	
Shih et al. 2018	Kinesiotaping	Placebo	AJPS	+	+	+	!	-	!	+
Burféind & Chimera 2015	Kinesiotaping	No tape	AJPS	+	+	+	!	-	!	!
dos Santos et al. 2017	Kinesiotaping	Sham taping	PJPS	+	+	+	+	-	+	+
Aaserth et al. 2015	Kinesiotaping	No tape	AJPS	+	+	+	-	-	-	-
Zanca et al. 2015	Kinesiotaping with tension	i) No taping ii) KT without tension	AJPS	+	+	+	!	-	!	!

547  
 548 **D1** = Randomisation process, **D2** = Deviations from the intended interventions, **D3** = Missing outcome data,  
 549 **D4** = Measurement of the outcome, **D5** = Selection of the reported result



**Table 2.** Methodological quality for non-randomized controlled trials studies of intervention assessed with ROBINS-I Tool (2016).

items	sub-items	Studies		
		Lin et al. 2011 <sup>9</sup>	Keenan et al. 2017 <sup>8</sup>	de Oliveira et al. 2019 <sup>5</sup>
1. Bias due to confounding	1.1	PY	PY	PY
	1.2	N	/	N
	1.3	/	/	/
	1.4	Y	PN	Y
	1.5	/	/	/
	1.6	N	PY	NI
	1.7	PY	PY	NI
	1.8	PY	PY	/
<b>risk of bias judgement (item 1)</b>		<b>low</b>	<b>moderate</b>	<b>low</b>
2. Bias in selection of participants into the study	2.1	N	N	N
	2.2	/	/	/
	2.3	/	/	/
	2.4	Y	PY	NI
	2.5	PY	/	/
<b>risk of bias judgement (item 2)</b>		<b>low/moderate</b>	<b>low/moderate</b>	<b>low</b>
3. Bias in classification of interventions	3.1	Y	Y	Y
	3.2	Y	Y	Y
	3.3	PN	Y	NI
<b>risk of bias judgement (item 3)</b>		<b>low</b>	<b>low</b>	<b>low</b>
4. Bias due to deviations from intended interventions	4.1	N	/	/
	4.2	N	/	/
	4.3	NI	NI	NI
	4.4	Y	Y	Y
	4.5	Y	Y	Y
	4.6	/	/	/
<b>risk of bias judgement (item 4)</b>		<b>low</b>	<b>low</b>	<b>low</b>
5. Bias due to missing data	5.1	Y	Y	Y
	5.2	N	N	N
	5.3	N	Y	N
	5.4	/	PY	/
	5.5	/	PN	/
<b>risk of bias judgement (item 5)</b>		<b>low</b>	<b>moderate</b>	<b>low</b>
6. Bias in measurement of outcomes	6.1	PY	PY	PN
	6.2	Y	Y	Y
	6.3	Y	Y	Y
	6.4	PN	PN	PN
<b>risk of bias judgement (item 6)</b>		<b>moderate</b>	<b>moderate</b>	<b>low/moderate</b>
7. Bias in the selection of the reported result	7.1	NI	NI	NI
	7.2	NI	NI	NI
	7.3	NI	NI	NI
<b>risk of bias judgement (item 7)</b>		<b>No information</b>	<b>no information</b>	<b>no information</b>
<b>Overall risk of bias</b>		<b>moderate risk</b>	<b>moderate risk</b>	<b>low risk</b>
<b>Methodological quality</b>		<b>moderate quality (MQ)</b>	<b>moderate quality (MQ)</b>	<b>good quality (GQ)</b>

A description of the risk of bias is available in the Supplementary Material.

ROBINS-I scores were converted to a specific quality classification based on the classification suggested by de Oliveira et al.<sup>36</sup>

- low risk = good quality (GQ)
- moderate risk = moderate quality (MQ)
- serious risk = low quality (LQ)
- critical risk = very low quality (VLQ)

**High quality does not exist in this categorization since the studies are non-randomized controlled trials.**

553 **Table 3.** Summary of certainty of evidence of the included studies assessed following the GRADE guidelines.

554

GRADE evidence profile							
Proprioception outcomes & Population	Studies	Limitations in study design (risk of bias)	Inconsistency	Indirectness (generalizability) (PICO)	Imprecision (sparse data; group size)	Publication bias	GRADE certainty of evidence
<b>Active Joint Position Sense (AJPS)</b>							
<b>Healthy shoulders (n=79)</b> Conflicting results for the effects of elastic KT on shoulder proprioception	<b>4 studies</b>	<b>Serious</b>	<b>Serious</b>	<b>Serious</b>	<b>Very Serious</b>	<b>Not serious</b>	<b>Low</b> ⊕⊕○○
	Aarseth et al. 2015	RCT (cross-over)					
	Burfeind & Chimera 2015	RCT					
	Lin et al. 2011	cross-sectional non-RCT study of intervention					
	Zanca et al. 2015	RCT (cross-over)					
<b>Pathological shoulders (n=52)</b> Subacromial pain syndrome Rotator cuff tendinopathy	<b>2 studies</b>	<b>Serious</b>	<b>Serious</b>	<b>Serious</b>	<b>Very serious</b>	<b>Serious</b>	<b>Very low</b> ⊕○○○



Scapular proprioception was improved, but no reported change for GHJ proprioception	Shih et al. 2018	RCT cross-sectional					
	de Oliveira et al. 2019	non-RCT study of intervention					
<b>Passive Joint Position Sense (PJPS)</b>							
No studies were identified as having evaluated PJPS amongst healthy shoulders.							
<b>Pathological shoulders (n=13)</b> Chronic hemiparetic (post-stroke) Improvement in PJPS	dos Santos et al.	<b>Not serious</b>  RCT cross-over	<b>Serious</b>	<b>Not serious</b>	<b>Very serious</b>	<b>Very serious</b>	<b>Very low</b> ⊕○○○
<b>Kinesthesia (sense of movement)</b>							
No studies were identified as having evaluated the sense of kinesthesia amongst healthy shoulders.							
<b>Pathological shoulders (n=30)</b> Subacromial pain syndrome No effect.	Keenan et al.	<b>Serious</b>  cross-sectional Non-RCT study of intervention	<b>Not serious</b>	<b>Not serious</b>	<b>Very serious</b>	<b>Extremely serious</b>	<b>Very low</b> ⊕○○○

555

556 The certainty of evidence was assessed using the grading of recommendations assessment, development and evaluation (GRADE) framework.

557 (**Not serious**) = Quality not downgraded, (**Serious**) = Factor downgraded by one level, (**Very serious**) = Factor downgraded by two levels, (**Extremely serious**) = For non-  
558 randomized studies assessed with ROBINS-I, rating down by three levels.

559

560 **Abbreviations:** AJPS, active joint position sense; CRoB-2, Cochrane Risk of Bias Assessment (Version 2); CI, confidence interval; GHJ, glenohumeral joint, KT, kinesiology tape;  
 561 PJPS, passive joint position sense; RoB, risk of bias.

562 The body of evidence for an outcome may be determined to have serious or very serious issues for the affected domain (or critically serious for risk of bias when ROBINS-I is used).

563 See the Supplementary Material for the processed followed for the development and presentation of the GRADE evidence profile.

564

565

**Table 4.** Summary of findings of the effectiveness of elastic kinesiology taping on proprioception.

Study	Main Findings	Proprioception Outcome	Study Design	Risk of Bias	GRADE certainty of evidence
<b>Healthy shoulders</b>					
<b>Aarseth et al. (2015)</b> (n=27)	No change at 50° or 110° of scapular abduction. ↑ proprioceptive error at 90° of scapular abduction (2.65°, $p=0.01$ ).	AJPS	RCT (cross-over)	High risk	<b>Low</b> ⊕⊕○○
<b>Burfeind &amp; Chimera (2015)</b> (n=16)	↓ proprioception error in flexion ( $p=0.04$ ) and ER ( $p=0.03$ ). Control group (no tape): ↑ variability with their proprioception performance.	AJPS	RCT	Some concerns	
<b>Lin et al. (2011)</b> (n=12)	↓ proprioception error ( $11.9°±8.3°$ , $p<0.005$ ).	AJPS	RCT (cross-sectional)	Moderate risk	
<b>Zanca et al. (2015)</b> (n=24)	No effects on proprioception following a muscle fatigue protocol at any angle (50°, 70°, 90° scapular abduction) ( $p=0.41$ ).	AJPS	Randomized crossover single-blind study	Some concerns	
<b>Pathological shoulders</b>					
<b>Shih et al. (2018)</b> (n=30)	<i>Subacromial pain syndrome (Overhead athletes)</i> ↓ proprioceptive error of the scapular for up/down rotation ( $p=0.04$ ) and anterior/posterior tilting ( $p=0.03$ )	AJPS	RCT	Some concerns	<b>Very low</b> ⊕○○○

<i>Rotator cuff tendinopathy</i>					
<b>de Oliveira et al. (2019)</b> (n=22)	No reported change to proprioception with elastic KT at low or mid-amplitudes (45°-65°, 80°-100°) ( $p>0.05$ ).	AJPS	Cross-sectional	Low risk	
<i>Chronic hemiparetic (post-stroke)</i>					
<b>Dos Santos et al. (2017)</b> (n=13)	↓ PJPS error in abduction at 30° and 60° as well as flexion at 30° and 60° (all $p<0.010$ ). Proprioception improved regardless of the level of sensorimotor impairment.	PJPS	Randomized sham-controlled crossover study	Low risk	<b>Very low</b> ⊕○○○
<i>Subacromial pain syndrome</i>					
<b>Keenan et al. (2017)</b> (n=30)	Elastic KT did not have an effect on kinesthesia ( $0.033 \leq p \leq 0.77$ ).	Kinesthesia	Placebo controlled quasi-experiment	Moderate risk	<b>Very low</b> ⊕○○○

Abbreviations: AJPS, active joint position sense; ↑, increase; ↓, decrease; KT, kinesiology tape; PJPS, passive joint position sense. Methodological quality assessed with Risk of bias of randomized studies according to the Cochrane Risk of Bias Assessment (Version 2) and ROBINS-I (non-RCTs of intervention).

Level of evidence assessed with GRADE framework.

566 The GRADE certainty of evidence can be evaluated as very low, low, moderate or high certainty (See the Supplementary Material).

567

568 As the evidence has been evaluated to be of very low to low quality, and only a small number of studies have been identified which evaluated a shoulder proprioception  
 569 outcome, a strength of recommendation could not be determined. The aggregate of evidence is currently so low that any recommendation on the effectiveness of elastic  
 570 KT on shoulder proprioception outcomes remains speculative.


571

572

573

574

575



Click here to access/download

**Supporting File**

BJPT-D-22-00095 Supplementary material\_Final.docx

