

# Oral microbiome and nitric oxide biomarkers in older people with mild cognitive impairment and APOE4 genotype

Joanna E. L'Heureux<sup>a</sup>, Anne Corbett<sup>a</sup>, Clive Ballard<sup>a</sup>, David Vauzour<sup>b</sup>, Byron Creese<sup>c</sup>, Paul G. Winyard<sup>a</sup>, Andrew M. Jones<sup>a</sup> and Anni Vanhatalo<sup>a,\*</sup>

<sup>a</sup>Faculty of Health and Life Sciences, University of Exeter Medical School, University of Exeter, St Luke's campus, Exeter EX1 2LU, United Kingdom

<sup>b</sup>Norwich Medical School, University of East Anglia, Norwich NR4 7TJ, United Kingdom

<sup>c</sup>Department of Life Sciences, University of Brunel, London UB8 3PH, United Kingdom

\*To whom correspondence should be addressed: Email: [a.vanhatalo@exeter.ac.uk](mailto:a.vanhatalo@exeter.ac.uk)

Edited By Hülya Bayir

## Abstract

Apolipoprotein E4 (APOE4) genotype and nitric oxide (NO) deficiency are risk factors for age-associated cognitive decline. The oral microbiome plays a critical role in maintaining NO bioavailability during aging. The aim of this study was to assess interactions between the oral microbiome, NO biomarkers, and cognitive function in 60 participants with mild cognitive impairment (MCI) and 60 healthy controls using weighted gene co-occurrence network analysis and to compare the oral microbiomes between APOE4 carriers and noncarriers in a subgroup of 35 MCI participants. Within the MCI group, a high relative abundance of *Neisseria* was associated with better indices of cognition relating to executive function (Switching Stroop,  $r_s = 0.33$ ,  $P = 0.03$ ) and visual attention (Trail Making,  $r_s = -0.30$ ,  $P = 0.05$ ), and in the healthy group, *Neisseria* correlated with working memory (Digit Span,  $r_s = 0.26$ ,  $P = 0.04$ ). High abundances of *Haemophilus* ( $r_s = 0.38$ ,  $P = 0.01$ ) and *Haemophilus parainfluenzae* ( $r_s = 0.32$ ,  $P = 0.03$ ), that co-occurred with *Neisseria* correlated with better scores on executive function (Switching Stroop) in the MCI group. There were no differences in oral nitrate ( $P = 0.48$ ) or nitrite concentrations ( $P = 0.84$ ) between the MCI and healthy groups. Linear discriminant analysis Effect Size identified *Porphyromonas* as a predictor for MCI and *Prevotella intermedia* as a predictor of APOE4-carrier status. The principal findings of this study were that a greater prevalence of oral *P. intermedia* is linked to elevated genetic risk for dementia (APOE4 genotype) in individuals with MCI prior to dementia diagnosis and that interventions that promote the oral *Neisseria*–*Haemophilus* and suppress *Prevotella*-dominated modules have potential for delaying cognitive decline.

**Keywords:** cognitive status, aging, nitrate, *Prevotella*, genetic risk

## Significance Statement

Cognition typically declines during aging and mild cognitive impairment (MCI) may progress to the development of Alzheimer's disease (AD). Periodontal disease-causing bacteria have been linked to worsened cognitive function during aging and the development of AD, which may involve dysfunction of the nitrate–nitrite–nitric oxide pathway. We found that the oral *Porphyromonas* genus was associated with MCI and that the abundance of *Prevotella intermedia* was a predictor of apolipoprotein E4-carrier status. The balance between two metabolic pathways for oral nitrate reduction, denitrification, and dissimilatory nitrate reduction to ammonia (DNRA) was skewed toward DNRA in MCI. These findings have significant implications for understanding preclinical cognitive risk states and how cognitive decline could be delayed or prevented using prebiotic interventions.

OXFORD  
UNIVERSITY PRESS

**Competing Interest:** J.E.L., D.V., B.C., P.G.W., A.M.J., A.V. declare no competing interests. C.B. has received consulting fees from Acadia pharmaceutical company, AARP, Addex pharmaceutical company, Eli Lilly, Enterin pharmaceutical company, GWPharm, H. Lundbeck pharmaceutical company, Novartis pharmaceutical company, Janssen Pharmaceuticals, Johnson and Johnson pharmaceuticals, Novo Nordisk pharmaceutical company, Orion Corp pharmaceutical company, Otsuka America Pharm. Inc, Sunovion Pharm. Inc, Suven pharmaceutical company, Roche pharmaceutical company, Biogen pharmaceutical company, Synexus clinical research organization and tauX pharmaceutical company and research funding from Synexus clinical research organization, Roche pharmaceutical company, Novo Nordisk pharmaceutical company and Novartis pharmaceutical company. A.C. discloses financial relationships with Suven and Janssen pharmaceutical companies for consultancy work.

**Received:** June 6, 2024. **Accepted:** November 18, 2024

© The Author(s) 2025. Published by Oxford University Press on behalf of National Academy of Sciences. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

## Introduction

Cognitive decline is typically associated with aging (1). In some individuals, clinically significant decline leads to the development of mild cognitive impairment (MCI), which affects about 15% of older adults. MCI is the greatest risk factor for the development of dementia or Alzheimer's disease (AD), with an estimated 10% of people with MCI converting to dementia each year (2). Cognitive decline and dementia represent a major public health issue, and there is an urgent need to elucidate the risk factors for decline and explore means to reduce this risk (3).

Periodontitis has been associated with worsened cognitive function (4) and missing teeth in the oral cavity have been associated with lower Mini Mental State Examination test scores (5). A potential mechanism linking oral health and cognitive decline is the increased inflammation and damage caused by pathogenic oral bacteria, such as *Porphyromonas gingivalis*, *Treponema denticola*, and *Prevotella intermedia*, which in turn increase the risk for periodontal diseases (6). Patients with AD present with higher levels of *P. gingivalis* and lower bacterial diversity in the oral cavity compared with healthy controls (7, 8). Two routes have been suggested by which oral pathogens may cause cognitive decline. The direct route is via trauma in the mouth, whereby the oral bacteria gain access to the circulatory system and then relocate to the brain by traversing the blood–brain barrier (9), which is increasingly permeable in AD. Known oral disease-causing bacteria have been found in the cerebrospinal fluid of patients with a brain abscess (10), and *P. gingivalis* has been discovered in the brains of patients with AD (11). The oral bacteria may also indirectly affect the brain by impairing the oral mucosal barrier and allowing metabolites produced by the oral bacteria to enter the circulatory system or exacerbating inflammation through the overproduction of cytokines (12, 13).

One potential mechanism linking oral health and cognition is the production of nitric oxide (NO) via the nitrate–nitrite–NO pathway (14). NO is an important signaling molecule for many physiological processes, such as vasodilation, muscle contraction, neurotransmission, and host defense against microorganisms (15–17). There are two known methods of NO production in mammals. The L-arginine pathway endogenously produces NO via NO synthase (NOS) enzymes (18). The second mechanism is through the nitrate–nitrite–NO pathway, where commensal oral bacteria reduce nitrate to nitrite, which is further reduced to form NO in the circulation and in tissues (19). A main role of NO in the brain is binding to guanylyl cyclase which acts as a pre- or postsynaptic messenger. In preclinical models, NOS activity has been detected in the hippocampus, which is the brain region responsible for learning and memory, and inhibition of NOS resulted in impaired memory (20). NO has a role in synaptic plasticity and long-term potentiation, and a reduction in NO availability may be linked with the inability of patients with AD to retain new information (21, 22). During aging, endogenous NO production is reduced through decreased NOS gene expression and increased degradation of arginine, catalyzed by arginase, which is associated with cardiovascular diseases such as hypertension and risk of vascular AD (23–25). Decreased NO biomarker concentrations of nitrate and nitrite in the plasma and brains of patients with AD have been reported compared with healthy controls (26–28).

A key genetic risk factor for cognitive decline and AD is the apolipoprotein E4 (APOE4) allele, which has been associated with weakening of the blood–brain barrier (29). APOE4 carriers also have an elevated risk for other systemic diseases,

such as hypertension, atherosclerosis, and skeletal muscle weakness (30–33), and these conditions are also notably hallmarked by NO deficiency. This suggests that there may be interactions between APOE4, NO bioavailability and the oral microbiome during aging-associated cognitive decline. It is, however, unknown whether characteristics of the oral microbiome correlate with cognitive function, or whether changes in oral microbiome composition may already be detected in healthy older individuals with MCI but prior to dementia diagnosis.

The purpose of this study was, therefore, to compare the oral microbiomes and oral NO biomarkers between individuals with MCI and healthy controls, and explore relationships between co-occurring modules of oral bacteria, cognitive function, and NO biomarkers. A secondary aim was to perform an MCI subgroup analysis to compare oral microbiomes and NO biomarkers between APOE4 carriers and noncarriers.

## Results

### Participant characteristics

There were 120 participants recruited in the study. After data pre-processing, 5 samples were removed from further analysis due to low-quality data. Therefore, 115 samples were analyzed in total. Included in the data analysis were 60 participants in the healthy group (17 males; mean age  $\pm$  SD,  $67 \pm 5$  years; 43 females; mean age  $\pm$  SD,  $67 \pm 8$  years) and 55 participants in the MCI group (8 males; mean age  $\pm$  SD,  $70 \pm 6$  years; 47 females; mean age  $\pm$  SD,  $68 \pm 8$  years). APOE status was defined as “high-risk” APOE4 carriers (E3E4, E4E4) and “low-risk” APOE4 noncarriers (E2E3, E3E3) in a subset of MCI participants. E2E4 carriers ( $n = 2$ ) were excluded from this analysis due to ambiguous risk status, and therefore, 33 participants were included in the APOE4 subgroup analysis (Table 1).

### Mouth rinse nitrate and nitrite concentrations


There were no differences in mouth rinse nitrate (median  $\pm$  interquartile range [IQR]; healthy  $122 \pm 403$   $\mu$ M; MCI  $80 \pm 263$   $\mu$ M,  $P = 0.48$ ) or nitrite concentrations (median  $\pm$  IQR; healthy  $35 \pm 48$   $\mu$ M; MCI  $34 \pm 53$   $\mu$ M,  $P = 0.84$ ) between the healthy and MCI groups (Fig. 1A and B). Within the MCI group, there were no differences between APOE4 carriers ( $118 \pm 110$   $\mu$ M) and noncarriers ( $206 \pm 258$   $\mu$ M) in mouth rinse nitrate ( $P = 0.30$ ) or nitrite concentrations (carriers  $42 \pm 46$   $\mu$ M, noncarriers  $74 \pm 48$   $\mu$ M,  $P = 0.12$ ).

### Alpha and beta diversity of the oral microbiome

Species diversity was assessed using the Shannon H' index, and species richness was measured using Chao1. The Mann–Whitney U test revealed that there were no differences in

**Table 1.** Cohort APOE4 status characteristics.

Characteristic	APOE4 carrier (n = 14)	APOE4 noncarrier (n = 19)
Risk group	High	Low
APOE status (n)	E3E4 (n = 13) E4E4 (n = 1)	E2E3 (n = 4) E3E3 (n = 15)
Male sex (n)	3	3
Female sex (n)	11	16
Male age (mean years $\pm$ SD)	$68 \pm 6$	$70 \pm 7$
Female age (mean years $\pm$ SD)	$68 \pm 7$	$69 \pm 8$




**Fig. 1.** Mouth rinse nitrate and nitrite concentrations and species diversity. The median mouth rinse nitrate (A) and nitrite (B) concentrations were not different between healthy and MCI groups. There were also no differences in median Shannon H' index (C) or Chao1 species richness (D) between the healthy and MCI groups. Error bars in (A) and (B) indicate IQR. The box plots in (C) and (D) represent the median and IQR, while whiskers show the minimum and maximum values.

Shannon H' index ( $P=0.75$ , Fig. 1C) or Chao1 species richness ( $P=0.45$ , Fig. 1D) between the healthy and MCI groups. There were also no differences in Shannon H' diversity ( $P=0.32$ ) or in Chao1 species richness ( $P=0.92$ ) between APOE4 carriers and noncarriers.

Nonmetric multidimensional scaling (NMDS) was used to explore possible underlying patterns in the microbiome data. Stress was 0.01, meaning that the individual distances between the objects were well represented. The NMDS showed that the oral microbiomes of healthy and MCI participants displayed no distinct groupings and that there were no significant differences between the healthy and MCI groups ( $P=0.80$ , Fig. 2A) or between APOE4 carriers and noncarriers ( $P=0.70$ , Fig. 2B).


### Weighted gene co-occurrence network analysis of the oral microbiome and cognitive test outcomes

Due to the similarities in bacterial diversity between the healthy and MCI groups, a consensus correlation network was used to assess consistencies among the results of healthy and MCI groups. The weighted gene co-occurrence network analysis (WGCNA) analysis revealed seven consensus modules of oral bacteria, which were given arbitrary colors and labeled from ME0 to ME7 (Fig. 3). ME0 contained the operational taxonomic units (OTUs) which did not significantly correlate with other OTUs and were thus not assigned to a module. Table 2 provides a list of the species and genera assigned to modules ME1–ME7.



**Fig. 2.** NMDS of healthy and MCI groups. (A) NMDS showed no distinct groupings or significant differences in the oral microbiomes between healthy (blue) and MCI groups (orange;  $P=0.80$ ). (B) NMDS also indicated no distinct groupings or significant differences between the APOE4 carriers (red) and noncarriers (blue;  $P=0.30$ ) within the MCI group. NMDS was based on Bray–Curtis dissimilarity, and the data points represent the oral microbiome of each individual participant. Only selected species of highly abundant species are shown in the plot for visual clarity.

The cognitive test outcomes included in the network analysis included measures of working memory (Digit Span, Paired Associates Learning, Self-ordered Search), executive function (Verbal Reasoning, Switching Stroop), and visual attention (Trail Making). High scores in tasks on working memory and executive function indicate intact cognitive function, whereas high scores in the Trail Making test (which includes a time dimension) indicate impaired cognitive function in the visual attention domain. The network correlations were notably distinct between the healthy and MCI groups for microbiome modules ME2 (dominated by *Veillonella* and *Megasphaera*), ME3 (dominated by *Prevotella*), and ME6 (dominated by *Neisseria* and




**Fig. 3.** Consensus gene dendrogram, generated using a signed network where modules represent positively correlated taxonomic units. Seven distinct microbiome modules (ME1–ME7) were identified.

**Table 2.** OTUs assigned to consensus WGCNA modules.

Module	Module color	OTU
ME1	Black	<i>lg_Lachnoanaerobaculum</i> <i>lg_Mogibacterium</i> <i>lg_Lachnoanaerobaculum umeaense</i>
ME2	Blue	<i>lg_Mogibacterium diversum</i> <i>lg_Lancefieldella</i> <i>lg_Megasphaera</i> <i>lg_Veillonella</i> <i>lg_Actinomyces pacaensis</i> <i>lg_Lancefieldella parvula</i> <i>lg_Veillonella atypica</i> <i>lg_Veillonella parvula</i>
ME3	Red	<i>lg_Prevotella</i> <i>lg_P. intermedia</i> <i>lg_P. jejuni</i> <i>lg_P. melaninogenica</i>
ME4	Turquoise	<i>lg_Fusobacterium</i> <i>lg_Leptotrichia</i> <i>lg_Porphyrmonas</i> <i>lg_Tannerella</i> <i>lg_Fusobacterium nucleatum</i> <i>lg_Porphyrmonas gingivalis</i> <i>lg_Tannerella sp. oral taxon HOT-286</i>
ME5	Brown	<i>lg_Dialister</i> <i>lg_Parvimonas</i> <i>lg_Treponema</i> <i>lg_Parvimonas micra</i>
ME6	Yellow	<i>lg_Haemophilus</i> <i>lg_Neisseria</i> <i>lg_Haemophilus parainfluenzae</i>
ME7	Green	<i>lg_Capnocytophaga</i> <i>lg_Campylobacter showae</i> <i>lg_Capnocytophaga gingivalis</i> <i>lg_Capnocytophaga sputigena</i>


Only the assigned species and genus are shown for brevity.

*Haemophilus*; Fig. 4A and B). A consensus network indicated that there were no significantly conserved correlations when the groups were combined (Fig. 4C). In the healthy group, ME2 correlated positively with oral nitrite and ME7 (dominated by *Capnocytophaga*) with oral nitrate concentration, while there were no significant correlations between any of the microbiome modules and the cognitive function tests (Fig. 4A). In the MCI group, the ME2 module did not correlate with NO biomarkers,



**Fig. 4.** Signed WGCNA heat maps for the healthy (A) and MCI (B) groups, together with a consensus network (C) combining the two groups. Positive correlations between modules and traits are shown in green, whereas negative correlations are shown in red. The first cell value denotes the  $r$ -value and the second value (in parentheses) denotes the  $P$ -value. The cognitive tests included Digit Span (SPAN), Paired Associates Learning (PAL), Verbal Reasoning (VERB), Self-ordered Search (SEARCH), Switching Stroop (STROOP), and Trail Making (TRAILS).

but this module correlated negatively with working memory (Paired Associates Learning task;  $r_s = -0.36$ ,  $P = 0.007$ ) and executive function (Switching Stroop task;  $r_s = -0.32$ ,  $P = 0.02$ ), and positively with a Trail Making task of visual attention, where higher scores are indicative of cognitive decline ( $r_s = 0.32$ ,  $P = 0.02$ ; Fig. 4B). In the MCI group, the ME6 module (dominated by *Neisseria* and *Haemophilus*) correlated positively with oral nitrite concentration ( $r_s = 0.31$ ,  $P = 0.02$ ), cognitive test outcomes on working memory (Self-Ordered Search task;  $r_s = 0.27$ ,




**Fig. 5.** Correlation networks in the healthy (A) and MCI (B) groups show the significantly correlated clinical traits (gray), together with *Haemophilus* and *Neisseria* from ME6 (blue). The line color denotes the direction of the correlation where green is positive, and red is negative. The color brightness indicates the strength of the correlation. The cognitive variables included total scores (TOT) in Digit Span (SPAN), Paired Associates Learning (PAL), Verbal Reasoning (VERB), Self-ordered Search (SEARCH), Switching Stroop (STROOP), and Trail Making (TRAILS) tasks.

$P=0.05$ ), and executive function (Switching Stroop task;  $r_s=0.37$ ,  $P=0.005$ ; Fig. 4B), and correlated negatively with the Trail Making score (high Trail Making score indicating impaired visual attention;  $r_s=-0.27$ ,  $P=0.05$ ; Fig. 4B).


The ME6 module, with positive interactions with cognition in the MCI group, was explored further by creating genus- and species-level Spearman correlation networks with cognitive outcomes and NO biomarkers (Fig. 5). In the healthy group, *Neisseria* correlated with Digit Span summary score which assesses working memory ( $r_s=0.26$ ,  $P=0.04$ ; Fig. 5A). In the MCI group, *Neisseria* ( $r_s=0.33$ ,  $P=0.03$ ), *Haemophilus* ( $r_s=0.38$ ,  $P=0.01$ ), and *Haemophilus parainfluenzae* ( $r_s=0.32$ ,  $P=0.03$ ) correlated with Switching Stroop summary score which assesses executive function (Fig. 5B), and *Haemophilus* ( $r_s=0.35$ ,  $P=0.03$ ) and *H. parainfluenzae* ( $r_s=0.34$ ,  $P=0.04$ ) correlated with nitrite concentration (Fig. 5B). In the MCI group, *Neisseria* also negatively correlated with the Trail Making task score, where high scores are indicative of impaired visual attention ( $r_s=-0.30$ ,  $P=0.05$ ).

### Linear discriminant analysis effect size comparisons between APOE4 carriers and noncarriers

Linear discriminant analysis effect size (LEfSe) analysis identified *Porphyromonas* as a potential biomarker in the MCI group



**Fig. 6.** LEfSe showed five significantly abundant taxonomic units differed by APOE4 status. APOE4 carriers were defined as E3E4/E4E4 and noncarriers as E2E3/E3E3 allele carriers. Negative bars indicate that the taxonomic unit was significantly abundant in the APOE4 carriers, and positive bars indicate greater abundance in the noncarriers.



**Fig. 7.** Correlation networks in APOE4 carriers (A) and noncarriers (B) of the MCI group showing the significantly correlated clinical traits (gray) and OTUs (blue) from the microbiome module ME3 where *Prevotella* was a dominant genus. The line color denotes the direction of the correlation where green is positive, and red is negative. The color brightness indicates the strength of the correlation. The cognitive variables include total scores (TOT) in Paired Associates Learning (PAL) and Self-ordered Search (SEARCH).

compared with healthy controls (log-linear discriminant analysis [log LDA] score, 2.34;  $P=0.03$ ). APOE4 risk status was available for a subset of the participants (Table 1), where LEfSe showed that in APOE4 carriers, *P. intermedia* was more abundant (log LDA score, 2.96,  $P=0.04$ ; Fig. 6), and *Leptotrichia hongkongensis*, *Selenomonas* sp. oral taxon 920, *Actinomyces* sp. HMT 175, and *Prevotella oris* were less abundant compared with the APOE4 noncarriers (Fig. 6).

To further explore the relationships between *Prevotella*, NO biomarkers, cognitive performance, and APOE status, we generated a Spearman correlation network using the WGCNA ME3 module (Fig. 7). In the APOE4 carrier group, *Prevotella melaninogenica* correlated positively with working memory assessed by Self-ordered Search task ( $r_s=0.58$ ,  $P=0.03$ ), while

*Prevotella jejuni* correlated negatively with the Paired Associates Learning task which also assesses working memory ( $r_s = -0.56$ ,  $P = 0.04$ , Fig. 7A). In the APOE4 noncarrier group, mouth rinse nitrate ( $r_s = -0.73$ ,  $P = 0.007$ ) and nitrite concentrations ( $r_s = -0.64$ ,  $P = 0.04$ ) correlated negatively with *P. intermedia* (Fig. 7B).

## Discussion

This study characterized oral microbiome modules that are associated with cognitive status, NO homeostasis, and genetic risk for dementia in the Platform for Research Online to Investigate Genetics and Cognition in Ageing (PROTECT) cohort of people over 50 years of age who have not received a dementia diagnosis. The principal original findings were that in people with MCI, a high relative abundance of *Neisseria-Haemophilus* module of co-occurring oral bacteria (ME6) was associated with better cognitive outcomes on working memory, executive function, and visual attention, as well as a greater oral nitrite concentration, while a high abundance of the oral *Prevotella*-dominated module (ME3) was associated with low oral nitrate availability. *Prevotella intermedia* was identified as a predictor for elevated genetic risk for dementia (APOE4 carrier status). In contrast, in healthy controls, high concentrations of NO biomarkers were associated with high abundances of *Veillonella-Megasphaera* (ME2) and *Capnocytophaga* (ME7)-dominated microbiome modules, and in healthy controls, no microbiome module correlated with cognitive outcomes. We showed that the genus *Porphyromonas* was more abundant in MCI than in healthy controls, supporting the notion that the presence of elevated oral *Porphyromonas* precedes dementia diagnosis. These results provide a rationale for further research into underlying mechanisms that connect the oral microbiome to cognitive health through the trajectory from health to MCI, and ultimately to postdementia diagnosis. They also highlight the potential for interventions to ameliorate or delay aging-associated cognitive decline through promotion of the *Neisseria-Haemophilus* and eradication of the *Prevotella*-dominated modules of co-occurring oral bacteria.

Within the MCI group, the oral *Neisseria-Haemophilus* module (ME6) was associated with a greater oral nitrite concentration, and a high abundance of the *Prevotella*-dominated module (ME3) was associated with a low oral nitrate concentration. Several *Neisseria* species contain genes that encode proteins for nitrate reduction (*narG*, *napA*) and for the denitrification pathway (*nirK*, *norB*, or *nosZ*; search performed on <https://www.uniprot.org/>, accessed on 2023 November 20), which restores NO precursors to the circulation thus helping maintain systemic NO availability (34). In contrast, many *Prevotella* species, including *P. intermedia*, *P. jejuni*, and *P. melaninogenica* of the ME3 module in the present study, contain the *nrfA* gene of the dissimilatory pathway of nitrate reduction to ammonia (DNRA; search performed on <https://www.uniprot.org/>, accessed on 2023 November 20), which “short-circuits” the nitrate–nitrite–NO cycle by removing NO precursors from circulation and decreasing systemic NO availability (34). In healthy controls, the oral nitrite concentration correlated positively with the *Veillonella-Megasphaera* (ME2) module, and oral nitrate with a *Capnocytophaga*-dominated microbiome module (ME7). The latter is consistent with the increase in relative abundance of the *Capnocytophaga* genus following 10 days of high nitrate diet (35), while *Veillonella* species are among the most abundant and potent nitrate-reducing bacteria in the oral cavity (36). In addition to affecting systemic NO homeostasis via the enterosalivary circulation, oral nitrite production represents an

important part of host defense: NO and other reactive nitrogen intermediates formed from acidified nitrite inhibit the growth of a wide range of microorganisms (37), and nitrite is bactericidal against oral pathogens including *Fusobacterium nucleatum*, *Eikenella corrodens*, and *P. gingivalis* (38). Diet was not monitored in the present study, and future research is needed to explore how variation in habitual nitrate and macronutrient intakes may influence the relationships between the oral microbiome modules and NO bioavailability. A high oral nitrate:nitrite ratio, characteristic of a nitrate-rich diet (such as the Mediterranean and “Dietary Approaches to Stop Hypertension” diets), favors bacteria of the denitrification pathway (34) which we have shown to be associated with good cognitive outcomes ((35); present study).

A novel finding of the present study was that in the MCI group, the *Neisseria-Haemophilus* module was linked to a broader range of cognitive domains than was seen in the healthy controls. Although module-level correlations with cognitive outcomes did not reach significance in healthy controls, the OTU-level correlation network showed that *Neisseria* genus correlated with working memory. A previous study in healthy older people found a consensus WGCNA correlation between the *Neisseria-Haemophilus* module and a “sustained attention” cognitive domain, which was robust across dietary nitrate and placebo interventions (35). The *Neisseria-Haemophilus* module has been linked to characteristics associated with cognitive health, such as periodontal health, younger age, lower BMI, and non-smoking status (39). *Neisseria* and *Haemophilus* consistently co-occur in the oral ecosystems (35, 40, 41) and are found in high abundances in saliva (42, 43). Horizontal gene transfer has been shown to occur between *Haemophilus* and *Neisseria meningitidis* (44), suggesting that these genera have a mutually beneficial relationship. Collectively, these results indicate that a high relative abundance of bacteria belonging in the *Neisseria-Haemophilus* module is associated with better cognitive outcomes in individuals with MCI (present study) as well as in healthy older people (35).

There is a significant inflammatory component to the etiology of neurological damage that ultimately manifests as AD (45). *Prevotella* species play a role as pathobionts or pathogens in inducing periodontitis, a key risk factor for cognitive impairment and AD (46), and baseline antibodies for *P. intermedia* and *F. nucleatum* have been implicated as predictors of MCI and AD during a 10-year follow-up (47). The oral microbiome may be one of the factors that initiates the systemic inflammatory cascade ((47); present study), while lifestyle and environmental factors that modulate the oral microbiome may explain some of the individual variation in the progression of neurodegeneration. Within the MCI group, *P. intermedia* inversely correlated with mouth rinse nitrate and nitrite concentrations in the APOE4 noncarriers, and there was a greater abundance of *P. intermedia* in APOE4 carriers than in noncarriers. This is reflective of previous findings of higher abundances of *Prevotellaceae* family members in the gut microbiome of APOE4 carriers (48). APOE4 carrier status may therefore be associated with distinct features of both oral and gut microbiomes (48), with important implications for advancing microbiome-targeted preventative therapies. These data also suggest that oral bacteria relative abundances could be used for early detection of risk for cognitive impairment or APOE4 carrier status.

The *Porphyromonas* genus was identified as a risk predictor for MCI. This is consistent with previous studies showing an association between oral *Porphyromonas* and cognitive decline or AD symptoms in rats and humans (11, 49–52). *Porphyromonas* spp. are a driver of periodontal disease and have been shown to promote neuroinflammation and degradation through the activation of inflammatory pathways (53). In mice, *P. gingivalis*

lipopolysaccharide has been shown to impair learning and memory (54). In the present study, *P. gingivalis* was highly correlated with *F. nucleatum* which has been found to co-occur previously (55). *Fusobacterium nucleatum* is a pathogen that acts as a “scaffolding species” co-aggregating with obligate anaerobes in brain abscesses and in biofilm formation in oral infections (56), and the attachment of *P. gingivalis* to human fibroblasts is enhanced by the presence of *F. nucleatum* (57). *Fusobacterium nucleatum* and *P. gingivalis* are both ubiquitous in the oral cavity and are well-known candidates for causing oral inflammation and periodontal disease (58, 59). Moreover, serum antibodies against *F. nucleatum* and *P. intermedia* are elevated in AD compared with healthy controls (47). The present findings of a close co-occurrence of *F. nucleatum* and *P. gingivalis* in the ME4 module, and a significantly elevated *Porphyromonas* in MCI compared with healthy controls, suggest that the co-occurrence of these periodontal pathogens contributes to the development of cognitive impairment.

16S rRNA gene amplicon sequencing has moderate accuracy at species level and, unlike whole genome sequencing, it does not enable verification of nitrate-reductase genes in given OTUs. We relied on a data repository of previously reported bacterial nitrate-reductase genes (<https://www.uniprot.org/>) in interpreting the present data. Future research should assess the consistency of species-level relationships using metagenomics to improve species-level identification accuracy. It is also important to note that, in addition to the relative abundances of oral bacteria assessed in the present study, total bacterial mass may influence relationships between the oral microbiome and host cognitive function. Nevertheless, the novel findings presented herein based on relative abundance data highlight potential targets for pro- and prebiotic interventions and represent a significant advance in understanding how the oral bacterial ecosystem influences cognition in health and in MCI.

The study participants were comprised mostly of females, which could impact the generalizability of the present results. In females, estrogen may confer protection against stroke and cardiovascular disease by enhancing NO production through increasing endothelial NOS expression and activation (60–62). Sex differences have also been reported in the oral microbiome (63). One study found that after dietary nitrate supplementation, the increase in plasma nitrite concentration and the increase in the extent of the chemical reduction of oral nitrate were both greater in females compared with males, but there were no differences in oral microbiome composition (64). Despite evidence that female sex hormones may confer protection against vascular diseases, women have a higher lifetime risk of developing AD with potentially fewer modifiable risk factors (65, 66). More research is required to understand the impact of sex hormones on NO bioavailability and the oral microbiome, and to elucidate the potential consequences for the development of MCI and AD.

The clinical relevance of the present findings is significant, presenting a strong basis for applied research to test the efficacy of nitrate and potentially other nutrients as therapeutic interventions to alter the oral microbiome in MCI and following dementia diagnosis, as well as in APOE4 carriers and noncarriers. We have shown that dietary nitrate supplementation is a powerful intervention in healthy older people to decrease *P. intermedia* and other pathogenic oral bacteria, including *Clostridium difficile*, *P. gingivalis*, *T. denticola*, and *Tannerella forsythia*, and increase *Neisseria* species (35). There is a need to assess the efficacy of dietary nitrate in individuals at the early stages of cognitive decline, where increased dietary nitrate intake may help reverse the rise in oral *Prevotella*:*Neisseria* ratio and thus delay the onset of MCI and AD. Such an intervention may

be particularly important for APOE4 carriers with elevated oral *P. intermedia* preceding dementia diagnosis.

In conclusion, we identified a co-occurring module of oral bacteria dominated by *Neisseria* and *Haemophilus* and associated with oral nitrate reduction via the denitrification pathway, as a positive influence on cognitive outcomes in individuals with MCI. A *Prevotella*-dominated module favoring the DNRA nitrate reduction pathway was associated with low oral NO availability, and *P. intermedia* was revealed as a potential predictor for elevated genetic risk for dementia as indicated by APOE4 status. These results give rise to a novel hypothesis that the balance between two metabolic pathways for nitrate reduction within the oral ecosystem, denitrification and DNRA, is skewed toward DNRA in MCI and potentially modulated by APOE4 status.

## Materials and methods

### Participants

All participants were recruited from the PROTECT study, which is an online aging cohort established for tracking the cognitive health of older adults in the United Kingdom (ethics reference number 13/LO/1578, London Bridge National Health Research Ethics Committee). The present study received approval from the University of Exeter Sport and Health Sciences Ethics Committee (190206-B-03), and all participants provided electronic informed consent as part of a validated online registration to take part.

Inclusion criteria for enrollment in PROTECT at the time of this study were  $\geq 50$  years of age, no diagnosis of dementia, and access to the internet. This study recruited from the PROTECT cohort for whom genetic data were available in addition to longitudinal cognitive data. Additional eligibility criteria excluded any participants using tobacco, antibiotics, or mouthwash. Participants were randomly selected from healthy and MCI groups in the overall PROTECT cohort, defined by standardized thresholds for cognitive performance as described previously (67). APOE4 status was identified (E4E4, E3E4, E2E4, E3E3, and E2E3) and participants were grouped according to genotype.

### Sample collection

The participants were provided with a sample collection kit by post, which included 10 mL of mouthwash (Scope, 15 wt% alcohol, Procter & Gamble) for the self-collection of the oral microbiome and mouth rinse NO biomarker samples. The participants were instructed to collect the mouth rinse sample in the morning and refrain from eating or drinking for at least 2 h prior to sample collection. The participants were asked to swish 10 mL of bactericidal mouthwash (Scope, 15 wt% alcohol, Procter & Gamble) for 30 s. They expectorated the mouth rinse into a universal tube and posted the sample back to the laboratory on the same day as collection. Upon arrival, the samples were immediately stored at  $-80^{\circ}\text{C}$ .

### Oral microbiome 16S rRNA gene amplicon sequencing

The oral bacteria's genomic DNA was extracted from the mouth rinse samples using a Gentra Puregene Buccal Cell Kit (Qiagen, Germantown, MD, USA), according to the manufacturer's instructions. The sample libraries were prepared for sequencing using the NEXTflex 16S V1-V3 Amplicon-Seq Kit (Bioo Scientific, Austin, TX, USA) and sequenced using paired-end 300-bp Illumina MiSeq system, as described previously (68). Sequencing

data were trimmed using Trim-Galore! (Krueger F. Trim-Galore!, [http://www.bioinformatics.babraham.ac.uk/projects/trim\\_galore/](http://www.bioinformatics.babraham.ac.uk/projects/trim_galore/)).

Taxonomic classification of the metagenomic sequencing data was performed using the Kraken2 pipeline with the Kraken2 standard build database with a confidence score 0.05 (69).

## NO biomarkers

Mouth rinse nitrate and nitrite concentrations were measured in 1 mL aliquots using ozone-based chemiluminescence, as previously described (70). The mean nitrate and nitrite concentrations from a mouth rinse blank were subtracted from each sample.

## Cognitive tests

Participants performed a series of cognitive tests using a computerized cognitive test system embedded in the PROTECT study website. They were asked to complete the tests three times within a 7-day window, although this was not mandatory. The tests were measures of working memory (Digit Span, Paired Associates Learning, and Self-ordered Search), executive function (Verbal Reasoning, Switching Stroop), and visual attention (Trail Making), and are described in more detail in previous studies (67, 71, 72). The mean of the total scores for each repeat testing session was used for comparison between the two groups, as previously described (67).

## APOE

APOE genotyping was performed at deCODE Genetics. DNA was extracted from saliva samples collected by post and genotyping was performed using Illumina Global Screening Array with custom content (including directly genotyped single nucleotide polymorphisms, rs429358 and rs7412, to determine APOE status).

## Statistical analysis

All data processing steps were completed in R statistical software (73). The bacterial OTUs were processed by first removing rare OTUs if there were over 20 missing values across all samples and transformed into relative abundances. There were 148 OTUs used in further analysis. Sample outliers were removed if there were <1,000 bacterial reads present. The remaining samples (healthy,  $n = 60$ ; MCI,  $n = 55$ ) were checked for outliers using hierarchical clustering, NMDS, Shannon  $H'$  diversity index, and Chao1 species richness estimate were completed using the R vegan package (74). NMDS of the oral microbiome in healthy and MCI groups and APOE risk groups were based on Bray–Curtis dissimilarity. NMDS ordinations were compared using the ADONIS test in R vegan (74). LEfSe in Conda was used to uncover potential risk predictor OTUs in the samples by comparing the healthy and MCI groups. The samples were normalized, and the alpha value for the factorial Kruskal–Wallis sum-rank test was set to 0.05. The LDA threshold was set to 2.0 and the analysis was completed using the all-against-all “more-strict” method (75).

WGCNA was used to group positively correlated OTUs into modules for analysis against clinical data in a signed network (76). The scale-free topology was calculated for the OTUs. Soft-thresholding power did not reach 0.8, but the data patterns are similar to what has been shown previously (35). Therefore, an arbitrary soft-threshold power was selected according to the recommendations of Langfelder and Horvath (76). Signed correlation networks were used to construct an adjacency matrix for each OTU, where negative correlations were considered unconnected. This adjacency matrix was transformed into a Topological Overlap Matrix and the OTUs were clustered using

hierarchical clustering to produce modules. After the OTUs had been grouped into modules, each module eigenvalue was correlated against NO biomarkers and cognitive test results. Correlation network visualization was performed in Cytoscape (77). The significance level was set at  $\alpha < 0.05$ .

Mouth rinse nitrate and nitrite concentrations were not normally distributed. Therefore, mouth rinse nitrate and nitrite concentrations were compared between Healthy and MCI groups using the Mann–Whitney test, and concentration values are given as the median  $\pm$  IQR.

## Funding

This paper represents independent research part-funded by the National Institute for Health and Care Research as part of the Exeter Biomedical Research Centre (United Kingdom) (Grant Ref: NIHR203320). The views expressed are those of the authors and not necessarily those of the NIHR (United Kingdom) or the Department of Health and Social Care, United Kingdom. It was also supported by the National Institute of Health and Care Research Collaboration for Leadership in Applied Health Research and Care South-West Peninsula, United Kingdom. J.E.L.’s PhD was supported by the University of Exeter Carlotta Palmer studentship. Genotyping was performed at deCODE Genetics. This work was funded in part through the Medical Research Council through the Proximity to Discovery: Industry Engagement Fund [External Collaboration, Innovation and Entrepreneurism: Translational Medicine in Exeter 2 (EXCITEME2), ref. MC\_PC\_17189] awarded to B.C. This project utilized equipment funded by the Wellcome Trust Institutional Strategic Support Fund (WT097835MF), Wellcome Trust Multi User Equipment Award (WT101650MA), and Biotechnology and Biological Sciences Research Council LOLA award (BB/K003240/1).

## Author Contributions

J.E.L.: conceptualization, data curation, software, formal analysis, investigation, visualization, methodology, writing—original draft, review, and editing. A.C.: resources, funding acquisition, project administration, writing—review and editing. C.B.: data curation, funding acquisition, project administration, writing—review and editing. D.V.: writing—review and editing. B.C.: data curation, project administration; writing—review and editing. P.G.W.: writing—review and editing. A.M.J.: supervision, writing—review and editing. A.V.: conceptualization, supervision, funding acquisition, investigation, methodology, project administration, writing—review and editing.

## Data Availability

Sequencing data have been deposited in the National Center for Biotechnology Information Sequence Read Archive Database and are available at BioProject PRJNA1106018. Code used for the analysis is available at <https://github.com/jlhxx>.

## References

- Deary IJ, et al. 2009. Age-associated cognitive decline. *Br Med Bull*. 92:135–152.
- Prince M, et al. 2015. *World Alzheimer report 2015—the global impact of dementia: an analysis of prevalence, incidence, cost and trends*. Alzheimer’s Disease International.

- 3 Livingston G, et al. 2020. Dementia prevention, intervention, and care: 2020 report of the Lancet Commission. *Lancet*. 396:413–446.
- 4 Asher S, Stephen R, Mäntylä P, Suominen AL, Solomon A. 2022. Periodontal health, cognitive decline, and dementia: a systematic review and meta-analysis of longitudinal studies. *J Am Geriatr Soc*. 70:2695–2709.
- 5 Zhang S, et al. 2020. Poor oral health conditions and cognitive decline: studies in humans and rats. *PLoS One*. 15:e0234659.
- 6 Liccardo D, et al. 2020. Potential bidirectional relationship between periodontitis and Alzheimer's disease. *Front Physiol*. 11:683.
- 7 Beydoun MA, et al. 2020. Clinical and bacterial markers of periodontitis and their association with incident all-cause and Alzheimer's disease dementia in a large national survey. *J Alzheimers Dis*. 75:157–172.
- 8 Wu YF, et al. 2021. Oral microbiota changes in elderly patients, an indicator of Alzheimer's disease. *Int J Environ Res Public Health*. 18:4211.
- 9 Parahitiyawa NB, Jin LJ, Leung WK, Yam WC, Samaranyake LP. 2009. Microbiology of odontogenic bacteremia: beyond endocarditis. *Clin Microbiol Rev*. 22:46–64.
- 10 Iida Y, et al. 2004. Brain abscess in which *Porphyromonas gingivalis* was detected in cerebrospinal fluid. *Br J Oral Maxillofac Surg*. 42:180.
- 11 Dominy SS, et al. 2019. *Porphyromonas gingivalis* in Alzheimer's disease brains: evidence for disease causation and treatment with small-molecule inhibitors. *Sci Adv*. 5:eau3333.
- 12 Epstein SE, Zhou YF, Zhu J. 1999. Infection and atherosclerosis: emerging mechanistic paradigms. *Circulation*. 100:e20–e28.
- 13 Yoneda M, et al. 2012. Involvement of a periodontal pathogen, *Porphyromonas gingivalis* on the pathogenesis of non-alcoholic fatty liver disease. *BMC Gastroenterol*. 12:16.
- 14 Rocha BS. 2021. The nitrate-nitrite-nitric oxide pathway on healthy ageing: a review of pre-clinical and clinical data on the impact of dietary nitrate in the elderly. *Front Aging*. 2:778467.
- 15 Cohen RA, et al. 1999. Mechanism of nitric oxide-induced vasodilation: refilling of intracellular stores by sarcoplasmic reticulum Ca<sup>2+</sup> ATPase and inhibition of store-operated Ca<sup>2+</sup> influx. *Circ Res*. 84:210–219.
- 16 Garthwaite J. 1991. Glutamate, nitric oxide and cell-cell signaling in the nervous system. *Trends Neurosci*. 14:60–67.
- 17 Stamler JS, Meissner G. 2001. Physiology of nitric oxide in skeletal muscle. *Physiol Rev*. 81:209–237.
- 18 Moncada S, Higgs A. 1993. The L-arginine-nitric oxide pathway. *N Engl J Med*. 329:2002–2012.
- 19 Lundberg JO, Weitzberg E, Gladwin MT. 2008. The nitrate–nitrite–nitric oxide pathway in physiology and therapeutics. *Nat Rev Drug Discov*. 7:156–167.
- 20 Harooni HE, Naghdi N, Sepehri H, Rohani AH. 2009. The role of hippocampal nitric oxide (NO) on learning and immediate, short- and long-term memory retrieval in inhibitory avoidance task in male adult rats. *Behav Brain Res*. 201:166–172.
- 21 Puzzo D, et al. 2005. Amyloid-beta peptide inhibits activation of the nitric oxide/cGMP/cAMP-responsive element-binding protein pathway during hippocampal synaptic plasticity. *J Neurosci*. 25:6887–6897.
- 22 Wang HG, et al. 2005. Presynaptic and postsynaptic roles of NO, cGK, and RhoA in long-lasting potentiation and aggregation of synaptic proteins. *Neuron*. 45:389–403.
- 23 Pie JE, et al. 2002. Age-related decline of inducible nitric oxide synthase gene expression in primary cultured rat hepatocytes. *Mol Cells*. 13:399–406.
- 24 Berkowitz DE, et al. 2003. Arginase reciprocally regulates nitric oxide synthase activity and contributes to endothelial dysfunction in aging blood vessels. *Circulation*. 108:2000–2006.
- 25 Taddei S, et al. 2001. Age-related reduction of NO availability and oxidative stress in humans. *Hypertension*. 38:274–279.
- 26 Corzo L, Zas R, Rodríguez S, Fernández-Novoa L, Cacabelos R. 2007. Decreased levels of serum nitric oxide in different forms of dementia. *Neurosci Lett*. 420:263–267.
- 27 DiCiero Miranda M, de Bruin VM, Vale MR, Viana GS. 2000. Lipid peroxidation and nitrite plus nitrate levels in brain tissue from patients with Alzheimer's disease. *Gerontology*. 46:179–184.
- 28 Selley ML. 2003. Increased concentrations of homocysteine and asymmetric dimethylarginine and decreased concentrations of nitric oxide in the plasma of patients with Alzheimer's disease. *Neurobiol Aging*. 24:903–907.
- 29 Hultman K, Strickland S, Norris EH. 2013. The APOE  $\epsilon$ 4/ $\epsilon$ 4 genotype potentiates vascular fibrin(ogen) deposition in amyloid-laden vessels in the brains of Alzheimer's disease patients. *J Cereb Blood Flow Metab*. 33:1251–1258.
- 30 Biffi A, et al. 2011. APOE genotype and extent of bleeding and outcome in lobar intracerebral haemorrhage: a genetic association study. *Lancet Neurol*. 10:702–709.
- 31 Chen X, Miller NM, Afghah Z, Geiger JD. 2019. Development of AD-like pathology in skeletal muscle. *J Parkinsons Dis Alzheimers Dis*. 6:10.13188/2376-922x.1000028.
- 32 Linton MF, Fazio S. 1999. Macrophages, lipoprotein metabolism, and atherosclerosis: insights from murine bone marrow transplantation studies. *Curr Opin Lipidol*. 10:97–105.
- 33 Montagne A, et al. 2021. APOE4 accelerates advanced-stage vascular and neurodegenerative disorder in old Alzheimer's mice via cyclophilin A independently of amyloid- $\beta$ . *Nat Aging*. 1:506–520.
- 34 Morou-Bermúdez E, Torres-Colón JE, Bermúdez NS, Patel RP, Joshupura KJ. 2022. Pathways linking oral bacteria, nitric oxide metabolism, and health. *J Dent Res*. 101:623–631.
- 35 Vanhatalo A, et al. 2021. Network analysis of nitrate-sensitive oral microbiome reveals interactions with cognitive function and cardiovascular health across dietary interventions. *Redox Biol*. 41:101933.
- 36 Doel JJ, Benjamin N, Hector MP, Rogers M, Allaker RP. 2005. Evaluation of bacterial nitrate reduction in the human oral cavity. *Eur J Oral Sci*. 113:14–19.
- 37 De Groote MA, Fang FC. 1995. NO inhibitions: antimicrobial properties of nitric oxide. *Clin Infect Dis*. 21 Suppl 2:S162–S165.
- 38 Allaker RP, Silva Mendez LS, Hardie JM, Benjamin N. 2001. Antimicrobial effect of acidified nitrite on periodontal bacteria. *Oral Microbiol Immunol*. 16:253–256.
- 39 Takeshita T, et al. 2016. Bacterial diversity in saliva and oral health-related conditions: the Hisayama Study. *Sci Rep*. 6:22164.
- 40 De Filippis F, et al. 2014. The same microbiota and a potentially discriminant metabolome in the saliva of omnivore, ovo-lacto-vegetarian and vegan individuals. *PLoS One*. 9:e112373.
- 41 Espinoza JL, et al. 2022. Differential network analysis of oral microbiome metatranscriptomes identifies community scale metabolic restructuring in dental caries. *PNAS Nexus*. 1:pgac239.
- 42 Chen H, et al. 2015. A Filifactor alocis-centered co-occurrence group associates with periodontitis across different oral habitats. *Sci Rep*. 5:9053.
- 43 Einarsson GG, et al. 2019. Community analysis and co-occurrence patterns in airway microbial communities during health and disease. *ERJ Open Res*. 5:00128–2017.
- 44 Kroll JS, Wilks KE, Farrant JL, Langford PR. 1998. Natural genetic exchange between *Haemophilus* and *Neisseria*: intergeneric

- transfer of chromosomal genes between major human pathogens. *Proc Natl Acad Sci U S A*. 95:12381–12385.
- 45 Akiyama H, et al. 2000. Inflammation and Alzheimer's disease. *Neurobiol Aging*. 21:383–421.
  - 46 Jiao Y, Hasegawa M, Inohara N. 2014. The role of oral pathobionts in dysbiosis during periodontitis development. *J Dent Res*. 93: 539–546.
  - 47 Sparks Stein P, et al. 2012. Serum antibodies to periodontal pathogens are a risk factor for Alzheimer's disease. *Alzheimers Dement*. 8:196–203.
  - 48 Tran TTT, et al. 2019. APOE genotype influences the gut microbiome structure and function in humans and mice: relevance for Alzheimer's disease pathophysiology. *FASEB J*. 33: 8221–8231.
  - 49 Chi L, et al. 2021. *Porphyromonas gingivalis*-induced cognitive impairment is associated with gut dysbiosis, neuroinflammation, and glymphatic dysfunction. *Front Cell Infect Microbiol*. 11:755925.
  - 50 Choi S, et al. 2019. Association of chronic periodontitis on Alzheimer's disease or vascular dementia. *J Am Geriatr Soc*. 67: 1234–1239.
  - 51 Ding Y, Ren J, Yu H, Yu W, Zhou Y. 2018. *Porphyromonas gingivalis*, a periodontitis causing bacterium, induces memory impairment and age-dependent neuroinflammation in mice. *Immun Ageing*. 15:6.
  - 52 Hu Y, et al. 2020. Periodontitis induced by *P. gingivalis*-LPS is associated with neuroinflammation and learning and memory impairment in Sprague-Dawley rats. *Front Neurosci*. 14:658.
  - 53 Costa MJF, et al. 2021. Relationship of *Porphyromonas gingivalis* and Alzheimer's disease: a systematic review of pre-clinical studies. *Clin Oral Investig*. 25:797–806.
  - 54 Zhang J, et al. 2018. *Porphyromonas gingivalis* lipopolysaccharide induces cognitive dysfunction, mediated by neuronal inflammation via activation of the TLR4 signaling pathway in C57BL/6 mice. *J Neuroinflammation*. 15:37.
  - 55 Li J, et al. 2014. Comparative analysis of the human saliva microbiome from different climate zones: Alaska, Germany, and Africa. *BMC Microbiol*. 14:316.
  - 56 Kommedal Ø, et al. 2014. Massive parallel sequencing provides new perspectives on bacterial brain abscesses. *J Clin Microbiol*. 52:1990–1997.
  - 57 Metzger Z, Blasbalg J, Dotan M, Weiss EI. 2009. Enhanced attachment of *Porphyromonas gingivalis* to human fibroblasts mediated by *Fusobacterium nucleatum*. *J Endod*. 35:82–85.
  - 58 Kistler JO, Booth V, Bradshaw DJ, Wade WG. 2013. Bacterial community development in experimental gingivitis. *PLoS One*. 8: e71227.
  - 59 Rafiei M, et al. 2017. Study of *Porphyromonas gingivalis* in periodontal diseases: a systematic review and meta-analysis. *Med J Islam Repub Iran*. 31:62.
  - 60 Hurn PD, Macrae IM. 2000. Estrogen as a neuroprotectant in stroke. *J Cereb Blood Flow Metab*. 20:631–652.
  - 61 Dubey RK, Jackson EK. 2001. Cardiovascular protective effects of 17 $\beta$ -estradiol metabolites. *J Appl Physiol* (1985). 91:1868–1883.
  - 62 Chambliss KL, Shaul PW. 2002. Estrogen modulation of endothelial nitric oxide synthase. *Endocr Rev*. 23:665–686.
  - 63 Liu X, et al. 2023. Sex differences in the oral microbiome, host traits, and their causal relationships. *iScience*. 26:105839.
  - 64 Kapil V, et al. 2018. Sex differences in the nitrate-nitrite-NO( $\bullet$ ) pathway: role of oral nitrate-reducing bacteria. *Free Radic Biol Med*. 126:113–121.
  - 65 Nebel RA, et al. 2018. Understanding the impact of sex and gender in Alzheimer's disease: a call to action. *Alzheimers Dement*. 14: 1171–1183.
  - 66 Anstey KJ, et al. 2021. Association of sex differences in dementia risk factors with sex differences in memory decline in a population-based cohort spanning 20–76 years. *Sci Rep*. 11:7710.
  - 67 Brooker H, et al. 2020. FLAME: a computerized neuropsychological composite for trials in early dementia. *Alzheimers Dement (Amst)*. 12:e12098.
  - 68 Vanhatalo A, et al. 2018. Nitrate-responsive oral microbiome modulates nitric oxide homeostasis and blood pressure in humans. *Free Radic Biol Med*. 124:21–30.
  - 69 Wood DE, Lu J, Langmead B. 2019. Improved metagenomic analysis with Kraken 2. *Genome Biol*. 20:257.
  - 70 Kelly J, et al. 2013. Effects of short-term dietary nitrate supplementation on blood pressure, O<sub>2</sub> uptake kinetics, and muscle and cognitive function in older adults. *Am J Physiol Regul Integr Comp Physiol*. 304:R73–R83.
  - 71 Ward G, Roberts MJ, Phillips LH. 2001. Task-switching costs, Stroop-costs, and executive control: a correlational study. *Q J Exp Psychol A*. 54:491–511.
  - 72 Parsons OA, Maslow HI, Morris F, Denny JP. 1964. Trail-making test performance in relation to certain experimenter, test and subject variables. *Percept Mot Skills*. 19:199–206.
  - 73 R Core Team. 2020. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.
  - 74 Oksanen J, et al. 2022. vegan: Community Ecology Package. R package version 2.6-8. <https://CRAN.Rproject.org/package=vegan>.
  - 75 Segata N, et al. 2011. Metagenomic biomarker discovery and explanation. *Genome Biol*. 12:R60.
  - 76 Langfelder P, Horvath S. 2008. WGCNA: an R package for weighted correlation network analysis. *BMC Bioinformatics*. 9: 559.
  - 77 Shannon P, et al. 2003. Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res*. 13:2498–2504.