

Search for dark matter produced in association with a pair of bottom quarks in proton-proton collisions at $\sqrt{s} = 13$ TeV



The CMS collaboration

E-mail: cms-publication-committee-chair@cern.ch

ABSTRACT: A search for dark matter (DM) particles produced in association with bottom quarks is presented. The analysis uses proton-proton collision data at a center-of-mass energy of $\sqrt{s} = 13$ TeV, corresponding to an integrated luminosity of 138 fb^{-1} . The search is performed in a final state with large missing transverse momentum and a pair of jets originating from bottom quarks. No significant excess of data is observed with respect to the standard model expectation. Results are interpreted in the context of a type-II two-Higgs-doublet model with an additional light pseudoscalar (2HDM+a). An upper limit is set on the mass of the lighter pseudoscalar, probing masses up to 260 GeV at 95% confidence level. Sensitivity to the parameter space with the ratio of the vacuum expectation values of the two Higgs doublets, $\tan \beta$, greater than 15 is achieved, capitalizing on the enhancement of couplings between pseudoscalars and bottom quarks with high $\tan \beta$.

KEYWORDS: Dark Matter, Hadron-Hadron Scattering

ARXIV EPRINT: [2408.17336](https://arxiv.org/abs/2408.17336)

Contents

1	Introduction	1
2	Signal and background simulations	2
3	The CMS detector	4
4	Data samples and event reconstruction	4
5	Analysis strategy and event selection	5
6	Background estimation	7
7	Signal extraction	8
8	Systematic uncertainties	9
9	Results	10
10	Summary	12
	The CMS collaboration	19

1 Introduction

So far, the evidence for the existence of dark matter (DM) has come from astrophysics. According to the lambda cold dark matter (Λ CDM) model of cosmology [1], the high-precision Planck satellite map of the cosmic microwave background [2] indicates that DM constitutes about 26% of the total mass-energy of the known universe [3], however, the particle nature of DM has not been elucidated yet. If it is composed of subnuclear particles, then these could potentially be produced at the LHC. Such particles, whether elementary or composite, are weakly interacting with standard model (SM) particles, which means they might escape detection in collider experiments. Since these DM candidates are expected to be electrically neutral and stable, their presence could be inferred through an observed imbalance in the transverse momentum vectors of particles in the detector, known as missing transverse momentum ($\vec{p}_T^{\text{miss}}$) with magnitude represented by p_T^{miss} .

Many beyond-the-SM (BSM) theories hypothesize the particle nature of DM [4, 5]. Based on astrophysical and cosmological indications and the observed relic density of the universe, weakly interacting massive particles (WIMPs) [6], with masses in the range from a few GeV to a few TeV, are a popular type of DM candidates that can be searched for at colliders. Such searches can be performed by looking for the pair production of WIMPs through a mediator that recoils against one or more visible SM particles. WIMPs can give a detector signature of the type $X + p_T^{\text{miss}}$ [7], where X is one or more SM particles, e.g. a heavy-quark pair, produced in association with DM.

In this paper, we perform a search for DM particles produced in association with a pair of bottom quarks ($b\bar{b} + p_T^{\text{miss}}$) using the proton-proton (pp) collision data collected with the CMS detector in 2016–2018, corresponding to an integrated luminosity of 138 fb^{-1} . The search for DM produced in association with heavy-flavor quarks is complementary to other $X + p_T^{\text{miss}}$ searches and to searches for DM production in cascades in the context of supersymmetry (SUSY) at the LHC. The ATLAS and CMS Collaborations have performed DM searches using heavy-flavor jets as tags [7, 8], such as b-jets from single top quarks ($t/\bar{t}$) [9–12], b-jets from top quark pairs ($t\bar{t}$) [13, 14], or bottom quark pairs ($b\bar{b}$) [15–20].

2 Signal and background simulations

The model used for the interpretation of the results of this analysis is a type-II two-Higgs-doublet model with an additional pseudoscalar singlet interacting with DM particles, referred to as 2HDM+a [21, 22]. The model contains a scalar Higgs boson (h , assumed to be the observed Higgs boson with mass $m_h = 125 \text{ GeV}$), a pair of charged Higgs bosons ($H^\pm$), a heavy scalar boson (H), a heavy pseudoscalar boson (A), a light pseudoscalar boson (a), and a Dirac fermion (χ), which is assumed to be the DM particle. Figure 1 shows the leading Feynman diagrams for the production of the $b\bar{b} + \chi\bar{\chi}$ process within the 2HDM+a model.

The choice of the fixed and free parameters in this model follows the recommendations of ref. [23]. The model contains the following parameters: the electroweak vacuum expectation value ($v \approx 246 \text{ GeV}$), the mass of the DM candidate m_χ , the masses of the Higgs bosons ($m_h, m_H, m_{H^\pm}$), the masses of the pseudoscalar bosons (m_a, m_A), the mixing angle between the two CP-even weak spin-0 eigenstates (α), the mixing angle between the two CP-odd weak spin-0 eigenstates (θ), the ratio of the vacuum expectation values of the two Higgs doublets ($\tan\beta$), and the quartic couplings ($\lambda_3, \lambda_{P1}, \lambda_{P2}$) of the scalar potential, which governs the interactions between the scalar doublets and the pseudoscalar.

In the decoupling limit, where $\sin(\beta - \alpha) = 1$ and $\lambda_3 = \lambda_{P1} = \lambda_{P2} = 3$ ensure the boundedness of the Higgs potential, electroweak precision constraints give $m_A = m_H = m_{H^\pm}$, and the DM coupling y_χ is set to 1. This results in five free parameters, namely, $\tan\beta$, $\sin\theta$, m_A , m_a , and m_χ . This choice is adopted in many 2HDM+a interpretations of heavy spin-0 resonances searches performed at the LHC [17, 24–27]. Furthermore, the motivation for using the 2HDM+a model is that the couplings of both the heavier pseudoscalar ($g_{Ab\bar{b}}$) and the lighter pseudoscalar ($g_{ab\bar{b}}$) to bottom quarks are proportional to $\tan\beta$, resulting in increased interaction rates in the high- $\tan\beta$ region.

The production cross section is dominantly driven by the mass of the lighter mediator, a , as well as $\tan\beta$ and $\sin\theta$, with no strong dependence on the mass of A . Furthermore, as shown in figure 2, the distribution of p_T^{miss} strongly depends on m_a . On the other hand, $\tan\beta$, $\sin\theta$, and m_χ marginally change the p_T^{miss} distribution, as also shown in figure 2 for m_χ . Hence, signal events are generated with $m_A = 600 \text{ GeV}$, $\tan\beta = 35$, $\sin\theta = 0.7$, and $m_\chi = 1 \text{ GeV}$ for different m_a values using MADGRAPH5_aMC@NLO v2.6.0 [28] and the 5-flavor scheme at leading order (LO) in quantum chromodynamics (QCD) with no additional jets at the matrix element level.

A lower threshold of 150 GeV is set on p_T^{miss} at the generator level to constrain the kinematic phase space of simulated events to the region where the analysis is conducted. Various




Figure 1. Leading Feynman diagrams for the $b\bar{b} + \chi\bar{\chi}$ process in the 2HDM+a model.




Figure 2. Normalized (to unity) shape of generator-level p_T^{miss} distribution for two illustrative lighter pseudoscalar masses m_a (left) and for five illustrative DM masses m_χ (right).

SM processes have a similar final state and higher production cross sections. The dominant background comes from the $Z(\nu\nu) + \text{jets}$ production ($\approx 60\text{--}70\%$ of the total background), when the Z boson is produced in association with one or two b jets and decays to neutrinos. The $t\bar{t}$ (lepton + jets) and $W(\ell\nu) + \text{jets}$ backgrounds are the next largest backgrounds, where a lepton coming from these processes is either lost or not reconstructed properly in the detector. Dileptonic and hadronic $t\bar{t}$ decays, as well as single top quark and diboson (WW , WZ , ZZ) production events, are smaller contributions to the total background.

The $Z(\nu\nu) + \text{jets}$, $Z(\ell\ell) + \text{jets}$, and $W(\ell\nu) + \text{jets}$ background processes ($\ell = e, \mu, \tau$) are simulated using MADGRAPH5_aMC@NLO v2.4.2 [28] at next-to-LO (NLO) in QCD. The $t\bar{t}$ and single top quark production events are simulated using the POWHEG v2 generator [29, 30] at NLO in QCD. Diboson events are generated at LO in QCD using the PYTHIA 8.212 generator [31]. The SM Higgs boson with a mass of 125 GeV produced in association with a vector boson is considered as a background and is simulated using the POWHEG

generator at NLO in QCD. The NNPDF3.1 parton distribution functions (PDFs) [32] are used, depending on the QCD order of the generator for each physics process. Parton showering, fragmentation, and hadronization are simulated with PYTHIA 8.230 (8.212 for 2016) using the CP5 (CUETP8M1 for 2016) underlying event tune [33, 34].

Interactions of the resulting final state particles with the CMS detector are simulated using the GEANT4 software [35] and reconstructed using the same algorithms used for the data. Additional inelastic pp interactions in the same or a neighboring bunch crossing (pileup) are included in the simulation. The pileup distribution is adjusted to match the corresponding distribution observed in data.

3 The CMS detector

The central feature of the CMS apparatus is a superconducting solenoid of 6 m internal diameter, providing a magnetic field of 3.8 T. Within the solenoid volume are a silicon pixel and strip tracker, a lead tungstate crystal electromagnetic calorimeter (ECAL), and a brass and scintillator hadron calorimeter (HCAL), each composed of a barrel and two endcap sections. Forward calorimeters, made of steel and quartz fibers, extend the pseudorapidity (η) coverage provided by the barrel and endcap detectors. Muons are detected in gas-ionization chambers embedded in the steel flux-return yoke outside the solenoid.

Events of interest are selected using a two-tiered trigger system [36]. The first level, composed of custom hardware processors (L1), uses information from the calorimeters and muon detectors to select events at a rate of around 100 kHz in a fixed time interval of approximately $4\ \mu\text{s}$. The second level, known as the high-level trigger (HLT), consists of a farm of processors running a version of the full event reconstruction software optimized for fast processing, and reduces the event rate to around 1 kHz before data storage.

A more detailed description of the CMS detector, along with a definition of the coordinate system used and the relevant kinematic variables, can be found in ref. [37].

4 Data samples and event reconstruction

This search uses data collected in pp collisions between 2016 and 2018. In particular, physics data sets for which the events pass the $p_{\text{T}}^{\text{miss}}$ and single-electron triggers at the HLT, and electron-photon and calo- $p_{\text{T}}^{\text{miss}}$ triggers at L1 are used for final analysis.

A global “particle-flow” (PF) algorithm [38] aims to reconstruct all individual particles in an event, combining information provided by the inner tracker ECAL, HCAL, and muon detectors. The reconstructed particles are used to build τ leptons, jets, and missing transverse momentum [39–41]. The primary vertex is taken to be the vertex corresponding to the hardest scattering in the event, evaluated using tracking information alone, as described in section 9.4.1 of ref. [42].

Jets are clustered using the anti- k_{T} jet clustering algorithm [43] with a distance parameter ($\Delta R = \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2}$) of 0.4 (“AK4 jets”). Jet calibration [40] is used to account for the energy deposits of neutral particles and full nonlinear hadronic response of the detector. Pileup effects from the charged hadrons are mitigated by removing them from pileup interactions, with a technique called “charged-hadron subtraction” (CHS) [44]. For mitigation of neutral

PU particles, a new technique, pileup per particle identification (PUPPI) [45] is used, which builds on the existing CHS algorithm. Furthermore, to ensure that the energy and momentum of the reconstructed jet matches that of the quark/gluon-initiated jet, jet energy corrections are applied to the reconstructed jets. In this search, jets with $p_T > 30 \text{ GeV}$ and $|\eta| < 2.5$ are considered. To remove any overlap with leptons, all jets within $\Delta R = 0.4$ with leptons are removed.

Jets originating from bottom quark hadronization, b jets, are identified with the DEEPCSV algorithm [46], which uses a deep neural network (DNN) with 4 hidden layers and exploits the discriminating power of variables such as the position of the secondary vertex, the lifetime of the bottom quark, and the mass of b hadrons. Three working points are defined on the basis of the output of the DeepCSV algorithm. The loose working point corresponds to a 10% misidentification (“mistag”) rate with 84% efficiency. The medium working point gives a 1% mistag rate with 68% efficiency, and the tight working point gives a 0.1% mistag rate with 50% efficiency.

The hadron-plus-strips algorithm is employed to reconstruct hadronically decaying τ leptons, denoted as τ_h [39]. Jets reconstructed using AK4 jet clustering are used as an input to the algorithm and τ_h candidates are reconstructed on the basis of the number of charged hadrons and energy deposits in strips in the ECAL, which originate from the photon candidates and their electron-positron conversions. In order to minimize the misidentification rate of τ_h from quark and gluon jets, a convolutional discriminator based on a DNN called DeepTau [47] is employed. Events with a τ_h candidate with $p_T > 20 \text{ GeV}$ and $|\eta| < 2.3$ are vetoed.

Shower shape and isolation variables, as defined in section 7.1 of ref. [48], are used to identify electrons and photons. Isolated electrons and photons used in this analysis have $p_T > 10$ and 15 GeV , respectively, and $|\eta| < 2.5$. Isolated muons have $p_T > 10 \text{ GeV}$ and $|\eta| < 2.4$. Two parameters are considered while identifying the muons, namely the number of measurements in the tracker-muon system and the track reconstruction quality [49].

As mentioned in section 1, the negative vectorial p_T sum of all PF candidates is used to calculate $\vec{p}_T^{\text{miss}}$ with all jet energy corrections are applied. Events with anomalous $\vec{p}_T^{\text{miss}}$ can occur because of reconstruction failures, detector artifacts, or noncollision backgrounds [41]. Dedicated event filters, which can identify more than 90% of such events, are used, with a mistagging rate of less than 0.1%.

5 Analysis strategy and event selection

The final states reconstructed in this analysis consist of one or two b-tagged jets and large p_T^{miss} , which could arise from DM particles. The data are split into two categories based on the number of b-tagged jets: the 1b category contains events with only one b-tagged jet and is composed of one signal region (SR), called SR1, and two control regions (CRs), while the 2b category contains events with two b-tagged jets and is composed of one SR (SR2) and two CRs. Events that contain at least one well identified and isolated electron or muon with $p_T > 10 \text{ GeV}$ are vetoed from the SRs to reduce the background from the $t\bar{t}$ and $W(\ell\nu) + \text{jets}$ processes. These regions are illustrated in figure 3. The three dominant background sources $Z(\nu\nu) + \text{jets}$, $t\bar{t}(\ell\nu)$, and $W(\ell\nu) + \text{jets}$ are estimated by performing a simultaneous fit of the SRs and CRs, as detailed in section 7.




Figure 3. Definition of SR1 and SR2 in terms of jet multiplicity (left), and CR and SR definition in the lepton multiplicity and additional jet multiplicity plane (right). The categories depend on the number and flavor of leptons, and the number of jets in an event.

In the CRs, p_T^{miss} is replaced by the absolute value of the transverse momentum recoil, $\vec{U}$, to mimic the variation observed in p_T^{miss} within the SRs because of the presence of leptons. The recoil is defined as:

$$U = |\vec{U}| = \left| -\left(\vec{p}_T^{\text{miss}} + \sum \vec{p}_T^{\text{lep}} \right) \right| \quad (5.1)$$

In the 2b category, the p_T^{miss} distribution for signal events is similar to that of background events, such that there is little separation power for signal and background. The angular variable $\cos \Theta^*$ [50], defined as

$$\cos \Theta^* = \left| \tanh \left(\frac{\eta_1 - \eta_2}{2} \right) \right|, \quad (5.2)$$

where η_1 and η_2 are the pseudorapidities of the leading and subleading jets, gives an optimal discrimination between the signal and backgrounds in this category. In the longitudinally boosted frame, $\cos \Theta^*$ is the cosine of the polar angle between each jet and the beam axis, where the pseudorapidities of the jets are equal and opposite.

Events in the SRs and CRs are selected such that they are mutually exclusive. The events in the SRs and muon-based CRs are selected by a logical ‘OR’ of p_T^{miss} triggers with different thresholds (90, 100, 110, 120, or 140 GeV), whereas events in the electron-based CRs are selected by using a logical ‘OR’ of single electron triggers with minimum threshold of 27, 32, or 35 GeV [36]. Events are preselected using a set of requirements common to the SRs and CRs. In the CRs, the selection criteria incorporate U to mimic the variation observed in p_T^{miss} within the SRs because of the presence of leptons.

The event selection procedure starts with the preselection, which requires events passing trigger thresholds, followed by an offline selection of p_T^{miss} or $U > 250$ GeV. Events must include at least one jet, with the leading jet having $p_T > 100$ GeV, as optimized based on

Selection	Cut Value
Trigger	p_T^{miss} or single electron
p_T^{miss} (U in CR)	>250 GeV
Jets	multiplicity ≥ 1 , $p_T > 30$ GeV, $ \eta < 2.5$
Leading jet	$p_T > 100$ GeV
Overlap removal	no overlap with e, μ , τ_h , γ
γ	veto photon with $p_T > 15$ GeV and $ \eta < 2.5$
$\min(\Delta\phi(\text{jet}, \vec{p}_T^{\text{miss}}))$	>0.5
Δp_T^{miss} (PF - calorimeter)	<0.5

Table 1. Preselection criteria applied to all the events entering the SRs and CRs.

signal sensitivity. Events with a photon with $p_T > 15$ GeV and $|\eta| < 2.5$ are rejected to suppress photon-induced backgrounds ($Z(\ell\ell) + \gamma + \text{jet}$, $W(\ell\nu) + \gamma + \text{jet}$).

QCD multijet events can mimic the signal signature when the jet momentum is mismeasured, giving rise to high p_T^{miss} . The QCD multijet background is reduced by requiring the minimum $\Delta\phi$ between any jet and $\vec{p}_T^{\text{miss}}$ ($\min(\Delta\phi(\text{jet}, \vec{p}_T^{\text{miss}}))$) to be higher than 0.5 radians. This ensures that $\vec{p}_T^{\text{miss}}$ is not aligned with any jet, which is a common feature in events with mismeasured jets. The QCD multijet background is further reduced by requiring the relative difference in the value of p_T^{miss} measured with the PF algorithm and with calorimeter only information to be less than 0.5. This quantity is defined as

$$\Delta p_T^{\text{miss}}(\text{PF} - \text{calorimeter}) = \begin{cases} \left| p_T^{\text{miss}}(\text{PF})/p_T^{\text{miss}}(\text{calorimeter}) - 1 \right|, & \text{in the SRs and} \\ \left| p_T^{\text{miss}}(\text{PF})/U(\text{calorimeter}) - 1 \right|, & \text{in the CRs.} \end{cases}$$

Table 1 summarizes the requirements to preselect the events for all three data-taking years. A dedicated event selection for the SRs and CRs is applied on top of the preselection.

Events in SR2 have two jets tagged as originating from bottom quarks (using the medium working point) and large p_T^{miss} . Furthermore, up to one additional jet with $p_T > 30$ GeV, not b-tagged, is allowed in events entering this region. SR1 is foreseen to catch signal events where one of the b jets is lost because it is outside of the detector acceptance or is not tagged correctly, requiring exactly one b-tagged jet and up to one additional jet. This category helps to recover some lost signal efficiency at the cost of a relatively higher background.

6 Background estimation

To estimate the $Z(\nu\nu) + \text{jets}$ background, two CRs, targeting the $Z \rightarrow e^-e^+$ and $Z \rightarrow \mu^-\mu^+$ processes, are defined. In these CRs, out of exactly two leptons in the event, the leading lepton must have $p_T > 30$ GeV, the invariant mass of the dilepton candidate must be between 70 and 110 GeV, and there is no b tagging requirement on the jets. The CRs dedicated to the $W(\ell\nu) + \text{jets}$ and $t\bar{t}$ backgrounds are built for SR1 and SR2, respectively. They require a single lepton (electron or muon) with $p_T > 30$ GeV and a transverse mass $m_T = \sqrt{2p_T^{\text{miss}}p_T^l(1 - \cos[\Delta\phi(\vec{p}_T^{\text{miss}}, \vec{p}_T^l)])}$ less than 160 GeV, together with the same b tagging requirements as in the SRs. In addition, events in the $t\bar{t}$ CR have at least 1 additional




Figure 4. QCD background contribution in the QCD CR (black and pink dots) in the 2b category using 2017 data.

non b-tagged jet, while, for the $W(\ell\nu) + \text{jets}$ CR, exactly one b-tagged jet is required and p_T^{miss} must be greater than 100 GeV.

To estimate the QCD multijet background, a QCD CR is defined using the same selection as the SR except for requiring $\min(\Delta\phi(\text{jet}, \vec{p}_T^{\text{miss}})) < 0.5$. The QCD background in the CR is estimated as the difference between data and other simulated backgrounds, and is plotted against the $\min(\Delta\phi)$ threshold, i.e., for each entry of $\min(\Delta\phi)$ on the x axis, the y axis corresponds to the number of events passing the $\min(\Delta\phi)$ requirement. This distribution is plotted for each bin of p_T^{miss} or $\cos\Theta^*$ and then fitted with a function that describes the behavior of the distribution. The QCD contribution exhibits an exponential decrease with $\min(\Delta\phi)$, as shown for one example in figure 4. The QCD background is fitted with an exponential in the $\min(\Delta\phi)$ range between 0 to 0.3, and the resulting function is validated in the $\min(\Delta\phi)$ range between 0.3 to 0.5. The QCD contribution in the SR is calculated as the integral of the function for $\min(\Delta\phi(\text{jet}, \vec{p}_T^{\text{miss}})) > 0.5$.

7 Signal extraction

The fit for the 1b category is performed using p_T^{miss} in SR1 and U in corresponding CRs as observables, therefore, both SR1 and the corresponding CRs enter simultaneously in a binned profile likelihood fit to estimate the signal. Each bin of p_T^{miss} in SR1 is tied to the corresponding bin of U in the CR via a scale factor, treating the systematic uncertainties as nuisance parameters in the fit. A variable binning is used for p_T^{miss} and U with bin edges 250, 300, 400, 550, and 1000 GeV.

In the 2b category, the variable $\cos\Theta^*$ is used with bins of 0.25 width from 0.0 to 1.0, and a similar procedure is used to constrain the backgrounds as shown in figure 6 (Z and $t\bar{t}$) using equivalent bins in the CRs. For each category, the electron and muon CRs are combined to form a single CR for each background estimated from data.

Furthermore, the fit to the data determines the background post-fit yields in each bin, as shown in the lower panels of figure 5 and 6, displaying the ratio of the yields in data over the prediction, before and after the fit. The pre-fit yield values are determined by running the selection on simulated events to determine selection efficiency, and normalizing them to the integrated luminosity times their respective theoretical cross sections. The post-fit values are obtained as the pre-fit yields adjusted by the post-fit values of the nuisance parameters and scale factors. The lower panels of figure 5 and 6 show that the post-fit background distributions agree significantly better with the data in the control regions, confirming that the method is working.

8 Systematic uncertainties

The background and signal predictions are affected by systematic uncertainties. Only the systematic uncertainties that have an impact of at least half a percent are taken into account in the signal extraction fit. Uncertainties that do not impact the shape of the distribution, or are uniformly distributed, are treated as normalization uncertainties, whereas those that exhibit non-uniform variations are treated as shape and normalization uncertainties (except for scale and pdf uncertainties, where only shape has been taken into account).

Shape-based uncertainties in the b tagging efficiency are calculated separately for true b jets (having b flavor at the particle level) and incorrectly b -tagged jets for each event.

The uncertainty in the integrated luminosity amounts to 1.2–2.5%, depending on the data-taking year [51–53], while the overall uncertainty for the 2016–2018 period is 1.6%.

The p_T^{miss} -based trigger efficiency scale factors are measured in a phase space with single-muon and dimuon events, and the difference in the scale factors is considered as the systematic uncertainty. A shape-based systematic uncertainty of the order of 1% is applied to all the simulated samples in the SRs and muon CRs. The electron trigger efficiency scale factors are calculated using a tag-and-probe method [54] and a shape-based uncertainty is applied to all the simulated samples in the electron CRs. The overall effect is of the order of 2–3% on the normalization.

Shape uncertainties arising from jet energy corrections are applied to the simulations, separating different sources of uncertainty. The average effect is of the order of 2–3%.

An uncertainty in the total inelastic cross section of 4.6% [55] is used as a shape-based uncertainty to account for the effects of pileup modeling.

For the single top quark, diboson, and SM Higgs boson processes, a normalization uncertainty of 10%, 20%, and 20%, respectively, is used [56–58].

The systematic uncertainty in the cross section due to the renormalization and factorization scales is taken into account for signal and all backgrounds, using the envelope of the distributions treated as the shape uncertainty. The uncertainty related to the modeling of PDFs is estimated using the replicas provided in the corresponding PDF sets for the signal and different backgrounds [32, 59–61].

For the QCD multijet background, the total uncertainty comes from the combination of the uncertainty obtained from the 95% confidence intervals of the fitted extrapolation function and a flat uncertainty of 20% that covers the statistical uncertainty. The total effect on the QCD background estimation is about 50%. The maximum variations of the

Systematics	Type	Size
Pileup reweighting	shape+ lnN	3%
b tag efficiency	shape+ lnN	8%
Incorrectly b-tagged jets	shape+ lnN	7%
Electron identification	shape+ lnN	7%
Electron reconstruction efficiency	shape+ lnN	5%
Muon identification	shape+ lnN	<1%
Muon isolation	shape+ lnN	<1%
Electron trigger	shape+ lnN	1%
p_T^{miss} trigger	shape+ lnN	1%
Jet energy correction	shape+ lnN	3%
L1 prefiring	shape+ lnN	0.5%
Muon prefiring	lnN	0.5%
τ_h veto	lnN	6%
Integrated luminosity	lnN	2.5%
*Cross section diboson	lnN	20%
*Cross section Drell-Yan	lnN	5%
*Cross section Higgs boson	lnN	20%
*Cross section single top quark	lnN	10%
*QCD normalization	lnN	50%
Renormalization and factorization scales	shape	<42%
PDF	shape	<50%

Table 2. Maximum variations from systematic uncertainties, affecting the shape and normalization (lnN) of the signal and background predictions. Only the systematic sources with an impact greater than 0.5% are listed. *refers to corresponding individual processes.

sources of systematic uncertainties across individual bins and processes, along with their respective categories, is detailed in table 2.

9 Results

Figures 5 and 6 present the expected distributions of p_T^{miss} for SR1 and U for the $Z(\ell\ell) + \text{jets}$ and $W(\ell\nu) + \text{jets}$ CRs in the 1b category, as well as $\cos\Theta^*$ for the SR and CRs in the 2b category. These distributions are shown both before (prefit) and after (postfit) the maximum likelihood fit, with the observed data superimposed. No excess with respect to the SM prediction is observed.

Upper limits on the production crosssection of DM candidates in association with two bottom quarks as predicted by the theoretical model, σ /, are extracted at the 95% confidence level (CL) using a modified frequentist method [62–64] computed with an asymptotic approximation [65]. The combined limit is shown in figure 7. The observed exclusion range for m_a is up to 260 GeV, for $m_A = 600$ GeV, $\tan\beta = 35$, $\sin\theta = 0.7$, and $m_\chi = 1$ GeV.

The two-dimensional limit on signal strength in the plane of m_a - $\tan\beta$, using $m_A = 600$ GeV, $\sin\theta = 0.7$ and $m_\chi = 1$ GeV, is shown in figure 8 (upper left). The two-dimensional limit in the plane of m_a - $\sin\theta$, using $m_A = 600$ GeV, $\tan\beta = 35$ and $m_\chi = 1$ GeV, is shown in figure 8 (upper right). The two-dimensional limit in the plane of m_a - m_χ , using $\tan\beta = 35$, $\sin\theta = 0.7$, and $m_A = 600$ GeV is shown in figure 8 (lower).

Tabulated results are provided in the HEPData record for this analysis [66].




Figure 5. Observed and predicted p_T^{miss} (U) distributions in the 1b category. The red lines divide the plot into three parts: the first part shows the SR, the second part the $Z(\ell\ell) + \text{jets}$ CR, and the third part the $W(\ell\nu) + \text{jets}$ CR. The bottom plot shows the ratio of observed data to predicted distributions for both postfit and prefit predictions. The uncertainty band represents the combined systematic and statistical uncertainties in the predictions. Prefit points are shown with their combined systematic and statistical uncertainties.




Figure 6. Observed and predicted $\cos\Theta^*$ distributions in the 2b category. The red lines divide the plot into three parts: the first part shows the SR, the second part the $Z(\nu\nu) + \text{jets}$ CR, and the third part the $t\bar{t}$ CR. The bottom plot shows the ratio of observed data to predicted distributions for both postfit and prefit predictions. The uncertainty band represents the combined systematic and statistical uncertainties in the predictions. Prefit points are shown with their combined systematic and statistical uncertainties.




Figure 7. The 95% CL upper limit on the production cross-section of DM candidates in association with a pair of bottom quarks for $m_A = 600$ GeV, $\sin \theta = 0.7$, $m_\chi = 1$ GeV, and $\tan \beta = 35$, for the combination of SR1 and SR2. The green and yellow bands show the ± 1 and ± 2 standard deviations from expected limits. The mass points below the red line are excluded.

10 Summary

A search for dark matter produced in association with a pair of bottom quarks is performed using data collected by the CMS detector in 2016–2018, corresponding to an integrated luminosity of 138 fb^{-1} . The analysis searches for a possible signal by using two independent categories with different multiplicities of jets reconstructed as originating from a bottom quark. The results are interpreted in the framework of a simplified model, namely a type-II two-Higgs-doublet model with an additional pseudoscalar singlet (2HDM+a). It is the first search at the LHC to probe dark matter (DM) produced in association with two nonresonant bottom quarks in this model. This search performs best in the high- $\tan \beta$ phase space, where the signal production cross section is enhanced. This provides a complementary search to constrain the 2HDM+a model parameter phase space in the region where other DM searches are less sensitive. The lighter pseudoscalar mass below 260 GeV is excluded at 95% confidence level for $\tan \beta = 35$, $\sin \theta = 0.7$, and $m_A = 600$ GeV.

Acknowledgments

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centers and personnel of the Worldwide LHC Computing Grid and other centers for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC, the CMS detector, and the supporting computing infrastructure provided by the following funding agencies: SC (Armenia), BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil);




Figure 8. Observed and expected upper limits at 95% CL on the signal strength in the m_a — $\tan\beta$ (upper left), m_a — $\sin\theta$ (upper right), and m_a — m_χ (lower) planes. The shaded area bounded by solid black line is excluded, with, the dotted grey line ($m_a = 2m_\chi$) separating on-shell and off-shell productions for the lower plot.

MES and BNSF (Bulgaria); CERN; CAS, MoST, and NSFC (China); MINCIENCIAS (Colombia); MSES and CSF (Croatia); RIF (Cyprus); SENESCYT (Ecuador); ERC PRG, RVTT3 and MoER TK202 (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); SRNSF (Georgia); BMBF, DFG, and HGF (Germany); GSRI (Greece); NKFIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIP and NRF (Republic of Korea); MES (Latvia); LMTLT (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MES and NSC (Poland); FCT

(Portugal); MESTD (Serbia); MCIN/AEI and PCTI (Spain); MOSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); MHESI and NSTDA (Thailand); TUBITAK and TENMAK (Turkey); NASU (Ukraine); STFC (United Kingdom); DOE and NSF (U.S.A.).

Individuals have received support from the Marie-Curie program and the European Research Council and Horizon 2020 Grant, contract Nos. 675440, 724704, 752730, 758316, 765710, 824093, 101115353, 101002207, and COST Action CA16108 (European Union); the Leventis Foundation; the Alfred P. Sloan Foundation; the Alexander von Humboldt Foundation; the Science Committee, project no. 22rl-037 (Armenia); the Belgian Federal Science Policy Office; the Fonds pour la Formation à la Recherche dans l’Industrie et dans l’Agriculture (FRIA-Belgium); the F.R.S.-FNRS and FWO (Belgium) under the “Excellence of Science — EOS” — be.h project n. 30820817; the Beijing Municipal Science & Technology Commission, No. Z191100007219010 and Fundamental Research Funds for the Central Universities (China); the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Shota Rustaveli National Science Foundation, grant FR-22-985 (Georgia); the Deutsche Forschungsgemeinschaft (DFG), among others, under Germany’s Excellence Strategy — EXC 2121 “Quantum Universe” — 390833306, and under project number 400140256 — GRK2497; the Hellenic Foundation for Research and Innovation (HFRI), Project Number 2288 (Greece); the Hungarian Academy of Sciences, the New National Excellence Program — ÚNKP, the NKFIH research grants K 131991, K 133046, K 138136, K 143460, K 143477, K 146913, K 146914, K 147048, 2020-2.2.1-ED-2021-00181, and TKP2021-NKTA-64 (Hungary); the Council of Science and Industrial Research, India; ICSC — National Research Center for High Performance Computing, Big Data and Quantum Computing and FAIR — Future Artificial Intelligence Research, funded by the NextGenerationEU program (Italy); the Latvian Council of Science; the Ministry of Education and Science, project no. 2022/WK/14, and the National Science Center, contracts Opus 2021/41/B/ST2/01369 and 2021/43/B/ST2/01552 (Poland); the Fundação para a Ciência e a Tecnologia, grant CEECIND/01334/2018 (Portugal); the National Priorities Research Program by Qatar National Research Fund; MCIN/AEI/10.13039/501100011033, ERDF “a way of making Europe”, and the Programa Estatal de Fomento de la Investigación Científica y Técnica de Excelencia María de Maeztu, grant MDM-2017-0765 and Programa Severo Ochoa del Principado de Asturias (Spain); the Chulalongkorn Academic into Its 2nd Century Project Advancement Project, and the National Science, Research and Innovation Fund via the Program Management Unit for Human Resources & Institutional Development, Research and Innovation, grant B39G670016 (Thailand); the Kavli Foundation; the Nvidia Corporation; the SuperMicro Corporation; the Welch Foundation, contract C-1845; and the Weston Havens Foundation (U.S.A.).

Data Availability Statement. Release and preservation of data used by the CMS Collaboration as the basis for publications is guided by the [CMS data preservation, re-use, and open access policy](#).

Code Availability Statement. The CMS core software is publicly available on [GitHub](#).

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] N. Deruelle and J.-P. Uzan, *The Λ CDM model of the hot Big Bang*, in *Relativity in modern physics*, Oxford University Press, Oxford, U.K. (2018), p. 605–610 [[DOI:10.1093/oso/9780198786399.003.0059](https://doi.org/10.1093/oso/9780198786399.003.0059)].
- [2] PLANCK collaboration, *Planck 2018 results. I. Overview and the cosmological legacy of Planck*, *Astron. Astrophys.* **641** (2020) A1 [[arXiv:1807.06205](https://arxiv.org/abs/1807.06205)] [[INSPIRE](#)].
- [3] K. Freese, *Status of dark matter in the universe*, *Int. J. Mod. Phys.* **1** (2017) 325 [[arXiv:1701.01840](https://arxiv.org/abs/1701.01840)] [[INSPIRE](#)].
- [4] J.L. Feng, *Dark matter candidates from particle physics and methods of detection*, *Ann. Rev. Astron. Astrophys.* **48** (2010) 495 [[arXiv:1003.0904](https://arxiv.org/abs/1003.0904)] [[INSPIRE](#)].
- [5] V. Trimble, *Existence and nature of dark matter in the universe*, *Ann. Rev. Astron. Astrophys.* **25** (1987) 425 [[INSPIRE](#)].
- [6] G. Jungman, M. Kamionkowski and K. Griest, *Supersymmetric dark matter*, *Phys. Rept.* **267** (1996) 195 [[hep-ph/9506380](https://arxiv.org/abs/hep-ph/9506380)] [[INSPIRE](#)].
- [7] CMS collaboration, *Dark sector searches with the CMS experiment*, [arXiv:2405.13778](https://arxiv.org/abs/2405.13778) [[INSPIRE](#)].
- [8] ATLAS collaboration, *Constraints on simplified dark matter models involving an s-channel mediator with the ATLAS detector in pp collisions at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **84** (2024) 1102 [[arXiv:2404.15930](https://arxiv.org/abs/2404.15930)] [[INSPIRE](#)].
- [9] CMS collaboration, *Search for dark matter produced in association with a single top quark or a top quark pair in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **03** (2019) 141 [[arXiv:1901.01553](https://arxiv.org/abs/1901.01553)] [[INSPIRE](#)].
- [10] ATLAS collaboration, *Search for dark matter produced in association with a single top quark in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector*, *Eur. Phys. J. C* **81** (2021) 860 [[arXiv:2011.09308](https://arxiv.org/abs/2011.09308)] [[INSPIRE](#)].
- [11] ATLAS collaboration, *Search for dark matter produced in association with a single top quark and an energetic W boson in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector*, *Eur. Phys. J. C* **83** (2023) 603 [[arXiv:2211.13138](https://arxiv.org/abs/2211.13138)] [[INSPIRE](#)].
- [12] ATLAS collaboration, *Search for new particles in final states with a boosted top quark and missing transverse momentum in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *JHEP* **05** (2024) 263 [[arXiv:2402.16561](https://arxiv.org/abs/2402.16561)] [[INSPIRE](#)].
- [13] CMS collaboration, *Search for dark matter particles produced in association with a top quark pair at $\sqrt{s} = 13$ TeV*, *Phys. Rev. Lett.* **122** (2019) 011803 [[arXiv:1807.06522](https://arxiv.org/abs/1807.06522)] [[INSPIRE](#)].
- [14] ATLAS collaboration, *Constraints on spin-0 dark matter mediators and invisible Higgs decays using ATLAS 13 TeV pp collision data with two top quarks and missing transverse momentum in the final state*, *Eur. Phys. J. C* **83** (2023) 503 [[arXiv:2211.05426](https://arxiv.org/abs/2211.05426)] [[INSPIRE](#)].
- [15] CMS collaboration, *Search for dark matter produced in association with heavy-flavor quark pairs in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **77** (2017) 845 [[arXiv:1706.02581](https://arxiv.org/abs/1706.02581)] [[INSPIRE](#)].

- [16] CMS collaboration, *Searches for physics beyond the standard model with the M_{T2} variable in hadronic final states with and without disappearing tracks in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **80** (2020) 3 [[arXiv:1909.03460](#)] [[INSPIRE](#)].
- [17] CMS collaboration, *Search for dark matter produced in association with a Higgs boson decaying to a pair of bottom quarks in proton-proton collisions at $\sqrt{s} = 13$ TeV*, *Eur. Phys. J. C* **79** (2019) 280 [[arXiv:1811.06562](#)] [[INSPIRE](#)].
- [18] ATLAS collaboration, *Search for dark matter produced in association with bottom or top quarks in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector*, *Eur. Phys. J. C* **78** (2018) 18 [[arXiv:1710.11412](#)] [[INSPIRE](#)].
- [19] ATLAS collaboration, *Search for new phenomena in final states with b-jets and missing transverse momentum in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector*, *JHEP* **05** (2021) 093 [[arXiv:2101.12527](#)] [[INSPIRE](#)].
- [20] ATLAS collaboration, *Search for dark matter produced in association with a Standard Model Higgs boson decaying into b-quarks using the full run 2 dataset from the ATLAS detector*, *JHEP* **11** (2021) 209 [[arXiv:2108.13391](#)] [[INSPIRE](#)].
- [21] D. Abercrombie et al., *Dark matter benchmark models for early LHC run-2 searches: report of the ATLAS/CMS dark matter forum*, *Phys. Dark Univ.* **27** (2020) 100371 [[arXiv:1507.00966](#)] [[INSPIRE](#)].
- [22] M. Bauer, U. Haisch and F. Kahlhoefer, *Simplified dark matter models with two Higgs doublets. Part I. Pseudoscalar mediators*, *JHEP* **05** (2017) 138 [[arXiv:1701.07427](#)] [[INSPIRE](#)].
- [23] LHC DARK MATTER WORKING GROUP collaboration, *LHC dark matter working group: next-generation spin-0 dark matter models*, *Phys. Dark Univ.* **27** (2020) 100351 [[arXiv:1810.09420](#)] [[INSPIRE](#)].
- [24] ATLAS collaboration, *Search for heavy ZZ resonances in the $\ell^+\ell^-\ell^+\ell^-$ and $\ell^+\ell^-\nu\bar{\nu}$ final states using proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Eur. Phys. J. C* **78** (2018) 293 [[arXiv:1712.06386](#)] [[INSPIRE](#)].
- [25] ATLAS collaboration, *Search for heavy resonances decaying into WW in the $e\nu\mu\nu$ final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector*, *Eur. Phys. J. C* **78** (2018) 24 [[arXiv:1710.01123](#)] [[INSPIRE](#)].
- [26] CMS collaboration, *Search for beyond the standard model Higgs bosons decaying into a $b\bar{b}$ pair in pp collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **08** (2018) 113 [[arXiv:1805.12191](#)] [[INSPIRE](#)].
- [27] ATLAS collaboration, *Constraints on mediator-based dark matter and scalar dark energy models using $\sqrt{s} = 13$ TeV pp collision data collected by the ATLAS detector*, *JHEP* **05** (2019) 142 [[arXiv:1903.01400](#)] [[INSPIRE](#)].
- [28] J. Alwall et al., *The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations*, *JHEP* **07** (2014) 079 [[arXiv:1405.0301](#)] [[INSPIRE](#)].
- [29] S. Frixione, P. Nason and C. Oleari, *Matching NLO QCD computations with parton shower simulations: the POWHEG method*, *JHEP* **11** (2007) 070 [[arXiv:0709.2092](#)] [[INSPIRE](#)].
- [30] S. Alioli, P. Nason, C. Oleari and E. Re, *A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG BOX*, *JHEP* **06** (2010) 043 [[arXiv:1002.2581](#)] [[INSPIRE](#)].
- [31] T. Sjöstrand et al., *An introduction to PYTHIA 8.2*, *Comput. Phys. Commun.* **191** (2015) 159 [[arXiv:1410.3012](#)] [[INSPIRE](#)].

- [32] NNPDF collaboration, *Parton distributions for the LHC run II*, *JHEP* **04** (2015) 040 [[arXiv:1410.8849](#)] [[INSPIRE](#)].
- [33] CMS collaboration, *Event generator tunes obtained from underlying event and multiparton scattering measurements*, *Eur. Phys. J. C* **76** (2016) 155 [[arXiv:1512.00815](#)] [[INSPIRE](#)].
- [34] P. Skands, S. Carrazza and J. Rojo, *Tuning PYTHIA 8.1: the Monash 2013 tune*, *Eur. Phys. J. C* **74** (2014) 3024 [[arXiv:1404.5630](#)] [[INSPIRE](#)].
- [35] GEANT4 collaboration, *GEANT4 — a simulation toolkit*, *Nucl. Instrum. Meth. A* **506** (2003) 250 [[INSPIRE](#)].
- [36] CMS collaboration, *The CMS trigger system*, *2017 JINST* **12** P01020 [[arXiv:1609.02366](#)] [[INSPIRE](#)].
- [37] CMS collaboration, *The CMS experiment at the CERN LHC*, *2008 JINST* **3** S08004 [[INSPIRE](#)].
- [38] CMS collaboration, *Particle-flow reconstruction and global event description with the CMS detector*, *2017 JINST* **12** P10003 [[arXiv:1706.04965](#)] [[INSPIRE](#)].
- [39] CMS collaboration, *Performance of reconstruction and identification of τ leptons decaying to hadrons and ν_τ in pp collisions at $\sqrt{s} = 13$ TeV*, *2018 JINST* **13** P10005 [[arXiv:1809.02816](#)] [[INSPIRE](#)].
- [40] CMS collaboration, *Jet energy scale and resolution in the CMS experiment in pp collisions at 8 TeV*, *2017 JINST* **12** P02014 [[arXiv:1607.03663](#)] [[INSPIRE](#)].
- [41] CMS collaboration, *Performance of missing transverse momentum reconstruction in proton-proton collisions at $\sqrt{s} = 13$ TeV using the CMS detector*, *2019 JINST* **14** P07004 [[arXiv:1903.06078](#)] [[INSPIRE](#)].
- [42] D. Contardo et al., *Technical proposal for the phase-II upgrade of the CMS detector*, CERN-LHCC-2015-010, CERN, Geneva, Switzerland (2015) [[DOI:10.17181/CERN.VU8I.D59J](#)] [[INSPIRE](#)].
- [43] M. Cacciari, G.P. Salam and G. Soyez, *The anti- k_t jet clustering algorithm*, *JHEP* **04** (2008) 063 [[arXiv:0802.1189](#)] [[INSPIRE](#)].
- [44] CMS collaboration, *Pileup mitigation at CMS in 13 TeV data*, *2020 JINST* **15** P09018 [[arXiv:2003.00503](#)] [[INSPIRE](#)].
- [45] D. Bertolini, P. Harris, M. Low and N. Tran, *Pileup per particle identification*, *JHEP* **10** (2014) 059 [[arXiv:1407.6013](#)] [[INSPIRE](#)].
- [46] CMS collaboration, *Identification of heavy-flavour jets with the CMS detector in pp collisions at 13 TeV*, *2018 JINST* **13** P05011 [[arXiv:1712.07158](#)] [[INSPIRE](#)].
- [47] CMS collaboration, *Identification of hadronic tau lepton decays using a deep neural network*, *2022 JINST* **17** P07023 [[arXiv:2201.08458](#)] [[INSPIRE](#)].
- [48] CMS collaboration, *Electron and photon reconstruction and identification with the CMS experiment at the CERN LHC*, *2021 JINST* **16** P05014 [[arXiv:2012.06888](#)] [[INSPIRE](#)].
- [49] CMS collaboration, *Performance of the CMS muon detector and muon reconstruction with proton-proton collisions at $\sqrt{s} = 13$ TeV*, *2018 JINST* **13** P06015 [[arXiv:1804.04528](#)] [[INSPIRE](#)].
- [50] A.J. Barr, *Measuring slepton spin at the LHC*, *JHEP* **02** (2006) 042 [[hep-ph/0511115](#)] [[INSPIRE](#)].

- [51] CMS collaboration, *Precision luminosity measurement in proton-proton collisions at $\sqrt{s} = 13$ TeV in 2015 and 2016 at CMS*, *Eur. Phys. J. C* **81** (2021) 800 [[arXiv:2104.01927](#)] [[INSPIRE](#)].
- [52] CMS collaboration, *CMS luminosity measurement for the 2017 data-taking period at $\sqrt{s} = 13$ TeV*, *CMS-PAS-LUM-17-004*, CERN, Geneva, Switzerland (2018).
- [53] CMS collaboration, *CMS luminosity measurement for the 2018 data-taking period at $\sqrt{s} = 13$ TeV*, *CMS-PAS-LUM-18-002*, CERN, Geneva, Switzerland (2019).
- [54] CMS collaboration, *Measurements of inclusive W and Z cross sections in pp collisions at $\sqrt{s} = 7$ TeV*, *JHEP* **01** (2011) 080 [[arXiv:1012.2466](#)] [[INSPIRE](#)].
- [55] CMS collaboration, *Measurement of the inelastic proton-proton cross section at $\sqrt{s} = 13$ TeV*, *JHEP* **07** (2018) 161 [[arXiv:1802.02613](#)] [[INSPIRE](#)].
- [56] CMS collaboration, *Observation of the associated production of a single top quark and a W boson in pp collisions at $\sqrt{s} = 8$ TeV*, *Phys. Rev. Lett.* **112** (2014) 231802 [[arXiv:1401.2942](#)] [[INSPIRE](#)].
- [57] CMS collaboration, *Measurement of the ZZ production cross section and $Z \rightarrow \ell^+ \ell^- \ell'^+ \ell'^-$ branching fraction in pp collisions at $\sqrt{s} = 13$ TeV*, *Phys. Lett. B* **763** (2016) 280 [Erratum *ibid.* **772** (2017) 884] [[arXiv:1607.08834](#)] [[INSPIRE](#)].
- [58] CMS collaboration, *Measurement of the inclusive and differential WZ production cross sections, polarization angles, and triple gauge couplings in pp collisions at $\sqrt{s} = 13$ TeV*, *JHEP* **07** (2022) 032 [[arXiv:2110.11231](#)] [[INSPIRE](#)].
- [59] J. Butterworth et al., *PDF4LHC recommendations for LHC run II*, *J. Phys. G* **43** (2016) 023001 [[arXiv:1510.03865](#)] [[INSPIRE](#)].
- [60] S. Dulat et al., *New parton distribution functions from a global analysis of quantum chromodynamics*, *Phys. Rev. D* **93** (2016) 033006 [[arXiv:1506.07443](#)] [[INSPIRE](#)].
- [61] L.A. Harland-Lang, A.D. Martin, P. Motylinski and R.S. Thorne, *Parton distributions in the LHC era: MMHT 2014 PDFs*, *Eur. Phys. J. C* **75** (2015) 204 [[arXiv:1412.3989](#)] [[INSPIRE](#)].
- [62] LHC HIGGS CROSS SECTION WORKING GROUP collaboration, *Handbook of LHC Higgs cross sections: 3. Higgs properties*, [arXiv:1307.1347](#) [[DOI:10.5170/CERN-2013-004](#)] [[INSPIRE](#)].
- [63] T. Junk, *Confidence level computation for combining searches with small statistics*, *Nucl. Instrum. Meth. A* **434** (1999) 435 [[hep-ex/9902006](#)] [[INSPIRE](#)].
- [64] A.L. Read, *Presentation of search results: the CL_s technique*, *J. Phys. G* **28** (2002) 2693 [[INSPIRE](#)].
- [65] G. Cowan, K. Cranmer, E. Gross and O. Vitells, *Asymptotic formulae for likelihood-based tests of new physics*, *Eur. Phys. J. C* **71** (2011) 1554 [Erratum *ibid.* **73** (2013) 2501] [[arXiv:1007.1727](#)] [[INSPIRE](#)].
- [66] *HEPData record for this analysis*, <https://doi.org/10.17182/hepdata.152385>, (2024).

The CMS collaboration

Yerevan Physics Institute, Yerevan, Armenia

A. Hayrapetyan, A. Tumasyan ¹

Institut für Hochenergiephysik, Vienna, Austria

W. Adam , J.W. Andrejkovic , T. Bergauer , S. Chatterjee , K. Damanakis , M. Dragicevic ,
P.S. Hussain , M. Jeitler ², N. Krammer , A. Li , D. Liko , I. Mikulec , J. Schieck ²,
R. Schöfbeck , D. Schwarz , M. Sonawane , W. Waltenberger , C.-E. Wulz ²

Universiteit Antwerpen, Antwerpen, Belgium

T. Janssen , T. Van Laer , P. Van Mechelen 

Vrije Universiteit Brussel, Brussel, Belgium

N. Breugelmans , J. D'Hondt , S. Dansana , A. De Moor , M. Delcourt , F. Heyen ,
S. Lowette , I. Makarenko , D. Müller , S. Tavernier , M. Tytgat ³, G.P. Van Onsem ,
S. Van Putte , D. Vannerom 

Université Libre de Bruxelles, Bruxelles, Belgium

B. Bilin , B. Clerbaux , A.K. Das , G. De Lentdecker , H. Evard , L. Favart , P. Gianneios ,
J. Jaramillo , A. Khalilzadeh , F.A. Khan , K. Lee , M. Mahdavihorrani , A. Malara ,
S. Paredes , M.A. Shahzad , L. Thomas , M. Vanden Bemden , C. Vander Velde , P. Vanlaer 

Ghent University, Ghent, Belgium

M. De Coen , D. Dobur , G. Gokbulut , Y. Hong , J. Knolle , L. Lambrecht , D. Marckx ,
K. Mota Amarilo , A. Samalan , K. Skovpen , N. Van Den Bossche , J. van der Linden ,
L. Wezenbeek 

Université Catholique de Louvain, Louvain-la-Neuve, Belgium

A. Benecke , A. Bethani , G. Bruno , C. Caputo , J. De Favereau De Jeneret , C. Delaere ,
I.S. Donertas , A. Giammanco , A.O. Guzel , Sa. Jain , V. Lemaitre , J. Lidrych ,
P. Mastrapasqua , T.T. Tran , S. Wertz 

Centro Brasileiro de Pesquisas Fisicas, Rio de Janeiro, Brazil

G.A. Alves , M. Alves Gallo Pereira , E. Coelho , G. Correia Silva , C. Hensel ,
T. Menezes De Oliveira , C. Mora Herrera ⁴, A. Moraes , P. Rebello Teles , M. Soeiro ,
A. Vilela Pereira ⁴

Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

W.L. Aldá Júnior , M. Barroso Ferreira Filho , H. Brandao Malbouisson , W. Carvalho ,
J. Chinellato⁵, E.M. Da Costa , G.G. Da Silveira ⁶, D. De Jesus Damiao ,
S. Fonseca De Souza , R. Gomes De Souza , M. Macedo , J. Martins ⁷, L. Mundim ,
H. Nogima , J.P. Pinheiro , A. Santoro , A. Sznajder , M. Thiel 

Universidade Estadual Paulista, Universidade Federal do ABC, São Paulo, Brazil

C.A. Bernardes ⁶, L. Calligaris , T.R. Fernandez Perez Tomei , E.M. Gregores ,
I. Maietto Silverio , P.G. Mercadante , S.F. Novaes , B. Orzari , Sandra S. Padula 

Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria

A. Aleksandrov , G. Antchev , R. Hadjiiska , P. Iaydjiev , M. Misheva , M. Shopova ,
G. Sultanov 

University of Sofia, Sofia, Bulgaria

A. Dimitrov , L. Litov , B. Pavlov , P. Petkov , A. Petrov , E. Shumka 

Instituto De Alta Investigación, Universidad de Tarapacá, Casilla 7 D, Arica, Chile

S. Keshri , D. Laroze , S. Thakur 

Beihang University, Beijing, China

T. Cheng , T. Javaid , L. Yuan 

Department of Physics, Tsinghua University, Beijing, China

Z. Hu , Z. Liang, J. Liu, K. Yi ^{8,9}

Institute of High Energy Physics, Beijing, China

G.M. Chen ¹⁰, H.S. Chen ¹⁰, M. Chen ¹⁰, F. Iemmi , C.H. Jiang, A. Kapoor ¹¹, H. Liao ,
Z.-A. Liu ¹², R. Sharma ¹³, J.N. Song¹², J. Tao , C. Wang¹⁰, J. Wang , Z. Wang¹⁰, H. Zhang ,
J. Zhao 

State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China

A. Agapitos , Y. Ban , S. Deng , B. Guo, C. Jiang , A. Levin , C. Li , Q. Li , Y. Mao,
S. Qian, S.J. Qian , X. Qin, X. Sun , D. Wang , H. Yang, L. Zhang , Y. Zhao, C. Zhou 

Guangdong Provincial Key Laboratory of Nuclear Science and Guangdong-Hong Kong Joint Laboratory of Quantum Matter, South China Normal University, Guangzhou, China

S. Yang 

Sun Yat-Sen University, Guangzhou, China

Z. You 

University of Science and Technology of China, Hefei, China

K. Jaffel , N. Lu 

Nanjing Normal University, Nanjing, China

G. Bauer¹⁴, B. Li, J. Zhang 

Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) — Fudan University, Shanghai, China

X. Gao¹⁵, Y. Li

Zhejiang University, Hangzhou, Zhejiang, China

Z. Lin, C. Lu, M. Xiao

Universidad de Los Andes, Bogota, Colombia

C. Avila, D.A. Barbosa Trujillo, A. Cabrera, C. Florez, J. Fraga, J.A. Reyes Vega

Universidad de Antioquia, Medellin, Colombia

F. Ramirez, C. Rendón, M. Rodriguez, A.A. Ruales Barbosa, J.D. Ruiz Alvarez

University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia

D. Giljanovic, N. Godinovic, D. Lelas, A. Sculac

University of Split, Faculty of Science, Split, Croatia

M. Kovac, A. Petkovic, T. Sculac

Institute Rudjer Boskovic, Zagreb, Croatia

P. Bargassa, V. Brigljevic, B.K. Chitroda, D. Ferencek, K. Jakovcic, S. Mishra, A. Starodumov¹⁶, T. Susa

University of Cyprus, Nicosia, Cyprus

A. Attikis, K. Christoforou, A. Hadjiagapiou, C. Leonidou, J. Mousa, C. Nicolaou, L. Paizanos, F. Ptochos, P.A. Razis, H. Rykaczewski, H. Saka, A. Stepennov

Charles University, Prague, Czech Republic

M. Finger, M. Finger Jr., A. Kveton

Universidad San Francisco de Quito, Quito, Ecuador

E. Carrera Jarrin

Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt

A.A. Abdelalim^{17,18}, S. Elgammal¹⁹, A. Ellithi Kamel²⁰

Center for High Energy Physics (CHEP-FU), Fayoum University, El-Fayoum, Egypt

M. Abdullah Al-Mashad, M.A. Mahmoud

National Institute of Chemical Physics and Biophysics, Tallinn, Estonia

K. Ehataht, M. Kadastik, T. Lange, S. Nandan, C. Nielsen, J. Pata, M. Raidal, L. Tani, C. Veelken

Department of Physics, University of Helsinki, Helsinki, Finland

H. Kirschenmann, K. Osterberg, M. Voutilainen

Helsinki Institute of Physics, Helsinki, Finland

S. Bharthuar , N. Bin Norjoharuddeen , E. Brücken , F. Garcia , P. Inkaew ,
K.T.S. Kallonen , T. Lampén , K. Lassila-Perini , S. Lehti , T. Lindén , L. Martikainen ,
M. Myllymäki , M.m. Rantanen , H. Siikonen , J. Tuominiemi 

Lappeenranta-Lahti University of Technology, Lappeenranta, Finland

P. Luukka , H. Petrow 

IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France

M. Besancon , F. Couderc , M. Dejardin , D. Denegri, J.L. Faure, F. Ferri , S. Ganjour ,
P. Gras , G. Hamel de Monchenault , M. Kumar , V. Lohezic , J. Malcles , F. Orlandi ,
L. Portales , A. Rosowsky , M.Ö. Sahin , A. Savoy-Navarro ²¹, P. Simkina , M. Titov ,
M. Tornago 

Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France

F. Beaudette , G. Boldrini , P. Busson , A. Cappati , C. Charlot , M. Chiusi , F. Damas ,
O. Davignon , A. De Wit , I.T. Ehle , B.A. Fontana Santos Alves , S. Ghosh , A. Gilbert ,
R. Granier de Cassagnac , A. Hakimi , B. Harikrishnan , L. Kalipoliti , G. Liu , M. Nguyen ,
C. Ochando , R. Salerno , J.B. Sauvan , Y. Sirois , L. Urda Gómez , E. Vernazza ,
A. Zabi , A. Zghiche 

Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France

J.-L. Agram ²², J. Andrea , D. Apparu , D. Bloch , J.-M. Brom , E.C. Chabert ,
C. Collard , S. Falke , U. Goerlach , R. Haeberle , A.-C. Le Bihan , M. Meena , O. Poncet ,
G. Saha , M.A. Sessini , P. Van Hove , P. Vaucelle 

Centre de Calcul de l'Institut National de Physique Nucleaire et de Physique des Particules, CNRS/IN2P3, Villeurbanne, France

A. Di Florio 

Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France

D. Amram, S. Beauceron , B. Blancon , G. Boudoul , N. Chanon , D. Contardo ,
P. Depasse , C. Dozen ²³, H. El Mamouni, J. Fay , S. Gascon , M. Gouzevitch , C. Greenberg,
G. Grenier , B. Ille , E. Jourdhuy, I.B. Laktineh, M. Lethuillier , L. Mirabito, S. Perries,
A. Purohit , M. Vander Donckt , P. Verdier , J. Xiao 

Georgian Technical University, Tbilisi, Georgia

A. Khvedelidze ¹⁶, I. Lomidze , Z. Tsamalaidze ¹⁶

RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany

V. Botta , S. Consuegra Rodríguez , L. Feld , K. Klein , M. Lipinski , D. Meuser ,
A. Pauls , D. Pérez Adán , N. Röwert , M. Teroerde 

RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

S. Diekmann , A. Dodonova , N. Eich , D. Eliseev , F. Engelke , J. Erdmann ,
M. Erdmann , P. Fackeldey , B. Fischer , T. Hebbeker , K. Hoepfner , F. Ivone , A. Jung 

M.y. Lee , F. Mausolf , M. Merschmeyer , A. Meyer , S. Mukherjee , D. Noll , F. Nowotny, A. Pozdnyakov , Y. Rath, W. Redjeb , F. Rehm, H. Reithler , V. Sarkisovi , A. Schmidt , C. Seth, A. Sharma , J.L. Spah , A. Stein , F. Torres Da Silva De Araujo ²⁴, S. Wiedenbeck , S. Zaleski

RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany

C. Dziwok , G. Flügge , T. Kress , A. Nowack , O. Pooth , A. Stahl , T. Ziemons , A. Zotz 

Deutsches Elektronen-Synchrotron, Hamburg, Germany

H. Aarup Petersen , M. Aldaya Martin , J. Alimena , S. Amoroso, Y. An , J. Bach , S. Baxter , M. Bayatmakou , H. Becerril Gonzalez , O. Behnke , A. Belvedere , F. Blekman ²⁵, K. Borras ²⁶, A. Campbell , A. Cardini , C. Cheng, F. Colombina , M. De Silva , G. Eckerlin, D. Eckstein , L.I. Estevez Banos , O. Filatov , E. Gallo ²⁵, A. Geiser , V. Guglielmi , M. Guthoff , A. Hinzmann , L. Jeppe , B. Kaech , M. Kasemann , C. Kleinwort , R. Kogler , M. Komm , D. Krücker , W. Lange, D. Leyva Pernia , K. Lipka ²⁷, W. Lohmann ²⁸, F. Lorkowski , R. Mankel , I.-A. Melzer-Pellmann , M. Mendizabal Morentin , A.B. Meyer , G. Milella , K. Moral Figueroa , A. Mussgiller , L.P. Nair , J. Niedziela , A. Nürnberg , Y. Otarid, J. Park , E. Ranken , A. Raspereza , D. Rastorguev , J. Rübenach, L. Rygaard, A. Saggio , M. Scham ^{29,26}, S. Schnake ²⁶, P. Schütze , C. Schwanenberger ²⁵, D. Selivanova , K. Sharko , M. Shchedrolosiev , D. Stafford, F. Vazzoler , A. Ventura Barroso , R. Walsh , D. Wang , Q. Wang , Y. Wen , K. Wichmann, L. Wiens ²⁶, C. Wissing , Y. Yang , A. Zimmermann Castro Santos 

University of Hamburg, Hamburg, Germany

A. Albrecht , S. Albrecht , M. Antonello , S. Bein , L. Benato , S. Bollweg, M. Bonanomi , P. Connor , K. El Morabit , Y. Fischer , E. Garutti , A. Grohsjean , J. Haller , H.R. Jabusch , G. Kasieczka , P. Keicher, R. Klanner , W. Korcari , T. Kramer , C.c. Kuo, V. Kutzner , F. Labe , J. Lange , A. Lobanov , C. Matthies , L. Moureaux , M. Mrowietz, A. Nigamova , Y. Nissan, A. Paasch , K.J. Pena Rodriguez , T. Quadfasel , B. Raciti , M. Rieger , D. Savoie , J. Schindler , P. Schleper , M. Schröder , J. Schwandt , M. Sommerhalder , H. Stadie , G. Steinbrück , A. Tews, M. Wolf 

Karlsruher Institut fuer Technologie, Karlsruhe, Germany

S. Brommer , M. Burkart, E. Butz , T. Chwalek , A. Dierlamm , A. Droll, U. Elicabuk, N. Faltermann , M. Giffels , A. Gottmann , F. Hartmann ³⁰, R. Hofsaess , M. Horzela , U. Husemann , J. Kieseler , M. Klute , R. Koppenhöfer , J.M. Lawhorn , M. Link, A. Lintuluoto , B. Maier , S. Maier , S. Mitra , M. Mormile , Th. Müller , M. Neukum, M. Oh , E. Pfeffer , M. Presilla , G. Quast , K. Rabbertz , B. Regnery , N. Shadskiy , I. Shvetsov , H.J. Simonis , L. Sowa, L. Stockmeier, K. Tauqeer, M. Toms , N. Trevisani , R.F. Von Cube , M. Wassmer , S. Wieland , F. Wittig, R. Wolf , X. Zuo 

Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece

G. Anagnostou, G. Daskalakis , A. Kyriakis, A. Papadopoulos³⁰, A. Stakia 

National and Kapodistrian University of Athens, Athens, Greece

P. Kontaxakis , G. Melachroinos, Z. Painesis , I. Papavergou , I. Paraskevas , N. Saoulidou , K. Theofilatos , E. Tziaferi , K. Vellidis , I. Zisopoulos 

National Technical University of Athens, Athens, Greece

G. Bakas , T. Chatzistavrou, G. Karapostoli , K. Kousouris , I. Papakrivopoulos , E. Siamarkou, G. Tsipolitis , A. Zacharopoulou

University of Ioánnina, Ioánnina, Greece

K. Adamidis, I. Bestintzanos, I. Evangelou , C. Foudas, C. Kamtsikis, P. Katsoulis, P. Kokkas , P.G. Kosmoglou Kioseoglou , N. Manthos , I. Papadopoulos , J. Strologas 

HUN-REN Wigner Research Centre for Physics, Budapest, Hungary

C. Hajdu , D. Horvath ^{31,32}, K. Márton, A.J. Rádl ³³, F. Sikler , V. Veszpremi 

MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

M. Csanád , K. Farkas , A. Fehérkúti ³⁴, M.M.A. Gadallah ³⁵, Á. Kadlecik , P. Major , G. Pásztor , G.I. Veres 

Faculty of Informatics, University of Debrecen, Debrecen, Hungary

B. Ujvari , G. Zilizi 

Institute of Nuclear Research ATOMKI, Debrecen, Hungary

G. Bencze, S. Czellar, J. Molnar, Z. Szillasi

Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary

F. Nemes ³⁴, T. Novak 

Panjab University, Chandigarh, India

S. Bansal , S.B. Beri, V. Bhatnagar , G. Chaudhary , S. Chauhan , N. Dhingra ³⁶, A. Kaur , A. Kaur , H. Kaur , M. Kaur , S. Kumar , K. Sandeep , T. Sheokand, J.B. Singh , A. Singla 

University of Delhi, Delhi, India

A. Ahmed , A. Bhardwaj , A. Chhetri , B.C. Choudhary , A. Kumar , A. Kumar , M. Naimuddin , K. Ranjan , M.K. Saini, S. Saumya 

Saha Institute of Nuclear Physics, HBNI, Kolkata, India

S. Baradia , S. Barman ³⁷, S. Bhattacharya , S. Das Gupta, S. Dutta , S. Dutta, S. Sarkar

Indian Institute of Technology Madras, Madras, India

M.M. Ameen , P.K. Behera , S.C. Behera , S. Chatterjee , G. Dash , P. Jana , P. Kalbhor ,
S. Kamble , J.R. Komaragiri ³⁸, D. Kumar ³⁸, P.R. Pujahari , N.R. Saha , A. Sharma ,
A.K. Sikdar , R.K. Singh, P. Verma, S. Verma , A. Vijay

Tata Institute of Fundamental Research-A, Mumbai, India

S. Dugad, G.B. Mohanty , B. Parida , M. Shelake, P. Suryadevara

Tata Institute of Fundamental Research-B, Mumbai, India

A. Bala , S. Banerjee , R.M. Chatterjee, M. Guchait , Sh. Jain , A. Jaiswal, S. Kumar ,
G. Majumder , K. Mazumdar , S. Parolia , A. Thachayath 

National Institute of Science Education and Research, An OCC of Homi Bhabha National Institute, Bhubaneswar, Odisha, India

S. Bahinipati ³⁹, C. Kar , D. Maity ⁴⁰, P. Mal , T. Mishra ,
V.K. Muraleedharan Nair Bindhu ⁴⁰, K. Naskar ⁴⁰, A. Nayak ⁴⁰, S. Nayak, K. Pal, P. Sadangi,
S.K. Swain , S. Varghese ⁴⁰, D. Vats ⁴⁰

Indian Institute of Science Education and Research (IISER), Pune, India

S. Acharya ⁴¹, A. Alpna , S. Dube , B. Gomber ⁴¹, P. Hazarika , B. Kansal , A. Laha ,
B. Sahu ⁴¹, S. Sharma , K.Y. Vaish 

Isfahan University of Technology, Isfahan, Iran

H. Bakhshiansohi ⁴², A. Jafari ⁴³, M. Zeinali ⁴⁴

Institute for Research in Fundamental Sciences (IPM), Tehran, Iran

S. Bashiri, S. Chenarani ⁴⁵, S.M. Etesami , Y. Hosseini , M. Khakzad , E. Khazaie ⁴⁶,
M. Mohammadi Najafabadi , S. Tizchang ⁴⁷

University College Dublin, Dublin, Ireland

M. Felcini , M. Grunewald 

INFN Sezione di Bari^a, Università di Bari^b, Politecnico di Bari^c, Bari, Italy

M. Abbrescia ^{a,b}, A. Colaleo ^{a,b}, D. Creanza ^{a,c}, B. D'Anzi ^{a,b}, N. De Filippis ^{a,c},
M. De Palma ^{a,b}, W. Elmetenawee ^{a,b,17}, L. Fiore ^a, G. Iaselli ^{a,c}, L. Longo ^a, M. Louka ^{a,b},
G. Maggi ^{a,c}, M. Maggi ^a, I. Margjeka ^a, V. Mastrapasqua ^{a,b}, S. My ^{a,b}, S. Nuzzo ^{a,b},
A. Pellecchia ^{a,b}, A. Pompili ^{a,b}, G. Pugliese ^{a,c}, R. Radogna ^{a,b}, D. Ramos ^a, A. Ranieri ^a,
L. Silvestris ^a, F.M. Simone ^{a,c}, Ü. Sözbilir ^a, A. Stamerra ^{a,b}, D. Troiano ^{a,b}, R. Venditti ^{a,b},
P. Verwilligen ^a, A. Zaza ^{a,b}

INFN Sezione di Bologna^a, Università di Bologna^b, Bologna, Italy

G. Abbiendi ^a, C. Battilana ^{a,b}, D. Bonacorsi ^{a,b}, P. Capiluppi ^{a,b}, A. Castro ^{†,a,b},
F.R. Cavallo ^a, M. Cuffiani ^{a,b}, G.M. Dallavalle ^a, T. Diotallevi ^{a,b}, F. Fabbri ^a,
A. Fanfani ^{a,b}, D. Fasanella ^a, P. Giacomelli ^a, L. Giommi ^{a,b}, C. Grandi ^a, L. Guiducci ^{a,b},
S. Lo Meo ^{a,48}, M. Lorusso ^{a,b}, L. Lunerti ^a, S. Marcellini ^a, G. Masetti ^a, F.L. Navarria ^{a,b},

G. Paggi^{a,b}, A. Perrotta^a, F. Primavera^{a,b}, A.M. Rossi^{a,b}, S. Rossi Tisbeni^{a,b},
T. Rovelli^{a,b}, G.P. Siroli^{a,b}

INFN Sezione di Catania^a, Università di Catania^b, Catania, Italy

S. Costa^{a,b,49}, A. Di Mattia^a, A. Lapertosa^a, R. Potenza^{a,b}, A. Tricomi^{a,b,49}, C. Tuve^{a,b}

INFN Sezione di Firenze^a, Università di Firenze^b, Firenze, Italy

P. Assiouras^a, G. Barbagli^a, G. Bardelli^{a,b}, B. Camaiani^{a,b}, A. Cassese^a, R. Ceccarelli^a,
V. Ciulli^{a,b}, C. Civinini^a, R. D'Alessandro^{a,b}, E. Focardi^{a,b}, T. Kello^a, G. Latino^{a,b},
P. Lenzi^{a,b}, M. Lizzo^a, M. Meschini^a, S. Paoletti^a, A. Papanastassiou^{a,b}, G. Sguazzoni^a,
L. Viliani^a

INFN Laboratori Nazionali di Frascati, Frascati, Italy

L. Benussi^a, S. Bianco^a, S. Meola^{a,50}, D. Piccolo^a

INFN Sezione di Genova^a, Università di Genova^b, Genova, Italy

P. Chatagnon^a, F. Ferro^a, E. Robutti^a, S. Tosi^{a,b}

INFN Sezione di Milano-Bicocca^a, Università di Milano-Bicocca^b, Milano, Italy

A. Benaglia^a, F. Brivio^a, F. Cetorelli^{a,b}, F. De Guio^{a,b}, M.E. Dinardo^{a,b}, P. Dini^a,
S. Gennai^a, R. Gerosa^{a,b}, A. Ghezzi^{a,b}, P. Govoni^{a,b}, L. Guzzi^a, M.T. Lucchini^{a,b},
M. Malberti^a, S. Malvezzi^a, A. Massironi^a, D. Menasce^a, L. Moroni^a, M. Paganoni^{a,b},
S. Palluotto^{a,b}, D. Pedrini^a, A. Perego^{a,b}, B.S. Pinolini^a, G. Pizzati^{a,b}, S. Ragazzi^{a,b},
T. Tabarelli de Fatis^{a,b}

INFN Sezione di Napoli^a, Università di Napoli ‘Federico II’^b, Napoli, Italy; Università della Basilicata^c, Potenza, Italy; Scuola Superiore Meridionale (SSM)^d, Napoli, Italy

S. Buontempo^a, A. Cagnotta^{a,b}, F. Carnevali^{a,b}, N. Cavallo^{a,c}, F. Fabozzi^{a,c},
A.O.M. Iorio^{a,b}, L. Lista^{a,b,51}, P. Paolucci^{a,30}, B. Rossi^a

INFN Sezione di Padova^a, Università di Padova^b, Padova, Italy; Università di Trento^c, Trento, Italy

R. Ardino^a, P. Azzi^a, N. Bacchetta^{a,52}, D. Bisello^{a,b}, P. Bortignon^a, G. Bortolato^{a,b},
A. Bragagnolo^{a,b}, A.C.M. Bulla^a, R. Carlin^{a,b}, P. Checchia^a, T. Dorigo^a,
F. Gasparini^{a,b}, U. Gasparini^{a,b}, E. Lusiani^a, M. Margoni^{a,b}, A.T. Meneguzzo^{a,b},
M. Michelotto^a, M. Migliorini^{a,b}, J. Pazzini^{a,b}, P. Ronchese^{a,b}, R. Rossin^{a,b}, M. Tosi^{a,b},
A. Triossi^{a,b}, S. Ventura^a, M. Zanetti^{a,b}, P. Zotto^{a,b}, A. Zucchetta^{a,b}, G. Zumerle^{a,b}

INFN Sezione di Pavia^a, Università di Pavia^b, Pavia, Italy

C. Aimè^a, A. Braghieri^a, S. Calzaferri^a, D. Fiorina^a, P. Montagna^{a,b}, V. Re^a,
C. Riccardi^{a,b}, P. Salvini^a, I. Vai^{a,b}, P. Vitulo^{a,b}

INFN Sezione di Perugia^a, Università di Perugia^b, Perugia, Italy

S. Ajmal^{a,b}, M.E. Ascioti^{a,b}, G.M. Bilei^a, C. Carrivale^{a,b}, D. Ciangottini^{a,b}, L. Fanò^{a,b},
M. Magherini^{a,b}, V. Mariani^{a,b}, M. Menichelli^a, F. Moscatelli^{a,53}, A. Rossi^{a,b},
A. Santocchia^{a,b}, D. Spiga^a, T. Tedeschi^{a,b}

INFN Sezione di Pisa^a, Università di Pisa^b, Scuola Normale Superiore di Pisa^c, Pisa, Italy; Università di Siena^d, Siena, Italy

C.A. Alexe , P. Asenov , P. Azzurri , G. Bagliesi , R. Bhattacharya , L. Bianchini , T. Boccali , E. Bossini , D. Bruschini , R. Castaldi , M.A. Ciocci , M. Cipriani , V. D'Amante , R. Dell'Orso , S. Donato , A. Giassi , F. Ligabue , A.C. Marini , D. Matos Figueiredo , A. Messineo , M. Musich , F. Palla , A. Rizzi , G. Rolandi , S. Roy Chowdhury , T. Sarkar , A. Scribano , P. Spagnolo , R. Tenchini , G. Tonelli , N. Turini , F. Vaselli , A. Venturi , P.G. Verdini 

INFN Sezione di Roma^a, Sapienza Università di Roma^b, Roma, Italy

C. Baldenegro Barrera , P. Barria , C. Basile , F. Cavallari , L. Cunqueiro Mendez , D. Del Re , E. Di Marco , M. Diemoz , F. Errico , E. Longo , J. Mijuskovic , G. Organtini , F. Pandolfi , R. Paramatti , C. Quaranta , S. Rahatlou , C. Rovelli , F. Santanastasio , L. Soffi 

INFN Sezione di Torino^a, Università di Torino^b, Torino, Italy; Università del Piemonte Orientale^c, Novara, Italy

N. Amapane , R. Arcidiacono , S. Argiro , M. Arneodo , N. Bartosik , R. Bellan , A. Bellora , C. Biino , C. Borca , N. Cartiglia , M. Costa , R. Covarelli , N. Demaria , L. Finco , M. Grippio , B. Kiani , F. Legger , F. Luongo , C. Mariotti , L. Markovic , S. Maselli , A. Mecca , L. Menzio , P. Meridiani , E. Migliore , M. Monteno , R. Mulargia , M.M. Obertino , G. Ortona , L. Pacher , N. Pastrone , M. Pelliccioni , M. Ruspa , F. Siviero , V. Sola , A. Solano , A. Staiano , C. Tarricone , D. Trocino , G. Umoret , R. White 

INFN Sezione di Trieste^a, Università di Trieste^b, Trieste, Italy

J. Babbar , S. Belforte , V. Candelise , M. Casarsa , F. Cossutti , K. De Leo , G. Della Ricca 

Kyungpook National University, Daegu, Korea

S. Dogra , J. Hong , B. Kim , J. Kim, D. Lee, H. Lee, S.W. Lee , C.S. Moon , Y.D. Oh , M.S. Ryu , S. Sekmen , B. Tae, Y.C. Yang 

Department of Mathematics and Physics — GWNu, Gangneung, Korea

M.S. Kim 

Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea

G. Bak , P. Gwak , H. Kim , D.H. Moon 

Hanyang University, Seoul, Korea

E. Asilar , J. Choi , D. Kim , T.J. Kim , J.A. Merlin, Y. Ryou

Korea University, Seoul, Korea

S. Choi , S. Han, B. Hong , K. Lee, K.S. Lee , S. Lee , J. Yoo 

Kyung Hee University, Department of Physics, Seoul, KoreaJ. Goh , S. Yang **Sejong University, Seoul, Korea**H. S. Kim , Y. Kim, S. Lee**Seoul National University, Seoul, Korea**J. Almond, J.H. Bhyun, J. Choi , J. Choi, W. Jun , J. Kim , S. Ko , H. Kwon , H. Lee ,
J. Lee , J. Lee , B.H. Oh , S.B. Oh , H. Seo , U.K. Yang, I. Yoon **University of Seoul, Seoul, Korea**W. Jang , D.Y. Kang, Y. Kang , S. Kim , B. Ko, J.S.H. Lee , Y. Lee , I.C. Park , Y. Roh,
I.J. Watson **Yonsei University, Department of Physics, Seoul, Korea**S. Ha , H.D. Yoo **Sungkyunkwan University, Suwon, Korea**M. Choi , M.R. Kim , H. Lee, Y. Lee , I. Yu **College of Engineering and Technology, American University of the Middle East (AUM), Dasman, Kuwait**

T. Beyrouthy, Y. Gharbia

Kuwait University — College of Science — Department of Physics, Safat, KuwaitF. Alazemi **Riga Technical University, Riga, Latvia**K. Dreimanis , A. Gaile , C. Munoz Diaz, D. Osite , G. Pikurs, A. Potrebko , M. Seidel ,
D. Sidiropoulos Kontos**University of Latvia (LU), Riga, Latvia**N.R. Strautnieks **Vilnius University, Vilnius, Lithuania**M. Ambrozas , A. Juodagalvis , A. Rinkevicius , G. Tamulaitis **National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia**I. Yusuff ⁵⁴, Z. Zolkapli**Universidad de Sonora (UNISON), Hermosillo, Mexico**J.F. Benitez , A. Castaneda Hernandez , H.A. Encinas Acosta, L.G. Gallegos Maríñez,
M. León Cello , J.A. Murillo Quijada , A. Sehrawat , L. Valencia Palomo 

Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico

G. Ayala , H. Castilla-Valdez , H. Crotte Ledesma, E. De La Cruz-Burelo ,
I. Heredia-De La Cruz ⁵⁵, R. Lopez-Fernandez , J. Mejia Guisao , C.A. Mondragon Herrera,
A. Sánchez Hernández 

Universidad Iberoamericana, Mexico City, Mexico

C. Oropeza Barrera , D.L. Ramirez Guadarrama, M. Ramírez García 

Benemerita Universidad Autonoma de Puebla, Puebla, Mexico

I. Bautista , I. Pedraza , H.A. Salazar Ibarguen , C. Uribe Estrada 

University of Montenegro, Podgorica, Montenegro

I. Bubanja , N. Raicevic 

University of Canterbury, Christchurch, New Zealand

P.H. Butler 

National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan

A. Ahmad , M.I. Asghar, A. Awais , M.I.M. Awan, H.R. Hoorani , W.A. Khan 

AGH University of Krakow, Faculty of Computer Science, Electronics and Telecommunications, Krakow, Poland

V. Avati, L. Grzanka , M. Malawski 

National Centre for Nuclear Research, Swierk, Poland

H. Bialkowska , M. Bluj , M. Górski , M. Kazana , M. Szleper , P. Zalewski 

Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland

K. Bunkowski , K. Doroba , A. Kalinowski , M. Konecki , J. Krolikowski , A. Muhammad 

Warsaw University of Technology, Warsaw, Poland

K. Pozniak , W. Zabolotny 

Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal

M. Araujo , D. Bastos , C. Beirão Da Cruz E Silva , A. Boletti , M. Bozzo , T. Camporesi ,
G. Da Molin , P. Faccioli , M. Gallinaro , J. Hollar , N. Leonardo , G.B. Marozzo,
T. Niknejad , A. Petrilli , M. Pisano , J. Seixas , J. Varela , J.W. Wulff

Faculty of Physics, University of Belgrade, Belgrade, Serbia

P. Adzic , P. Milenovic 

VINCA Institute of Nuclear Sciences, University of Belgrade, Belgrade, Serbia

D. Devetak, M. Dordevic , J. Milosevic , V. Rekovic

Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain

J. Alcaraz Maestre , Cristina F. Bedoya , Oliver M. Carretero , M. Cepeda , M. Cerrada , N. Colino , B. De La Cruz , A. Delgado Peris , A. Escalante Del Valle , D. Fernández Del Val , J.P. Fernández Ramos , J. Flix , M.C. Fouz , O. Gonzalez Lopez , S. Goy Lopez , J.M. Hernandez , M.I. Josa , E. Martin Viscasillas , D. Moran , C. M. Morcillo Perez , Á. Navarro Tobar , C. Perez Dengra , A. Pérez-Calero Yzquierdo , J. Puerta Pelayo , I. Redondo , S. Sánchez Navas , J. Sastre , J. Vazquez Escobar 

Universidad Autónoma de Madrid, Madrid, Spain

J.F. de Trocóniz 

Universidad de Oviedo, Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA), Oviedo, Spain

B. Alvarez Gonzalez , J. Cuevas , J. Fernandez Menendez , S. Folgueras , I. Gonzalez Caballero , J.R. González Fernández , P. Leguina , E. Palencia Cortezon , J. Prado Pico, C. Ramón Álvarez , V. Rodríguez Bouza , A. Soto Rodríguez , A. Trapote , C. Vico Villalba , P. Vischia 

Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain

S. Bhowmik , S. Blanco Fernández , J.A. Brochero Cifuentes , I.J. Cabrillo , A. Calderon , J. Duarte Campderros , M. Fernandez , G. Gomez , C. Lasosa García , R. Lopez Ruiz , C. Martinez Rivero , P. Martinez Ruiz del Arbol , F. Matorras , P. Matorras Cuevas , E. Navarrete Ramos , J. Piedra Gomez , L. Scodellaro , I. Vila , J.M. Vizan Garcia 

University of Colombo, Colombo, Sri Lanka

B. Kailasapathy ⁵⁶, D.D.C. Wickramaratna 

University of Ruhuna, Department of Physics, Matara, Sri Lanka

W.G.D. Dharmaratna ⁵⁷, K. Liyanage , N. Perera 

CERN, European Organization for Nuclear Research, Geneva, Switzerland

D. Abbaneo , C. Amendola , E. Auffray , G. Auzinger , J. Baechler, D. Barney , A. Bermúdez Martínez , M. Bianco , A.A. Bin Anuar , A. Bocci , L. Borgonovi , C. Botta , E. Brondolin , C. Caillol , G. Cerminara , N. Chernyavskaya , D. d’Enterria , A. Dabrowski , A. David , A. De Roeck , M.M. Defranchis , M. Deile , M. Dobson , G. Franzoni , W. Funk , S. Giani, D. Gigi, K. Gill , F. Glege , J. Hegeman , J.K. Heikkilä , B. Huber, V. Innocente , T. James , P. Janot , O. Kaluzinska , O. Karacheban ²⁸, S. Laurila , P. Lecoq , E. Leutgeb , C. Lourenço , L. Malgeri , M. Mannelli , M. Matthewman, A. Mehta , F. Meijers , S. Mersi , E. Meschi , V. Milosevic , F. Monti , F. Moortgat , M. Mulders , I. Neutelings , S. Orfanelli, F. Pantaleo , G. Petrucciani , A. Pfeiffer , M. Pierini , H. Qu , D. Rabady , B. Ribeiro Lopes , M. Rovere , H. Sakulin , S. Sanchez Cruz , S. Scarfi , C. Schwick, M. Selvaggi , A. Sharma , K. Shchelina , P. Silva , P. Sphicas ⁵⁸, A.G. Stahl Leiton , A. Steen , S. Summers , D. Treille , P. Tropea 

D. Walter , J. Wanczyk ⁵⁹, J. Wang, K.A. Wozniak ⁶⁰, S. Wuchterl , P. Zehetner , P. Zejdl ,
W.D. Zeuner

Paul Scherrer Institut, Villigen, Switzerland

T. Bevilacqua ⁶¹, L. Caminada ⁶¹, A. Ebrahimi , W. Erdmann , R. Horisberger , Q. Ingram ,
H.C. Kaestli , D. Kotlinski , C. Lange , M. Missiroli ⁶¹, L. Noehte ⁶¹, T. Rohe 

ETH Zurich — Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland

T.K. Aarrestad , K. Androsov ⁵⁹, M. Backhaus , G. Bonomelli, A. Calandri , C. Cazzaniga ,
K. Datta , P. De Bryas Dexmiers D'archiac ⁵⁹, A. De Cosa , G. Dissertori , M. Dittmar,
M. Donegà , F. Eble , M. Galli , K. Gedia , F. Glessgen , C. Grab , N. Härringer ,
T.G. Harte, D. Hits , W. Lustermann , A.-M. Lyon , R.A. Manzoni , M. Marchegiani ,
L. Marchese , C. Martin Perez , A. Mascellani ⁵⁹, F. Nessi-Tedaldi , F. Pauss , V. Perovic ,
S. Pigazzini , B. Ristic , F. Riti , R. Seidita , J. Steggemann ⁵⁹, A. Tarabini , D. Valsecchi ,
R. Wallny 

Universität Zürich, Zurich, Switzerland

C. Amsler ⁶², P. Bärtshi , M.F. Canelli , K. Cormier , M. Huwiler , W. Jin , A. Jofrehei ,
B. Kilminster , S. Leontsinis , S.P. Liechti , A. Macchiolo , P. Meiring , F. Meng ,
U. Molinatti , J. Motta , A. Reimers , P. Robmann, M. Senger , E. Shokr, F. Stäger ,
R. Tramontano 

National Central University, Chung-Li, Taiwan

C. Adloff ⁶³, D. Bhowmik, C.M. Kuo, W. Lin, P.K. Rout , P.C. Tiwari ³⁸, S.S. Yu 

National Taiwan University (NTU), Taipei, Taiwan

L. Ceard, K.F. Chen , P.s. Chen, Z.g. Chen, A. De Iorio , W.-S. Hou , T.h. Hsu, Y.w. Kao,
S. Karmakar , R. Khurana, G. Kole , Y.y. Li , R.-S. Lu , E. Paganis , X.f. Su ,
J. Thomas-Wilsker , L.s. Tsai, D. Tsionou, H.y. Wu, E. Yazgan 

High Energy Physics Research Unit, Department of Physics, Faculty of Science, Chulalongkorn University, Bangkok, Thailand

C. Asawatangtrakuldee , N. Srimanobhas , V. Wachirapusanand 

Çukurova University, Physics Department, Science and Art Faculty, Adana, Turkey

D. Agyel , F. Boran , F. Dolek , I. Dumanoglu ⁶⁴, E. Eskut , Y. Guler ⁶⁵,
E. Gurpinar Guler ⁶⁵, C. Isik , O. Kara, A. Kayis Topaksu , U. Kiminsu , G. Onengut ,
K. Ozdemir ⁶⁶, A. Polatoz , B. Tali ⁶⁷, U.G. Tok , S. Turkcapar , E. Uslan , I.S. Zorbakir 

Middle East Technical University, Physics Department, Ankara, Turkey

G. Sokmen, M. Yalvac ⁶⁸

Bogazici University, Istanbul, Turkey

B. Akgun , I.O. Atakisi , E. Gülmez , M. Kaya ⁶⁹, O. Kaya ⁷⁰, S. Tekten ⁷¹

Istanbul Technical University, Istanbul, Turkey

A. Cakir , K. Cankocak ^{64,72}, G.G. Dincer ⁶⁴, Y. Komurcu , S. Sen ⁷³

Istanbul University, Istanbul, Turkey

O. Aydılek ⁷⁴, B. Hacisahinoglu , I. Hos ⁷⁵, B. Kaynak , S. Ozkorucuklu , O. Potok ,
H. Sert , C. Simsek , C. Zorbilmez 

Yildiz Technical University, Istanbul, Turkey

S. Cerci ⁶⁷, B. Isildak ⁷⁶, D. Sunar Cerci , T. Yetkin 

Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkiv, Ukraine

A. Boyaryntsev , B. Grynyov 

National Science Centre, Kharkiv Institute of Physics and Technology, Kharkiv, Ukraine

L. Levchuk 

University of Bristol, Bristol, United Kingdom

D. Anthony , J.J. Brooke , A. Bundock , F. Bury , E. Clement , D. Cussans , H. Flacher ,
M. Glowacki, J. Goldstein , H.F. Heath , M.-L. Holmberg , L. Kreczko , S. Paramesvaran ,
L. Robertshaw, S. Seif El Nasr-Storey, V.J. Smith , N. Stylianou ⁷⁷, K. Walkingshaw Pass

Rutherford Appleton Laboratory, Didcot, United Kingdom

A.H. Ball, K.W. Bell , A. Belyaev ⁷⁸, C. Brew , R.M. Brown , D.J.A. Cockerill , C. Cooke ,
A. Elliot , K.V. Ellis, K. Harder , S. Harper , J. Linacre , K. Manolopoulos, D.M. Newbold ,
E. Olaiya, D. Petyt , T. Reis , A.R. Sahasransu , G. Salvi , T. Schuh,
C.H. Shepherd-Themistocleous , I.R. Tomalin , K.C. Whalen , T. Williams 

Imperial College, London, United Kingdom

I. Andreou , R. Bainbridge , P. Bloch , C.E. Brown , O. Buchmuller, V. Cacchio,
C.A. Carrillo Montoya , G.S. Chahal ⁷⁹, D. Colling , J.S. Dancu, I. Das , P. Dauncey ,
G. Davies , J. Davies, M. Della Negra , S. Fayer, G. Fedi , G. Hall , M.H. Hassanshahi ,
A. Howard, G. Iles , C.R. Knight , J. Langford , J. León Holgado , L. Lyons ,
A.-M. Magnan , S. Mallios, M. Mieskolainen , J. Nash ⁸⁰, M. Pesaresi , P.B. Pradeep,
B.C. Radburn-Smith , A. Richards, A. Rose , K. Savva , C. Seez , R. Shukla , A. Tapper ,
K. Uchida , G.P. Uttley , L.H. Vage, T. Virdee ³⁰, M. Vojinovic , N. Wardle ,
D. Winterbottom 

Brunel University, Uxbridge, United Kingdom

K. Coldham, J.E. Cole , A. Khan, P. Kyberd , I.D. Reid 

Baylor University, Waco, Texas, U.S.A.

S. Abdullin , A. Brinkerhoff , E. Collins , M.R. Darwish ⁸¹, J. Dittmann , K. Hatakeyama ,
J. Hiltbrand , B. McMaster , J. Samudio , S. Sawant , C. Sutantawibul , J. Wilson 

Catholic University of America, Washington, DC, U.S.A.

R. Bartek^{[ID](#)}, A. Dominguez^{[ID](#)}, C. Huerta Escamilla, A.E. Simsek^{[ID](#)}, R. Uniyal^{[ID](#)},
A.M. Vargas Hernandez^{[ID](#)}

The University of Alabama, Tuscaloosa, Alabama, U.S.A.

B. Bam^{[ID](#)}, A. Buchot Perraguin^{[ID](#)}, R. Chudasama^{[ID](#)}, S.I. Cooper^{[ID](#)}, C. Crovella^{[ID](#)}, S.V. Gleyzer^{[ID](#)},
E. Pearson, C.U. Perez^{[ID](#)}, P. Rumerio^{[ID](#)}⁸², E. Usai^{[ID](#)}, R. Yi^{[ID](#)}

Boston University, Boston, Massachusetts, U.S.A.

A. Akpinar^{[ID](#)}, C. Cosby^{[ID](#)}, G. De Castro, Z. Demiragli^{[ID](#)}, C. Erice^{[ID](#)}, C. Fangmeier^{[ID](#)},
C. Fernandez Madrazo^{[ID](#)}, E. Fontanesi^{[ID](#)}, D. Gastler^{[ID](#)}, F. Golf^{[ID](#)}, S. Jeon^{[ID](#)}, J. O'cain, I. Reed^{[ID](#)},
J. Rohlf^{[ID](#)}, K. Salyer^{[ID](#)}, D. Sperka^{[ID](#)}, D. Spitzbart^{[ID](#)}, I. Suarez^{[ID](#)}, A. Tsatsos^{[ID](#)}, A.G. Zecchinelli^{[ID](#)}

Brown University, Providence, Rhode Island, U.S.A.

G. Benelli^{[ID](#)}, D. Cutts^{[ID](#)}, L. Gouskos^{[ID](#)}, M. Hadley^{[ID](#)}, U. Heintz^{[ID](#)}, J.M. Hogan^{[ID](#)}⁸³, T. Kwon^{[ID](#)},
G. Landsberg^{[ID](#)}, K.T. Lau^{[ID](#)}, D. Li^{[ID](#)}, J. Luo^{[ID](#)}, S. Mondal^{[ID](#)}, N. Pervan^{[ID](#)}, T. Russell, S. Sagir^{[ID](#)}⁸⁴,
X. Shen, F. Simpson^{[ID](#)}, M. Stamenkovic^{[ID](#)}, N. Venkatasubramanian, X. Yan^{[ID](#)}

University of California, Davis, Davis, California, U.S.A.

S. Abbott^{[ID](#)}, C. Brainerd^{[ID](#)}, R. Breedon^{[ID](#)}, H. Cai^{[ID](#)}, M. Calderon De La Barca Sanchez^{[ID](#)},
M. Chertok^{[ID](#)}, M. Citron^{[ID](#)}, J. Conway^{[ID](#)}, P.T. Cox^{[ID](#)}, R. Erbacher^{[ID](#)}, F. Jensen^{[ID](#)}, O. Kukral^{[ID](#)},
G. Mocellin^{[ID](#)}, M. Mulhearn^{[ID](#)}, S. Ostrom^{[ID](#)}, W. Wei^{[ID](#)}, S. Yoo^{[ID](#)}, F. Zhang^{[ID](#)}

University of California, Los Angeles, California, U.S.A.

M. Bachtis^{[ID](#)}, R. Cousins^{[ID](#)}, A. Datta^{[ID](#)}, G. Flores Avila^{[ID](#)}, J. Hauser^{[ID](#)}, M. Ignatenko^{[ID](#)},
M.A. Iqbal^{[ID](#)}, T. Lam^{[ID](#)}, E. Manca^{[ID](#)}, A. Nunez Del Prado, D. Saltzberg^{[ID](#)}, V. Valuev^{[ID](#)}

University of California, Riverside, Riverside, California, U.S.A.

R. Clare^{[ID](#)}, J.W. Gary^{[ID](#)}, M. Gordon, G. Hanson^{[ID](#)}, W. Si^{[ID](#)}

University of California, San Diego, La Jolla, California, U.S.A.

A. Aportela, A. Arora^{[ID](#)}, J.G. Branson^{[ID](#)}, S. Cittolin^{[ID](#)}, S. Cooperstein^{[ID](#)}, D. Diaz^{[ID](#)}, J. Duarte^{[ID](#)},
L. Giannini^{[ID](#)}, Y. Gu, J. Guiang^{[ID](#)}, R. Kansal^{[ID](#)}, V. Krutelyov^{[ID](#)}, R. Lee^{[ID](#)}, J. Letts^{[ID](#)},
M. Masciovecchio^{[ID](#)}, F. Mokhtar^{[ID](#)}, S. Mukherjee^{[ID](#)}, M. Pieri^{[ID](#)}, M. Quinnan^{[ID](#)},
B.V. Sathia Narayanan^{[ID](#)}, V. Sharma^{[ID](#)}, M. Tadel^{[ID](#)}, E. Vourliotis^{[ID](#)}, F. Würthwein^{[ID](#)}, Y. Xiang^{[ID](#)},
A. Yagil^{[ID](#)}

University of California, Santa Barbara — Department of Physics, Santa Barbara, California, U.S.A.

A. Barzdukas^{[ID](#)}, L. Brennan^{[ID](#)}, C. Campagnari^{[ID](#)}, K. Downham^{[ID](#)}, C. Grieco^{[ID](#)}, J. Incandela^{[ID](#)},
J. Kim^{[ID](#)}, A.J. Li^{[ID](#)}, P. Masterson^{[ID](#)}, H. Mei^{[ID](#)}, J. Richman^{[ID](#)}, S.N. Santpur^{[ID](#)}, U. Sarica^{[ID](#)},
R. Schmitz^{[ID](#)}, F. Setti^{[ID](#)}, J. Sheplock^{[ID](#)}, D. Stuart^{[ID](#)}, T.Á. Vámi^{[ID](#)}, S. Wang^{[ID](#)}, D. Zhang

California Institute of Technology, Pasadena, California, U.S.A.

S. Bhattacharya^{[ID](#)}, A. Bornheim^{[ID](#)}, O. Cerri, A. Latorre, J. Mao^{[ID](#)}, H.B. Newman^{[ID](#)},
G. Reales Gutiérrez, M. Spiropulu^{[ID](#)}, J.R. Vlimant^{[ID](#)}, C. Wang^{[ID](#)}, S. Xie^{[ID](#)}, R.Y. Zhu^{[ID](#)}

Carnegie Mellon University, Pittsburgh, Pennsylvania, U.S.A.

J. Alison , S. An , P. Bryant , M. Cremonesi , V. Dutta , T. Ferguson ,
T.A. Gómez Espinosa , A. Harilal , A. Kallil Tharayil , C. Liu , T. Mudholkar , S. Murthy ,
P. Palit , K. Park , M. Paulini , A. Roberts , A. Sanchez , W. Terrill 

University of Colorado Boulder, Boulder, Colorado, U.S.A.

J.P. Cumalat , W.T. Ford , A. Hart , A. Hassani , G. Karathanasis , N. Manganelli ,
J. Pearkes , C. Savard , N. Schonbeck , K. Stenson , K.A. Ulmer , S.R. Wagner ,
N. Zipper , D. Zuolo 

Cornell University, Ithaca, New York, U.S.A.

J. Alexander , S. Bright-Thonney , X. Chen , D.J. Cranshaw , J. Fan , X. Fan , S. Hogan ,
P. Kotamnives , J. Monroy , M. Oshiro , J.R. Patterson , M. Reid , A. Ryd , J. Thom ,
P. Wittich , R. Zou 

Fermi National Accelerator Laboratory, Batavia, Illinois, U.S.A.

M. Albrow , M. Alyari , O. Amram , G. Apollinari , A. Apresyan , L.A.T. Bauerick ,
D. Berry , J. Berryhill , P.C. Bhat , K. Burkett , J.N. Butler , A. Canepa , G.B. Cerati ,
H.W.K. Cheung , F. Chlebana , G. Cummings , J. Dickinson , I. Dutta , V.D. Elvira ,
Y. Feng , J. Freeman , A. Gandrakota , Z. Gece , L. Gray , D. Green , A. Grummer ,
S. Grünendahl , D. Guerrero , O. Gutsche , R.M. Harris , R. Heller , T.C. Herwig ,
J. Hirschauer , B. Jayatilaka , S. Jindariani , M. Johnson , U. Joshi , T. Klijnsma ,
B. Klima , K.H.M. Kwok , S. Lammel , D. Lincoln , R. Lipton , T. Liu , C. Madrid ,
K. Maeshima , C. Mantilla , D. Mason , P. McBride , P. Merkel , S. Mrenna , S. Nahn ,
J. Ngadiuba , D. Noonan , S. Norberg , V. Papadimitriou , N. Pastika , K. Pedro ,
C. Pena ⁸⁵, F. Ravera , A. Reinsvold Hall ⁸⁶, L. Ristori , M. Safdari , E. Sexton-Kennedy ,
N. Smith , A. Soha , L. Spiegel , S. Stoynev , J. Strait , L. Taylor , S. Tkaczyk ,
N.V. Tran , L. Uplegger , E.W. Vaandering , I. Zoi 

University of Florida, Gainesville, Florida, U.S.A.

C. Aruta , P. Avery , D. Bourilkov , P. Chang , V. Cherepanov , R.D. Field , C. Huh ,
E. Koenig , M. Kolosova , J. Konigsberg , A. Korytov , K. Matchev , N. Menendez ,
G. Mitselmakher , K. Mohrman , A. Muthirakalayil Madhu , N. Rawal , S. Rosenzweig ,
Y. Takahashi , J. Wang 

Florida State University, Tallahassee, Florida, U.S.A.

T. Adams , A. Al Kadhimi , A. Askew , S. Bower , V. Hagopian , R. Hashmi , R.S. Kim ,
S. Kim , T. Kolberg , G. Martinez , H. Prosper , P.R. Prova , M. Wulansatiti , R. Yohay ,
J. Zhang 

Florida Institute of Technology, Melbourne, Florida, U.S.A.

B. Alsufyani , M.M. Baarmand , S. Butalla , S. Das , T. Elkafrawy ⁸⁷, M. Hohlmann ,
E. Yanes 

University of Illinois Chicago, Chicago, U.S.A., Chicago, U.S.A.

M.R. Adams , A. Baty , C. Bennett, R. Cavanaugh , R. Escobar Franco , O. Evdokimov ,
C.E. Gerber , M. Hawksworth, A. Hingrajiya, D.J. Hofman , J.h. Lee , D. S. Lemos ,
A.H. Merrit , C. Mills , S. Nanda , G. Oh , B. Ozek , D. Pilipovic , R. Pradhan , E. Prifti,
T. Roy , S. Rudrabhatla , M.B. Tonjes , N. Varelas , M.A. Wadud , Z. Ye , J. Yoo 

The University of Iowa, Iowa City, Iowa, U.S.A.

M. Alhousseini , D. Blend, K. Dilsiz ⁸⁸, L. Emediato , G. Karaman , O.K. Köseyan ,
J.-P. Merlo, A. Mestvirishvili ⁸⁹, O. Neogi, H. Ogul ⁹⁰, Y. Onel , A. Penzo , C. Snyder,
E. Tiras ⁹¹

Johns Hopkins University, Baltimore, Maryland, U.S.A.

B. Blumenfeld , L. Corcodilos , J. Davis , A.V. Gritsan , L. Kang , S. Kyriacou ,
P. Maksimovic , M. Roguljic , J. Roskes , S. Sekhar , M. Swartz 

The University of Kansas, Lawrence, Kansas, U.S.A.

A. Abreu , L.F. Alcerro Alcerro , J. Anguiano , S. Arteaga Escatel , P. Baringer , A. Bean ,
Z. Flowers , D. Grove , J. King , G. Krintiras , M. Lazarovits , C. Le Mahieu ,
J. Marquez , M. Murray , M. Nickel , M. Pitt , S. Popescu ⁹², C. Rogan , C. Royon ,
R. Salvatico , S. Sanders , C. Smith , G. Wilson 

Kansas State University, Manhattan, Kansas, U.S.A.

B. Allmond , R. Gujju Gurunadha , A. Ivanov , K. Kaadze , Y. Maravin , J. Natoli ,
D. Roy , G. Sorrentino 

University of Maryland, College Park, Maryland, U.S.A.

A. Baden , A. Belloni , J. Bistany-riebman, Y.M. Chen , S.C. Eno , N.J. Hadley , S. Jabeen ,
R.G. Kellogg , T. Koeth , B. Kronheim, Y. Lai , S. Lascio , A.C. Mignerey , S. Nabili ,
C. Palmer , C. Papageorgakis , M.M. Paranjpe, E. Popova ⁹³, A. Shevelev , L. Wang 

Massachusetts Institute of Technology, Cambridge, Massachusetts, U.S.A.

J. Bendavid , I.A. Cali , P.c. Chou , M. D'Alfonso , J. Eysermans , C. Freer ,
G. Gomez-Ceballos , M. Goncharov, G. Grosso, P. Harris, D. Hoang, D. Kovalskyi , J. Krupa ,
L. Lavezzo , Y.-J. Lee , K. Long , C. McGinn, A. Novak , M.I. Park , C. Paus , C. Reissel ,
C. Roland , G. Roland , S. Rothman , G.S.F. Stephans , Z. Wang , B. Wyslouch ,
T. J. Yang 

University of Minnesota, Minneapolis, Minnesota, U.S.A.

B. Crossman , B.M. Joshi , C. Kapsiak , M. Krohn , D. Mahon , J. Mans , B. Marzocchi ,
M. Revering , R. Rusack , R. Saradhy , N. Strobbe 

University of Nebraska-Lincoln, Lincoln, Nebraska, U.S.A.

K. Bloom , D.R. Claes , G. Haza , J. Hossain , C. Joo , I. Kravchenko , J.E. Siado ,
W. Tabb , A. Vagnerini , A. Wightman , F. Yan , D. Yu 

State University of New York at Buffalo, Buffalo, New York, U.S.A.

H. Bandyopadhyay , L. Hay , H.w. Hsia, I. Iashvili , A. Kalogeropoulos , A. Kharchilava ,
M. Morris , D. Nguyen , J. Pekkanen , S. Rappoccio , H. Rejeb Sfar, A. Williams , P. Young 

Northeastern University, Boston, Massachusetts, U.S.A.

G. Alverson , E. Barberis , J. Bonilla , M. Campana , J. Dervan, Y. Haddad , Y. Han ,
A. Krishna , J. Li , M. Lu , G. Madigan , R. McCarthy , D.M. Morse , V. Nguyen ,
T. Orimoto , A. Parker , L. Skinnari , D. Wood 

Northwestern University, Evanston, Illinois, U.S.A.

J. Bueghly, S. Dittmer , K.A. Hahn , Y. Liu , Y. Miao , D.G. Monk , M.H. Schmitt ,
A. Taliencio , M. Velasco

University of Notre Dame, Notre Dame, Indiana, U.S.A.

G. Agarwal , R. Band , R. Bucci, S. Castells , A. Das , R. Goldouzian , M. Hildreth ,
K.W. Ho , K. Hurtado Anampa , T. Ivanov , C. Jessop , K. Lannon , J. Lawrence ,
N. Loukas , L. Lutton , J. Mariano, N. Marinelli, I. Mcalister, T. McCauley , C. Mcgrady ,
C. Moore , Y. Musienko ¹⁶, H. Nelson , M. Osherson , A. Piccinelli , R. Ruchti ,
A. Townsend , Y. Wan, M. Wayne , H. Yockey, M. Zarucki , L. Zygala 

The Ohio State University, Columbus, Ohio, U.S.A.

A. Basnet , B. Bylsma, M. Carrigan , L.S. Durkin , C. Hill , M. Joyce , M. Nunez Ornelas ,
K. Wei, B.L. Winer , B. R. Yates 

Princeton University, Princeton, New Jersey, U.S.A.

H. Bouchamaoui , P. Das , G. Dezoort , P. Elmer , A. Frankenthal , B. Greenberg ,
N. Haubrich , K. Kennedy, G. Kopp , S. Kwan , D. Lange , A. Loeliger , D. Marlow ,
I. Ojalvo , J. Olsen , D. Stickland , C. Tully 

University of Puerto Rico, Mayaguez, Puerto Rico, U.S.A.

S. Malik 

Purdue University, West Lafayette, Indiana, U.S.A.

A.S. Bakshi , S. Chandra , R. Chawla , A. Gu , L. Gutay, M. Jones , A.W. Jung ,
A.M. Koshy, M. Liu , G. Negro , N. Neumeister , G. Paspalaki , S. Piperov , V. Scheurer,
J.F. Schulte , M. Stojanovic , J. Thieman , A. K. Virdi , F. Wang , A. Wildridge , W. Xie ,
Y. Yao 

Purdue University Northwest, Hammond, Indiana, U.S.A.

J. Dolen , N. Parashar , A. Pathak 

Rice University, Houston, Texas, U.S.A.

D. Acosta , T. Carnahan , K.M. Ecklund , P.J. Fernández Manteca , S. Freed, P. Gardner,
F.J.M. Geurts , I. Krommydas , W. Li , J. Lin , O. Miguel Colin , B.P. Padley , R. Redjimi,
J. Rotter , E. Yigitbasi , Y. Zhang 

University of Rochester, Rochester, New York, U.S.A.

A. Bodek , P. de Barbaro , R. Demina , J.L. Dulemba , A. Garcia-Bellido , O. Hindrichs ,
A. Khukhunaishvili , N. Parmar, P. Parygin ⁹³, R. Taus 

Rutgers, The State University of New Jersey, Piscataway, New Jersey, U.S.A.

B. Chiarito, J.P. Chou , S.V. Clark , D. Gadkari , Y. Gershtein , E. Halkiadakis ,
M. Heindl , C. Houghton , D. Jaroslawski , S. Konstantinou , I. Laflotte , A. Lath ,
R. Montalvo, K. Nash, J. Reichert , H. Routray , P. Saha , S. Salur , S. Schnetzer,
S. Somalwar , R. Stone , S.A. Thayil , S. Thomas, J. Vora , H. Wang 

University of Tennessee, Knoxville, Tennessee, U.S.A.

D. Ally , A.G. Delannoy , S. Fiorendi , S. Higginbotham , T. Holmes , A.R. Kanuganti ,
N. Karunarathna , L. Lee , E. Nibigira , S. Spanier 

Texas A&M University, College Station, Texas, U.S.A.

D. Aebi , M. Ahmad , T. Akhter , O. Bouhali ⁹⁴, R. Eusebi , J. Gilmore , T. Huang ,
T. Kamon ⁹⁵, H. Kim , S. Luo , R. Mueller , D. Overton , D. Rathjens , A. Safonov 

Texas Tech University, Lubbock, Texas, U.S.A.

N. Akchurin , J. Damgov , N. Gogate , V. Hegde , A. Hussain , Y. Kazhykarim,
K. Lamichhane , S.W. Lee , A. Mankel , T. Peltola , I. Volobouev 

Vanderbilt University, Nashville, Tennessee, U.S.A.

E. Appelt , Y. Chen , S. Greene, A. Gurrola , W. Johns , R. Kunnawalkam Elayavalli ,
A. Melo , F. Romeo , P. Sheldon , S. Tuo , J. Velkovska , J. Viinikainen 

University of Virginia, Charlottesville, Virginia, U.S.A.

B. Cardwell , H. Chung, B. Cox , J. Hakala , R. Hirosky , A. Ledovsky , C. Neu 

Wayne State University, Detroit, Michigan, U.S.A.

S. Bhattacharya , P.E. Karchin 

University of Wisconsin — Madison, Madison, Wisconsin, U.S.A.

A. Aravind, S. Banerjee , K. Black , T. Bose , S. Dasu , I. De Bruyn , P. Everaerts ,
C. Galloni, H. He , M. Herndon , A. Herve , C.K. Koraka , A. Lanaro, R. Loveless ,
J. Madhusudanan Sreekala , A. Mallampalli , A. Mohammadi , S. Mondal, G. Parida ,
L. Pétré , D. Pinna, A. Savin, V. Shang , V. Sharma , W.H. Smith , D. Teague, H.F. Tsoi ,
W. Vetens , A. Warden 

Authors affiliated with an institute or an international laboratory covered by a cooperation agreement with CERN

S. Afanasiev , V. Alexakhin , D. Budkouski , I. Golutvin [†], I. Gorbunov , V. Karjavine ,
V. Korenkov , A. Lanev , A. Malakhov , V. Matveev ⁹⁶, V. Palichik , V. Perelygin ,
M. Savina , V. Shalaev , S. Shmatov , S. Shulha , V. Smirnov , O. Teryaev , N. Voytishin ,
B.S. Yuldashev⁹⁷, A. Zarubin , I. Zhizhin , G. Gavrilo , V. Golovtcov , Y. Ivanov ,
V. Kim ⁹⁶, P. Levchenko ⁹⁸, V. Murzin , V. Oreshkin , D. Sosnov , V. Sulimov 

L. Uvarov^{}, A. Vorobyev[†], Yu. Andreev^{}, A. Dermenev^{}, S. Gninenko^{}, N. Golubev^{},
A. Karneyeu^{}, D. Kirpichnikov^{}, M. Kirsanov^{}, N. Krasnikov^{}, I. Tlisova^{}, A. Toropin^{},
T. Aushev^{}, V. Gavrilov^{}, N. Lychkovskaya^{}, A. Nikitenko^{}^{99,100}, V. Popov^{}, A. Zhokin^{},
R. Chistov^{}⁹⁶, M. Danilov^{}⁹⁶, S. Polikarpov^{}⁹⁶, V. Andreev^{}, M. Azarkin^{}, M. Kirakosyan,
A. Terkulov^{}, E. Boos^{}, V. Bunichev^{}, M. Dubinin^{}⁸⁵, L. Dudko^{}, A. Ershov^{}, V. Klyukhin^{},
O. Kodolova^{}¹⁰⁰, S. Obraztsov^{}, M. Perfilov, V. Savrin^{}, P. Volkov^{}, G. Vorotnikov^{},
V. Blinov^{}⁹⁶, T. Dimova^{}⁹⁶, A. Kozyrev^{}⁹⁶, O. Radchenko^{}⁹⁶, Y. Skovpen^{}⁹⁶, V. Kachanov^{},
D. Konstantinov^{}, S. Slabospitskii^{}, A. Uzunian^{}, A. Babaev^{}, V. Borshch^{}, D. Druzhkin^{}¹⁰¹

Authors affiliated with an institute formerly covered by a cooperation agreement with CERN

V. Chekhovsky, V. Makarenko^{}

[†] Deceased

¹ Also at Yerevan State University, Yerevan, Armenia

² Also at TU Wien, Vienna, Austria

³ Also at Ghent University, Ghent, Belgium

⁴ Also at Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

⁵ Also at Universidade Estadual de Campinas, Campinas, Brazil

⁶ Also at Federal University of Rio Grande do Sul, Porto Alegre, Brazil

⁷ Also at UFMS, Nova Andradina, Brazil

⁸ Also at Nanjing Normal University, Nanjing, China

⁹ Now at The University of Iowa, Iowa City, Iowa, U.S.A.

¹⁰ Also at University of Chinese Academy of Sciences, Beijing, China

¹¹ Also at China Center of Advanced Science and Technology, Beijing, China

¹² Also at University of Chinese Academy of Sciences, Beijing, China

¹³ Also at China Spallation Neutron Source, Guangdong, China

¹⁴ Now at Henan Normal University, Xinxiang, China

¹⁵ Also at Université Libre de Bruxelles, Bruxelles, Belgium

¹⁶ Also at an institute or an international laboratory covered by a cooperation agreement with CERN

¹⁷ Also at Helwan University, Cairo, Egypt

¹⁸ Now at Zewail City of Science and Technology, Zewail, Egypt

¹⁹ Now at British University in Egypt, Cairo, Egypt

²⁰ Now at Cairo University, Cairo, Egypt

²¹ Also at Purdue University, West Lafayette, Indiana, U.S.A.

²² Also at Université de Haute Alsace, Mulhouse, France

²³ Also at Istinye University, Istanbul, Turkey

²⁴ Also at The University of the State of Amazonas, Manaus, Brazil

²⁵ Also at University of Hamburg, Hamburg, Germany

²⁶ Also at RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

²⁷ Also at Bergische University Wuppertal (BUW), Wuppertal, Germany

²⁸ Also at Brandenburg University of Technology, Cottbus, Germany

²⁹ Also at Forschungszentrum Jülich, Juelich, Germany

³⁰ Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland

³¹ Also at Institute of Nuclear Research ATOMKI, Debrecen, Hungary

³² Now at Universitatea Babes-Bolyai — Facultatea de Fizica, Cluj-Napoca, Romania

³³ Also at MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

³⁴ Also at HUN-REN Wigner Research Centre for Physics, Budapest, Hungary

³⁵ Also at Physics Department, Faculty of Science, Assiut University, Assiut, Egypt

³⁶ Also at Punjab Agricultural University, Ludhiana, India

- ³⁷ Also at *University of Visva-Bharati, Santiniketan, India*
- ³⁸ Also at *Indian Institute of Science (IISc), Bangalore, India*
- ³⁹ Also at *IIT Bhubaneswar, Bhubaneswar, India*
- ⁴⁰ Also at *Institute of Physics, Bhubaneswar, India*
- ⁴¹ Also at *University of Hyderabad, Hyderabad, India*
- ⁴² Also at *Deutsches Elektronen-Synchrotron, Hamburg, Germany*
- ⁴³ Also at *Isfahan University of Technology, Isfahan, Iran*
- ⁴⁴ Also at *Sharif University of Technology, Tehran, Iran*
- ⁴⁵ Also at *Department of Physics, University of Science and Technology of Mazandaran, Behshahr, Iran*
- ⁴⁶ Also at *Department of Physics, Isfahan University of Technology, Isfahan, Iran*
- ⁴⁷ Also at *Department of Physics, Faculty of Science, Arak University, ARAK, Iran*
- ⁴⁸ Also at *Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Bologna, Italy*
- ⁴⁹ Also at *Centro Siciliano di Fisica Nucleare e di Struttura Della Materia, Catania, Italy*
- ⁵⁰ Also at *Università degli Studi Guglielmo Marconi, Roma, Italy*
- ⁵¹ Also at *Scuola Superiore Meridionale, Università di Napoli ‘Federico II’, Napoli, Italy*
- ⁵² Also at *Fermi National Accelerator Laboratory, Batavia, Illinois, U.S.A.*
- ⁵³ Also at *Consiglio Nazionale delle Ricerche — Istituto Officina dei Materiali, Perugia, Italy*
- ⁵⁴ Also at *Department of Applied Physics, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Malaysia*
- ⁵⁵ Also at *Consejo Nacional de Ciencia y Tecnología, Mexico City, Mexico*
- ⁵⁶ Also at *Trincomalee Campus, Eastern University, Sri Lanka, Nilaveli, Sri Lanka*
- ⁵⁷ Also at *Saegis Campus, Nugegoda, Sri Lanka*
- ⁵⁸ Also at *National and Kapodistrian University of Athens, Athens, Greece*
- ⁵⁹ Also at *Ecole Polytechnique Fédérale Lausanne, Lausanne, Switzerland*
- ⁶⁰ Also at *University of Vienna, Vienna, Austria*
- ⁶¹ Also at *Universität Zürich, Zurich, Switzerland*
- ⁶² Also at *Stefan Meyer Institute for Subatomic Physics, Vienna, Austria*
- ⁶³ Also at *Laboratoire d’Annecy-le-Vieux de Physique des Particules, IN2P3-CNRS, Annecy-le-Vieux, France*
- ⁶⁴ Also at *Near East University, Research Center of Experimental Health Science, Mersin, Turkey*
- ⁶⁵ Also at *Konya Technical University, Konya, Turkey*
- ⁶⁶ Also at *Izmir Bakircay University, Izmir, Turkey*
- ⁶⁷ Also at *Adiyaman University, Adiyaman, Turkey*
- ⁶⁸ Also at *Bozok Universitetsi Rektörlüğü, Yozgat, Turkey*
- ⁶⁹ Also at *Marmara University, Istanbul, Turkey*
- ⁷⁰ Also at *Milli Savunma University, Istanbul, Turkey*
- ⁷¹ Also at *Kafkas University, Kars, Turkey*
- ⁷² Now at *Istanbul Okan University, Istanbul, Turkey*
- ⁷³ Also at *Hacettepe University, Ankara, Turkey*
- ⁷⁴ Also at *Erzincan Binali Yildirim University, Erzincan, Turkey*
- ⁷⁵ Also at *Istanbul University — Cerrahpasa, Faculty of Engineering, Istanbul, Turkey*
- ⁷⁶ Also at *Yildiz Technical University, Istanbul, Turkey*
- ⁷⁷ Also at *Vrije Universiteit Brussel, Brussel, Belgium*
- ⁷⁸ Also at *School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom*
- ⁷⁹ Also at *IPPP Durham University, Durham, United Kingdom*
- ⁸⁰ Also at *Monash University, Faculty of Science, Clayton, Australia*
- ⁸¹ Also at *Institute of Basic and Applied Sciences, Faculty of Engineering, Arab Academy for Science, Technology and Maritime Transport, Alexandria, Egypt*
- ⁸² Also at *Università di Torino, Torino, Italy*
- ⁸³ Also at *Bethel University, St. Paul, Minnesota, U.S.A.*
- ⁸⁴ Also at *Karamanoğlu Mehmetbey University, Karaman, Turkey*
- ⁸⁵ Also at *California Institute of Technology, Pasadena, California, U.S.A.*
- ⁸⁶ Also at *United States Naval Academy, Annapolis, Maryland, U.S.A.*

- ⁸⁷ *Also at Ain Shams University, Cairo, Egypt*
⁸⁸ *Also at Bingol University, Bingol, Turkey*
⁸⁹ *Also at Georgian Technical University, Tbilisi, Georgia*
⁹⁰ *Also at Sinop University, Sinop, Turkey*
⁹¹ *Also at Erciyes University, Kayseri, Turkey*
⁹² *Also at Horia Hulubei National Institute of Physics and Nuclear Engineering (IFIN-HH), Bucharest, Romania*
⁹³ *Now at another institute or international laboratory covered by a cooperation agreement with CERN*
⁹⁴ *Also at Texas A&M University at Qatar, Doha, Qatar*
⁹⁵ *Also at Kyungpook National University, Daegu, Korea*
⁹⁶ *Also at another institute or international laboratory covered by a cooperation agreement with CERN*
⁹⁷ *Also at Institute of Nuclear Physics of the Uzbekistan Academy of Sciences, Tashkent, Uzbekistan*
⁹⁸ *Also at Northeastern University, Boston, Massachusetts, U.S.A.*
⁹⁹ *Also at Imperial College, London, United Kingdom*
¹⁰⁰ *Now at Yerevan Physics Institute, Yerevan, Armenia*
¹⁰¹ *Also at Universiteit Antwerpen, Antwerpen, Belgium*