

## The geometric measure of entanglement for a symmetric pure state with non-negative amplitudes

Masahito Hayashi, Damian Markham, Mio Murao, Masaki Owari, and Shashank Virmani

Citation: *Journal of Mathematical Physics* **50**, 122104 (2009); doi: 10.1063/1.3271041

View online: <http://dx.doi.org/10.1063/1.3271041>

View Table of Contents: <http://scitation.aip.org/content/aip/journal/jmp/50/12?ver=pdfcov>

Published by the [AIP Publishing](#)

---

### Articles you may be interested in

[Geometric descriptions of entangled states by auxiliary varieties](#)

*J. Math. Phys.* **53**, 102203 (2012); 10.1063/1.4753989

[Non Zero Moments of Some Entangled Three Qubit Symmetric States](#)

*AIP Conf. Proc.* **1384**, 125 (2011); 10.1063/1.3635854

[Two-sided bounds on minimum-error quantum measurement, on the reversibility of quantum dynamics, and on maximum overlap using directional iterates](#)

*J. Math. Phys.* **51**, 092204 (2010); 10.1063/1.3463451

[Asymptotic performance of optimal state estimation in qubit system](#)

*J. Math. Phys.* **49**, 102101 (2008); 10.1063/1.2988130

[Minimally disturbing Heisenberg–Weyl symmetric measurements using hard-core collisions of Schrödinger particles](#)

*J. Math. Phys.* **47**, 082102 (2006); 10.1063/1.2222080

---

The logo for the AIP Journal of Applied Physics. It features the letters 'AIP' in a large, white, sans-serif font on the left. To its right is a vertical orange bar, followed by the text 'Journal of Applied Physics' in a smaller, white, sans-serif font. The background is a solid orange color.

*Journal of Applied Physics* is pleased to announce **André Anders** as its new Editor-in-Chief

## The geometric measure of entanglement for a symmetric pure state with non-negative amplitudes

Masahito Hayashi,<sup>1,a)</sup> Damian Markham,<sup>2</sup> Mio Muroa,<sup>3,4</sup> Masaki Owari,<sup>5</sup> and Shashank Virmani<sup>5,6</sup>

<sup>1</sup>Graduate School of Information Sciences, Tohoku University, Aoba-ku, Sendai 980-8579, Japan

<sup>2</sup>LTCI CNRS, TELECOM ParisTech, 37/39 rue Dareau, 75014 Paris, France

<sup>3</sup>Department of Physics, Graduate School of Science, The University of Tokyo, Tokyo 113-0033, Japan

<sup>4</sup>Institute for Nano Quantum Information Electronics, The University of Tokyo, Tokyo 113-0033, Japan

<sup>5</sup>Optics Section, Blackett Laboratory and Institute for Mathematical Sciences, Imperial College, London SW7 2AZ, United Kingdom

<sup>6</sup>Department Physics, SUPA, University of Strathclyde, Glasgow G4 0NG, United Kingdom

(Received 20 June 2009; accepted 10 November 2009; published online 28 December 2009)

In this paper for a class of symmetric multipart pure states, we consider a conjecture related to the geometric measure of entanglement: “for a symmetric pure state, the closest product state in terms of the fidelity can be chosen as a symmetric product state.” We show that this conjecture is true for symmetric pure states whose amplitudes are all non-negative in a computational basis. The more general conjecture is still open. © 2009 American Institute of Physics. [doi:10.1063/1.3271041]

### I. INTRODUCTION

The geometric measure of entanglement, which was first proposed by Shimony<sup>1</sup> and extended to multipartite systems by Wei *et al.*,<sup>2</sup> is one of the most natural entanglement measures for pure states in multipartite systems and has applications in various different topics, including many body physics,<sup>3–5</sup> local discrimination,<sup>6</sup> quantum computation,<sup>7,8</sup> condensed matter systems,<sup>9,10</sup> entanglement witnesses,<sup>11,12</sup> and the study of quantum channel capacities.<sup>13</sup>

Moreover, the same function (except its normalization) plays important roles in different fields of science apart from physics. First, the geometric measure of entanglement is nothing but the *injective tensor norm* itself, which appears in theory of operator algebra<sup>14</sup> and is now becoming increasingly important in theoretical physics—particularly in quantum information.<sup>15</sup> Second, in signal processing, especially in the field of multiway data analysis, high order statistics and independent component analysis, this function has been intensively studied under the name of *Rank one approximation to high order tensors*.<sup>16–21</sup>

In spite of its importance, its value has only been determined for limited classes of states with large symmetries, such as Greenberger-Horne-Zeilinger (GHZ) states, generalized W states, and certain families of stabilizer states.<sup>12,22,23</sup> This is because the geometric measure of entanglement is defined in terms of the maximum fidelity between the state and a pure product state, and therefore poses a difficult optimization problem.

In quantum information, there are several entanglement measures that attempt to quantify the “distance” between a quantum state and the set of separable states. For example, the relative entropy of entanglement and the robustness of entanglement. For such measures, when a given entangled state is invariant under a certain group action, we can normally choose a closest sepa-

<sup>a)</sup>Electronic mail: hayashi@math.is.tohoku.ac.jp.

erable state as it is invariant under the same group action. This property of measures helps us derive an exact value of these measures for such states with group symmetry.<sup>24</sup>

On the other hand, even though the geometric measure of entanglement is also defined in terms of the distance from a set of product states, it does not possess this property. In other words, for a given state, closest product states in terms of the fidelity do not inherit its group symmetry. For example, it is known that a translationally invariant closest product state for the translationally invariant GHZ-type state  $(|01\cdots 01\rangle + |10\cdots 10\rangle)/\sqrt{2}$  does not exist. Nevertheless, many researchers believed that permutation symmetry is exceptionally inherited by closest product states. In other words, there is one prominent conjecture: *for a symmetric pure state, the maximization can be attained by a symmetric product state*. If this conjecture was true, it could vastly reduce the computation of the geometric measure of entanglement for a symmetric pure state. To our knowledge, this conjecture first appeared in Ref. 2, where it was used in order to propose an analytical formula for the geometric measure of entanglement for GHZ and W states. Subsequently, this conjecture, and a stronger version (in which “symmetric” is replaced by “translationally invariant”), was used in calculations of the geometric measure for states of many body systems.<sup>4,5</sup> In Ref. 12, the authors attempted to prove this conjecture. However, it remained an open problem.

In this paper, we give a proof of this conjecture for a restricted but large class of symmetric states: symmetric states whose amplitudes are all non-negative in a given computational basis. This class involves many famous states such as GHZ states, W states, Dicke states, and also superposition of these states involving only non-negative coefficients. Our result is hence sufficient to give mathematical rigor to the computations of the entanglement of types of symmetric pure state that were presented in Refs. 2, 12, and 22.

## II. DEFINITIONS AND MAIN RESULT

Throughout this paper we will treat only finite dimensional Hilbert spaces obtained from tensor products of a single space  $\mathcal{H}$ . We start with the definition of the geometric measure of entanglement.

*Definition 1:* For a state  $|\Psi\rangle$ , the geometric measure of entanglement is defined as

$$E_g(|\psi\rangle) = \min_{|\Phi\rangle \in \text{Pro}(\mathcal{H}^{\otimes n})} -\log_2(|\langle \Phi | \psi \rangle|^2), \quad (1)$$

where  $\text{Pro}(\mathcal{H}^{\otimes n})$  is the set of product states on  $\mathcal{H}^{\otimes n}$ .

This is the distance between state  $|\psi\rangle$  and the closest product state  $|\Phi\rangle$  in terms of fidelity and has operational significance in several directions.<sup>3-8,11-13</sup> The measure can be extended to the mixed state case in a natural way via the convex roof method.<sup>2</sup> Several properties of this measure have already been studied and we know that it has many of the nice properties one might require from an entanglement measure.<sup>1,2,22,25</sup>

The main result of this paper is the following theorem.

**Theorem 1:** *If there exists a basis  $\{|i\rangle\}_{i=1}^{\dim \mathcal{H}}$  of  $\mathcal{H}$  such that a permutation invariant pure state  $|\Psi\rangle \in \mathcal{S}_n$  satisfies  $\langle \Psi | (|i_1\rangle \otimes \cdots \otimes |i_n\rangle) \geq 0$  for all  $i_1, \dots, i_n$ , then a closest product state  $|\Phi\rangle$  may be found in the symmetric Hilbert space. More precisely,*

$$E_g(|\Psi\rangle) = -\log_2 \max_{|\phi\rangle \in \mathcal{H}} |\langle \phi |^{\otimes n} | \Psi \rangle|^2, \quad (2)$$

where  $\mathcal{S}_n$  is the symmetric subspace of  $\mathcal{H}^{\otimes n}$ . In addition, we may choose an optimal state  $|\phi\rangle$  such that it satisfies  $\langle i | \phi \rangle \geq 0$  for all  $i$ .

As we have already mentioned, the preconditions of this theorem are satisfied by a large class of symmetric state, including GHZ and W states. The theorem hence gives a mathematically rigorous proof for the calculations in Refs. 2, 12, and 22. Intriguingly, an identical result to Theorem 1 has also been independently proven by Wei and Severini using methods from the theory of permanents.<sup>26</sup> It will be of interest to identify whether there are hidden similarities to the proofs, or whether they are truly distinct.

### III. PROOF OF MAIN RESULT

In order to prove Theorem 1, we will need to utilize Perron–Frobenius (PF) theorem (Ref. 27, Theorem 8.3.1) and another lemma, Lemma 1, which we now present and prove. We use the following notation. First, for a vector  $u \in \mathbb{R}^d$  expressed in a certain privileged basis, we say that  $u$  is non-negative if all elements of  $u$  are non-negative on that basis. We denote this using the notation  $u \geq 0$ . We use a similar notation for complex vector spaces: a pure state  $|v\rangle \in \mathcal{H}$  is said to be non-negative if  $\langle i|v\rangle \geq 0$  for a privileged basis  $\{|i\rangle\}_i$ , which we will define shortly. The non-negativity of a pure state will be denoted by  $|v\rangle \geq 0$ .

For a state  $|\Psi\rangle \in \mathcal{H}^{\otimes 2}$ , a product basis  $\{|i_1\rangle \otimes |i_2\rangle\}$  gives a natural isomorphism between all states  $|\Psi\rangle$  on  $\mathcal{H} \otimes \mathcal{H}$  and all  $d \times d$  complex matrices  $\Psi$  satisfying  $\text{Tr } \Psi^\dagger \Psi = 1$ , where  $d := \dim \mathcal{H}$ . If  $\{|i_1\rangle \otimes |i_2\rangle\}$  satisfies  $\langle \Psi | (|i_1\rangle \otimes |i_2\rangle) \in \mathbb{R}$ , then the state  $|\Psi\rangle$  corresponds to a *real matrix*  $\Psi$ . If such a bipartite state is permutationally invariant, then the matrix will also be symmetric. The real symmetricity of  $\Psi$  implies that its largest eigenvalue  $\lambda_1$  is also a singular value so that

$$\lambda_1 = \sup_{\|w\|=1} w^\dagger \Psi w = \sup_{\|w\|=1} \|\Psi w\|.$$

Since  $\Psi$  has non-negative elements, the (extended) PF theorem (Ref. 27, Theorem 8.3.1) implies that the largest singular value equals the largest eigenvalue and the corresponding eigenvector can be chosen to have non-negative elements so that

$$\lambda_1 = \sup\{w^T \Psi w | w_k \geq 0, w = 1\} = \sup\{u^T \Psi v | u_k, v_k \geq 0, \|u\| = \|v\| = 1\}. \quad (3)$$

Here, we give a lemma.

*Lemma 1:* If  $u$  and  $v$  are normalized vectors with non-negative elements and  $\lambda_1 = u^T \Psi v$ , then  $w = (u+v)/\|u+v\|$  is an eigenvector of  $\lambda_1$  with non-negative elements.

*Proof:* By the Cauchy–Schwarz inequality,

$$\lambda_1 = u^T \Psi v \leq \|\Psi v\| \leq \lambda_1.$$

Therefore, equality holds above and by the conditions for equality in Cauchy–Schwarz, we must have  $cu = \Psi v$ . Then, the conditions on  $u$  and  $v$  imply that  $c = \lambda$ . Applying the same argument to  $(\Psi u)^T v$ , we obtain  $\Psi u = \lambda v$ . So, the relation  $\Psi(u+v)/\|u+v\| = \lambda(u+v)/\|u+v\|$  holds. ■

Now, we are ready to prove Theorem 1.

*Proof of Theorem 1:* We start the proof by noting two important facts. First, if the statement of this theorem is valid, then the same statement is valid for a non-normalized state—the definition of the geometric measure of entanglement  $E_g(|\Psi\rangle)$  can be easily extended to a non-normalized state. Second, suppose that the assumption of the theorem is valid. Then because the amplitudes of  $\Psi$  are non-negative, we can easily see that

$$\max_{|\Phi\rangle \in \text{Pro}(\mathcal{H}^{\otimes n})} |\langle \Phi | \Psi \rangle| = \max_{|\Phi\rangle \in \text{Pro}(\mathcal{H}^{\otimes n})} \{ \langle \Phi | \Psi \rangle \langle i_1 \otimes \cdots \otimes |i_n\rangle | \Phi \rangle \geq 0, \forall i_1, \dots, i_n \}, \quad (4)$$

where  $\text{Pro}(\mathcal{H}^{\otimes n})$  is the set of all product states on  $\mathcal{H}^{\otimes n}$ . So, we just need to consider the optimization problem in the right hand side of the above equation. We will prove Theorem 1 by induction with respect to a number  $n$  of tensor copies of the Hilbert space.

For  $n=2$ , by means of the natural correspondence between bipartite states and matrices, we derive

$$\max_{|u\rangle, |v\rangle \in \mathcal{H}} \{ |u\rangle \otimes \langle v| | \Psi \rangle \langle i|u\rangle \geq 0 \text{ and } \langle i|v\rangle \geq 0, \forall i \} = \max_{u, v \in \mathbb{R}^d} \{ u^T \Psi v | u, v \geq 0 \}. \quad (5)$$

Thus, in the case  $n=2$ , Eq. (2) follows directly from (3).

Suppose that for all  $n \leq k$  the statement of this theorem is valid, and  $|\Psi\rangle \in \mathcal{H}^{\otimes k+1}$  satisfies the assumption of the theorem. Then, since  $|\Psi\rangle$  is non-negative, it satisfies Eq. (4). Thus, there exists a non-negative product state  $|a_1\rangle \otimes \cdots \otimes |a_{k+1}\rangle \geq 0$  satisfying

$$\langle a_1 | \otimes \cdots \otimes \langle a_{k+1} | | \Psi \rangle = \max_{|\Phi\rangle \in \text{Pro}(\mathcal{H}^{\otimes k+1})} |\langle \Phi | \Psi \rangle|. \tag{6}$$

We now define a non-normalized state  $|\Psi'_0\rangle \in \mathcal{H}^{\otimes k}$  as  $|\Psi'_0\rangle \stackrel{\text{def}}{=} I_{\mathcal{H}}^{\otimes k} \otimes \langle a_{k+1} | \Psi \rangle$ —clearly this state is also non-negative. Then, this state satisfies

$$\langle a_1 | \otimes \cdots \otimes \langle a_k | | \Psi'_0 \rangle = \max_{|\Phi\rangle \in \text{Pro}(\mathcal{H}^{\otimes k+1})} |\langle \Phi | \Psi \rangle|. \tag{7}$$

Now suppose that there exists a non-negative product state  $|a'_1\rangle \otimes \cdots \otimes |a'_k\rangle \geq 0$  satisfying  $\langle a'_1 | \otimes \cdots \otimes \langle a'_k | | \Psi'_0 \rangle > \langle a_1 | \otimes \cdots \otimes \langle a_k | | \Psi'_0 \rangle$ . Then, a non-negative product state  $|a'_1\rangle \otimes \cdots \otimes |a'_k\rangle \otimes |a_{k+1}\rangle \geq 0$  satisfies  $\langle a'_1 | \otimes \cdots \otimes \langle a'_k | \otimes \langle a_{k+1} | | \Psi \rangle > \max_{|\Phi\rangle \in \text{Pro}(\mathcal{H}^{\otimes k+1})} |\langle \Phi | \Psi \rangle|$ . However, since  $\langle a'_1 | \otimes \cdots \otimes \langle a'_k | \otimes \langle a_{k+1} | \in \text{Pro}(\mathcal{H}^{\otimes k+1})$ , this would be a contradiction. Hence, we obtain

$$\langle a_1 | \otimes \cdots \otimes \langle a_k | | \Psi'_0 \rangle = \max_{|\Phi'\rangle \in \text{Pro}(\mathcal{H}^{\otimes k})} |\langle \Phi' | \Psi'_0 \rangle|. \tag{8}$$

We now impose the assumption of the induction, that there exists a state  $|v_0\rangle \geq 0$  such that  $\langle v_0 | \otimes \langle a_{k+1} | | \Psi'_0 \rangle = \langle a_1 | \otimes \cdots \otimes \langle a_k | | \Psi'_0 \rangle = \max_{|\Phi'\rangle \in \text{Pro}(\mathcal{H}^{\otimes k})} |\langle \Phi' | \Psi'_0 \rangle|$ . From Eq. (7), we derive

$$\langle v_0 | \otimes \langle a_{k+1} | | \Psi'_0 \rangle = \max_{|\Phi\rangle \in \text{Pro}(\mathcal{H}^{\otimes k+1})} |\langle \Phi | \Psi \rangle| \tag{9}$$

Now, we define a finite sequence of non-negative states  $\{|C_p^{(0)}\}_{p=1}^{k+1}$  as

$$|C_p^{(0)}\rangle := |v_0\rangle \quad \text{for } 1 \leq p \leq k,$$

$$|C_{k+1}^{(0)}\rangle := |a_{k+1}\rangle.$$

By utilizing procedure detailed below, we will use this definition as a starting point for the construction of an infinite sequence of sets of non-negative states  $\{|C_p^{(i)}\}_{p=1}^{k+1}\}_{i=0}^{\infty}$  satisfying

$$\langle C_1^{(i)} | \otimes \cdots \otimes \langle C_{k+1}^{(i)} | | \Psi \rangle = \max_{|\Phi\rangle \in \text{Pro}(\mathcal{H}^{\otimes k+1})} |\langle \Phi | \Psi \rangle| \tag{10}$$

for all non-negative integers  $i$ . We note that because of the permutation symmetry of  $|\Psi\rangle$ , there is no significance to the order imposed by  $p$ .  $\{|C_p^{(i+1)}\}_{p=1}^{k+1}$  is defined from  $\{|C_p^{(i)}\}_{p=1}^{k+1}$  as follows. We choose a couple of states  $\{|C_\alpha^{(i)}\}, |C_\beta^{(i)}\}$  from  $\{|C_p^{(i)}\}_{p=1}^{k+1}$  such that their inner product  $\langle C_\alpha^{(i)} | C_\beta^{(i)} \rangle$  is the least among the inner products of all pairs of states selected from  $\{|C_p^{(i)}\}_{p=1}^{k+1}$ . Then,  $|C_\alpha^{(i+1)}\rangle$  and  $|C_\beta^{(i+1)}\rangle$  are defined as

$$|C_\alpha^{(i+1)}\rangle = |C_\beta^{(i+1)}\rangle := \frac{|C_\alpha^{(i)}\rangle + |C_\beta^{(i)}\rangle}{\| |C_\alpha^{(i)}\rangle + |C_\beta^{(i)}\rangle \|}. \tag{11}$$

For all other  $p \neq \alpha, \beta$ , we define  $|C_p^{(i+1)}\rangle$  as  $|C_p^{(i+1)}\rangle = |C_p^{(i)}\rangle$ . We need to show that the set of non-negative states  $\{|C_p^{(i+1)}\}_{p=1}^{k+1}$  defined as above actually satisfies Eq. (10) for all  $i$ . From the permutation symmetry of  $|\Psi\rangle$ , we can set  $\alpha=1$  and  $\beta=2$  without losing any generality. Then, we define a non-negative non-normalized bipartite state  $|\Psi'_i\rangle \in \mathcal{H}^{\otimes 2}$  as  $|\Psi'_i\rangle := I_{\mathcal{H}}^{\otimes 2} \otimes \langle C_3^{(i)} | \otimes \cdots \otimes \langle C_{k+1}^{(i)} | | \Psi \rangle$ . By the same discussion, we used to derive Eq. (8), we can conclude that

$$\langle C_1^{(i)} | \otimes \langle C_2^{(i)} | | \Psi'_i \rangle = \max_{|\Phi\rangle \in \text{Pro}(\mathcal{H}^{\otimes 2})} |\langle \Phi | \Psi'_i \rangle|. \tag{12}$$

By means of Lemma 1, we obtain  $\langle C_1^{(i+1)} | \otimes \langle C_2^{(i+1)} | | \Psi'_i \rangle = \langle C_1^{(i)} | \otimes \langle C_2^{(i)} | | \Psi'_i \rangle = \max_{|\Phi\rangle \in \text{Pro}(\mathcal{H}^{\otimes 2})} |\langle \Phi | \Psi'_i \rangle|$ . This means that  $\{|C_p^{(i+1)}\}_{p=1}^{k+1}$  satisfies Eq. (10) for all  $i$ . We are now at a stage where we have a symmetrization procedure that produces a sequence of product states that

all have the maximal inner product with the entangled state  $|\Psi\rangle$ . We must, however, show that this sequence of product states converges to a symmetric product state. This is the step that we now address.

Suppose  $\text{ang}(|u\rangle, |v\rangle)$  is the angle between two single-party states  $|u\rangle$  and  $|v\rangle$ . We define  $\theta_i$  as  $\theta_i := \max_{1 \leq p, q \leq k+1} \arg(|C_p^{(i)}\rangle, |C_q^{(i)}\rangle)$ . Then, by the definition of  $\{|C_p^{(i)}\rangle\}_{p=1}^{k+1}$ , we can easily see  $\theta_{i+1} \leq \theta_i$ . Moreover, we can prove  $\lim_{i \rightarrow \infty} \theta_i = 0$  as follows.

Suppose  $\{|C_p^{(i)}\rangle\}_{p=1}^{k+1}$  satisfies  $|C_p^{(i)}\rangle = |C_q^{(i)}\rangle = |u\rangle$  for all  $1 \leq p, q \leq \xi$ , and  $|C_p^{(i)}\rangle = |C_q^{(i)}\rangle = |v\rangle$  for all  $\xi+1 \leq p, q \leq k+1$ . Without loss of generality, we can assume  $\xi \geq (k+1)/2$ . Defining  $\eta$  as  $\eta := k+1-\xi$ , we can easily see that  $\{|C_p^{(i+\eta)}\rangle\}_{p=1}^{k+1}$  satisfies  $|C_p^{(i+\eta)}\rangle = |C_q^{(i+\eta)}\rangle = |u\rangle$  for all  $1 \leq p, q \leq \xi-\eta$ , and  $|C_p^{(i+\eta)}\rangle = |C_q^{(i+\eta)}\rangle = |u\rangle + |v\rangle / \sqrt{\| |u\rangle + |v\rangle \|^2}$  for all  $\xi-\eta+1 \leq p, q \leq k+1$ . Hence, if  $k+1$  is an even number and if  $\xi = (k+1)/2$ , then  $\theta_{i+\eta} = 0$ . Otherwise,  $\theta_{i+\eta} = \theta_i/2$ . Therefore, if  $\{|C_p^{(i)}\rangle\}_{p=1}^{k+1}$  satisfies  $|C_p^{(i)}\rangle = |C_q^{(i)}\rangle = |u\rangle$  for all  $1 \leq p, q \leq \xi$ , and  $|C_p^{(i)}\rangle = |C_q^{(i)}\rangle = |v\rangle$  for all  $\xi+1 \leq p, q \leq k+1$ , then,  $\theta_{i+f} \leq \theta_i/2$ , where  $f$  is the largest integer smaller than  $k+1/2$ . Since  $\{|C_p^{(0)}\rangle\}_{p=1}^{k+1}$  actually satisfies the above condition, we derive  $\theta_{i,f} \leq \theta_0/2^h$  for all positive integers  $h$ . Thus, we can conclude  $\lim_{i \rightarrow \infty} \theta_i = 0$ .

A sequence of non-negative states  $\{|C_p^{(i)}\rangle\}_{i=0}^{\infty}$  hence converges to the same non-negative state  $|C^\infty\rangle := \lim_{i \rightarrow \infty} |C_p^{(i)}\rangle$  without depending on  $p$ . Since Eq. (10) is valid for all non-negative integers  $i$ , by means of the continuity of the inner product, we obtain  $\langle C^\infty |^{\otimes k+1} | \Psi \rangle = \max_{|\Phi\rangle \in \text{Pro}(\mathcal{H}^{\otimes k+1})} |\langle \Phi | \Psi \rangle|$ . That is, the statement is valid for  $n=k+1$ . Therefore, by induction with respect of  $n$ , we have proved the statement of Theorem 1. ■

Here, we add one remark concerning the necessity of our Lemma 1. Indeed, from the PF theorem (Ref. 27, Theorem 8.3.1), we can immediately show the existence of a non-negative  $w_0$ , which satisfies Eq. (3). However, this is not enough to prove the theorem for the following reasons: for fixed  $\alpha \neq \beta$ , the PF theorem does enable symmetrization of the closest product state so that a new closest product state has the form of  $|C_\alpha^{(i+1)}\rangle = |C_\beta^{(i+1)}\rangle$ . However, since the symmetrization on a pair of particles  $\alpha$  and  $\beta$ , in principle, break the symmetry previously established among other pairs of particles, simple application of the PF theorem cannot conclude that there exists a sequence of closest product states converging to a completely symmetric state. Therefore, in order to demonstrate the convergence, we really need a relation between  $\{|C_\alpha^{(i)}\rangle, |C_\beta^{(i)}\rangle\}$  and  $\{|C_\alpha^{(i+1)}\rangle, |C_\beta^{(i+1)}\rangle\}$ , as in Lemma 1. Therefore, we need Lemma 1 to symmetrize a closest product state and Lemma 1 is essential for our proof.

#### IV. DISCUSSION AND CONCLUSIONS

In last part of this paper, we give several comments on the theorem. A stronger (and still unproven) version of the theorem could still be valid without the assumption of the “non-negativity” of the symmetric state  $|\Psi\rangle$ . As we have mentioned in the beginning of the paper, this stronger conjecture first appeared in the paper of Wei *et al.*;<sup>2</sup> they used this conjecture in order to propose an analytical formula for the geometric measure of entanglement for GHZ and W states. While we have not been able to prove the stronger version, the proof presented here of Theorem 1 applies to W and GHZ states, as they can be chosen to be non-negative in the sense that we require.

In fact, all specific instances of the geometric measure calculated in our previous paper<sup>12</sup> concern such “non-negative” states, and so the weaker version of the conjecture, Theorem 1, proved here is sufficient for those cases (see Sec. III.B of Ref. 12).

This weaker version of the conjecture proven above can be useful for calculations of the geometric measure of entanglement of various multipartite systems. For instance, recently several researches investigated possible connections between the behavior of the geometric measure of entanglement and existence of quantum phase transition in natural physical systems.<sup>4,5</sup> However, it is generally impossible to calculate a value of the geometric measure of entanglement for such large systems because the definition of the geometric measure involves a large optimization problem over all product states. Theorem 1 above provides a way to reduce the size of the optimization problem in those cases where the state is known to be symmetric and also non-



negative. Actually, in almost all the calculations to date of the geometric measure for ground states, the possibility of this type of reduction has been assumed. This paper gives a mathematically rigorous proof of this type of reduction for a restricted subset of pure states (“*the set all non-negative states*”) on a symmetric Hilbert space.

Finally, we mention the possibility of the extension of this theorem for a larger subset of symmetric states. The logic in the proof of Theorem 1 strongly depends on the reduction in the optimization problem described by Eq. (4). However, a similar reduction is no longer trivial for a state  $|\Psi\rangle$  having negative amplitudes. Moreover, when the state  $|\Psi\rangle$  has complex amplitudes, Lemma 1 is not valid (although it is, of course, clear that applying a local unitary  $U \otimes U \otimes \dots$  to a “non-negative” state gives a “nonpositive” state for which the conjecture is true). Hence, supplying either a proof or a counterexample to the original stronger statement of Theorem 1 (Ref. 12) is an interesting open problem.

## ACKNOWLEDGMENTS

We are very grateful to Robert Hübener and Otfried Gühne for pointing out the error in the argument presented in our original paper (Ref. 12). We also thank Tzu-Shieh Wei and Simone Severini for discussions. We are obliged to the editor for pointing out a very simple proof of Lemma 1. M.O. was supported by the EPSRC-GB Grant EP/C546237/1, and the European Union Integrated Project Qubit Applications (QAP). M.M. acknowledges support from JST and Nano-Quine. M.H. was supported by a Grant-in-Aid for Scientific Research in the Priority Area “Deepening and Expansion of Statistical Mechanical Informatics (DEX-SMI),” No. 18079014 and a MEXT Grant-in-Aid for Young Scientists (A) No. 20686026. S.V. acknowledges support from EU-STREP “Corner” and Quisco.

- <sup>1</sup>A. Shimony, *Ann. N.Y. Acad. Sci.* **755**, 675 (1995).
- <sup>2</sup>T.-C. Wei and P. M. Goldbart, *Phys. Rev. A* **68**, 042307 (2003).
- <sup>3</sup>D. Markham, J. Anders, V. Vedral, M. Murao, and A. Miyake, *Europhys. Lett.* **81**, 40006 (2008).
- <sup>4</sup>T.-C. Wei, D. Das, S. Mukhopadhyay, S. Vishveshwara, and P. M. Goldbart, *Phys. Rev. A* **71**, 060305 (2005).
- <sup>5</sup>Y. Nakata, D. Markham, and M. Murao, *Phys. Rev. A* **79**, 042313 (2009).
- <sup>6</sup>M. Hayashi, D. Markham, M. Murao, M. Owari, and S. Virmani, *Phys. Rev. Lett.* **96**, 040501 (2006).
- <sup>7</sup>D. Gross, S. Flammia, and J. Eisert, *Phys. Rev. Lett.* **102**, 190501 (2009).
- <sup>8</sup>C. E. Mora, M. Piani, A. Miyake, M. Van den Nest, W. Dür, and H. J. Briegel, *Phys. Rev. A* **80**, 033607 (2009).
- <sup>9</sup>R. Orús, S. Dusuel, and J. Vidal, *Phys. Rev. Lett.* **101**, 025701 (2008).
- <sup>10</sup>Q.-Q. Shi, R. Orus, J. O. Fjærstad, and H.-Q. Zhou, e-print arXiv:0901.2863v1 (2009).
- <sup>11</sup>O. Gühne and G. Toth, *Phys. Rep.* **474**, 1 (2009).
- <sup>12</sup>M. Hayashi, D. Markham, M. Murao, M. Owari, and S. Virmani, *Phys. Rev. A* **77**, 012104 (2008).
- <sup>13</sup>R. F. Werner and A. S. Holevo, *J. Math. Phys.* **43**, 4353 (2002).
- <sup>14</sup>A. Defant and K. Floret, *Tensor Norms and Operator Ideals* (North-Holland, Amsterdam, 1992).
- <sup>15</sup>D. Pérez-García, M. M. Wolf, C. Palazuelos, I. Villanueva, and M. Junge, *Commun. Math. Phys.* **279**, 455 (2008).
- <sup>16</sup>L. De Lathauwer, B. D. Moor, and J. Vandewalle, *SIAM J. Matrix Anal. Appl.* **21**, 1324 (2000).
- <sup>17</sup>T. Zhang and G. H. Golub, *SIAM J. Matrix Anal. Appl.* **23**, 534 (2001).
- <sup>18</sup>E. Kofidis and P. A. Regalia, *SIAM J. Matrix Anal. Appl.* **23**, 863 (2002).
- <sup>19</sup>H. Wang and N. Ahuja, IEEE International Conference on Pattern Recognition (ICPR), 2004 (unpublished).
- <sup>20</sup>G. Ni and Y. Wang, *Math. Comput. Modell.* **46**, 1345 (2007).
- <sup>21</sup>V. de Silva and L. H. Lim, *SIAM J. Matrix Anal. Appl.* **30**, 1084 (2008).
- <sup>22</sup>T.-C. Wei, M. Ericsson, P. M. Goldbart, and W. J. Munro, *Quantum Inf. Comput.* **4**, 252 (2004).
- <sup>23</sup>D. Markham, A. Miyake, and S. Virmani, *New J. Phys.* **9**, 194 (2007).
- <sup>24</sup>K. G. H. Vollbrecht and R. F. Werner, *Phys. Rev. A* **64**, 062307 (2001).
- <sup>25</sup>For a review of entanglement measures see M. B. Plenio and S. Virmani, *Quantum Inf. Comput.* **7**, 1 (2007); R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki *Rev. Mod. Phys.* **81**, 865 (2009).
- <sup>26</sup>T.-C. Wei and S. Severini, e-print arXiv:0905.0012v1.
- <sup>27</sup>R. A. Horn and C. R. Johnson, *Matrix Analysis* (Cambridge University Press, Cambridge, 1990).